

Hypertension (High Blood Pressure) - Comprehensive Guide

Introduction

Hypertension is a chronic condition in which blood pressure remains consistently high. It increases the risk of heart disease, stroke, kidney disease, and other complications. It is often called the silent killer because many people have no symptoms.

Hypertension is a chronic condition in which blood pressure remains consistently high. It increases the risk of heart disease, stroke, kidney disease, and other complications. It is often called the silent killer because many people have no symptoms.

Hypertension is a chronic condition in which blood pressure remains consistently high. It increases the risk of heart disease, stroke, kidney disease, and other complications. It is often called the silent killer because many people have no symptoms.

Hypertension is a chronic condition in which blood pressure remains consistently high. It increases the risk of heart disease, stroke, kidney disease, and other complications. It is often called the silent killer because many people have no symptoms.

Blood Pressure

Blood pressure has two values: systolic and diastolic. Normal is below 120/80 mmHg. Elevated and high values require monitoring and treatment.

Blood pressure has two values: systolic and diastolic. Normal is below 120/80 mmHg. Elevated and high values require monitoring and treatment.

Blood pressure has two values: systolic and diastolic. Normal is below 120/80 mmHg. Elevated and high values require monitoring and treatment.

Blood pressure has two values: systolic and diastolic. Normal is below 120/80 mmHg. Elevated and high values require monitoring and treatment.

Causes

Primary hypertension develops gradually. Secondary hypertension can be caused by kidney disease, hormone disorders, sleep apnea, or medications.

Primary hypertension develops gradually. Secondary hypertension can be caused by kidney disease, hormone disorders, sleep apnea, or medications.

Primary hypertension develops gradually. Secondary hypertension can be caused by kidney disease, hormone disorders, sleep apnea, or medications.

Primary hypertension develops gradually. Secondary hypertension can be caused by kidney disease, hormone disorders, sleep apnea, or medications.

Risk Factors

Age, obesity, family history, smoking, alcohol, stress, diabetes, kidney disease, lack of exercise, and excessive salt intake.

Age, obesity, family history, smoking, alcohol, stress, diabetes, kidney disease, lack of exercise, and excessive salt intake.

Age, obesity, family history, smoking, alcohol, stress, diabetes, kidney disease, lack of exercise, and excessive salt intake.

Age, obesity, family history, smoking, alcohol, stress, diabetes, kidney disease, lack of exercise, and excessive salt intake.

Symptoms

Most people have no symptoms. Severe hypertension may cause headache, blurred vision, chest pain, dizziness, nosebleeds, and shortness of breath.

Most people have no symptoms. Severe hypertension may cause headache, blurred vision, chest pain, dizziness, nosebleeds, and shortness of breath.

Most people have no symptoms. Severe hypertension may cause headache, blurred vision, chest pain, dizziness, nosebleeds, and shortness of breath.

Most people have no symptoms. Severe hypertension may cause headache, blurred vision, chest pain, dizziness, nosebleeds, and shortness of breath.

Diagnosis

Doctors diagnose hypertension using repeated blood pressure measurements. Blood tests, urine tests, ECG, and kidney function tests may also be recommended.

Doctors diagnose hypertension using repeated blood pressure measurements. Blood tests, urine tests, ECG, and kidney function tests may also be recommended.

Doctors diagnose hypertension using repeated blood pressure measurements. Blood tests, urine tests, ECG, and kidney function tests may also be recommended.

Doctors diagnose hypertension using repeated blood pressure measurements. Blood tests, urine tests, ECG, and kidney function tests may also be recommended.

Treatment

Treatment includes reducing salt intake, exercising regularly, maintaining healthy weight, avoiding smoking, limiting alcohol, and taking prescribed medicines.

Treatment includes reducing salt intake, exercising regularly, maintaining healthy weight, avoiding smoking, limiting alcohol, and taking prescribed medicines.

Treatment includes reducing salt intake, exercising regularly, maintaining healthy weight, avoiding smoking, limiting alcohol, and taking prescribed medicines.

Treatment includes reducing salt intake, exercising regularly, maintaining healthy weight, avoiding smoking, limiting alcohol, and taking prescribed medicines.

Complications

Untreated hypertension can lead to heart attack, stroke, kidney failure, eye damage, heart failure, and vascular disease.

Untreated hypertension can lead to heart attack, stroke, kidney failure, eye damage, heart failure, and vascular disease.

Untreated hypertension can lead to heart attack, stroke, kidney failure, eye damage, heart failure, and vascular disease.

Untreated hypertension can lead to heart attack, stroke, kidney failure, eye damage, heart failure, and vascular disease.

Prevention

Healthy diet, regular exercise, stress reduction, weight control, and routine health checkups reduce the risk.

Healthy diet, regular exercise, stress reduction, weight control, and routine health checkups reduce the risk.

Healthy diet, regular exercise, stress reduction, weight control, and routine health checkups reduce the risk.

Healthy diet, regular exercise, stress reduction, weight control, and routine health checkups reduce the risk.

Summary

Early diagnosis and consistent treatment greatly reduce complications and improve quality of life.

Early diagnosis and consistent treatment greatly reduce complications and improve quality of life.

Early diagnosis and consistent treatment greatly reduce complications and improve quality of life.

Early diagnosis and consistent treatment greatly reduce complications and improve quality of life.