

# Topological simplification problems

Ulrich Bauer

IST Austria

July 2, 2012


ATMCS 5, Edinburgh

Joint work with:

Carsten Lange, Max Wardetzky


Nina Amenta, Dominique Attali, Olivier Devillers, Marc Glisse, André Lieutier

# Topological denoising of a function




Noise (even small) can create lots of critical points

# Topological denoising of a function



Noise (even small) can create lots of critical points

# Topological denoising of a function




Noise (even small) can create lots of critical points

Topological denoising by simplification of critical set:

- ▶ removing critical points caused by noise

# Topological denoising of a function



Noise (even small) can create lots of critical points

Topological denoising by simplification of critical set:


- ▶ removing critical points caused by noise

## Problem



Given a function  $f$  and  $\delta > 0$ , find a function  $f_\delta$  that:

- ▶ minimizes number of critical points
- ▶ stays close to input function:  $\|f_\delta - f\|_\infty \leq \delta$

# Persistence diagrams [Cohen-Steiner et al., 2005]



# Persistence diagrams [Cohen-Steiner et al., 2005]




# Stability of persistence diagrams

Theorem (Cohen-Steiner et al., 2005)

Let  $\|f - g\|_\infty \leq \delta$ .

- ▶ The persistence pairs of  $f$  that have persistence  $> 2\delta$  can be mapped injectively to the persistence pairs of  $g$ .




# Stability of persistence diagrams

Theorem (Cohen-Steiner et al., 2005)

Let  $\|f - g\|_\infty \leq \delta$ .

- ▶ The persistence pairs of  $f$  that have persistence  $> 2\delta$  can be mapped injectively to the persistence pairs of  $g$ .
- ▶ Corresponding points  $p_f, p_g$  in the persistence diagrams have distance  $\|p_f - p_g\|_\infty \leq \delta$ .



# A bound on number of critical points

## Corollary

*Let  $f$  be a discrete Morse function and let  $\delta > 0$ .*

*Then for every function  $f_\delta$  with  $\|f_\delta - f\|_\infty \leq \delta$  we have:*

$$\begin{aligned} & \# \text{ critical points of } f_\delta \\ & \geq \# \text{ critical points of } f \text{ with persistence} > 2\delta. \end{aligned}$$

# A bound on number of critical points

## Corollary

*Let  $f$  be a discrete Morse function and let  $\delta > 0$ .*

*Then for every function  $f_\delta$  with  $\|f_\delta - f\|_\infty \leq \delta$  we have:*

$$\begin{aligned} & \# \text{ critical points of } f_\delta \\ & \geq \# \text{ critical points of } f \text{ with persistence} > 2\delta. \end{aligned}$$

## Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:

- ▶ remove all persistence pairs of  $f$  with persistence  $\leq 2\delta$

## Side-effects of elimination


Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:

- ▶ remove all persistence pairs of  $f$  with persistence  $\leq 2\delta$
- ▶ leave all other persistence pairs unmodified

# Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:


- ▶ remove all persistence pairs of  $f$  with persistence  $\leq 2\delta$
- ▶ leave all other persistence pairs unmodified



# Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:


- remove all persistence pairs of  $f$  with persistence  $\leq 2\delta$
- leave all other persistence pairs unmodified



# Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:


- remove all persistence pairs of  $f$  with persistence  $\leq 2\delta$
- leave all other persistence pairs unmodified



# Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:

- remove all persistence pairs of  $f$  with persistence  $\leq 2\delta$
- leave all other persistence pairs unmodified



For an optimal solution, we must allow the critical values to change!

## Interlude: discrete Morse theory

Since Vudit Nanda's talk today was canceled...

## Interlude: discrete Morse theory

What you need to know about discrete Morse theory for this talk:

## Interlude: discrete Morse theory

What you need to know about discrete Morse theory for this talk:

- ▶ it's a discrete version of Morse theory for cell complexes

## Interlude: discrete Morse theory

What you need to know about discrete Morse theory for this talk:

- ▶ it's a discrete version of Morse theory for cell complexes
- ▶ there are discrete notions of gradient vector fields and critical points

## Interlude: discrete Morse theory

What you need to know about discrete Morse theory for this talk:

- ▶ it's a discrete version of Morse theory for cell complexes
- ▶ there are discrete notions of gradient vector fields and critical points
- ▶ Morse functions are generic, nondegenerate functions (isolated critical points)

## Interlude: discrete Morse theory

What you need to know about discrete Morse theory for this talk:


- ▶ it's a discrete version of Morse theory for cell complexes
- ▶ there are discrete notions of gradient vector fields and critical points
- ▶ Morse functions are generic, nondegenerate functions (isolated critical points)
- ▶ discrete vector field: set of inequality constraints on function values

# Interlude: discrete Morse theory


What you need to know about discrete Morse theory for this talk:

- ▶ it's a discrete version of Morse theory for cell complexes
- ▶ there are discrete notions of gradient vector fields and critical points
- ▶ Morse functions are generic, nondegenerate functions (isolated critical points)
- ▶ discrete vector field: set of inequality constraints on function values
- ▶ homotopy type of sublevel sets changes only at critical values

## Canceling critical points of a gradient field



## Canceling critical points of a gradient field



# Persistence pairs and Morse cancellations

Theorem (B., Lange, Wardetzky, 2011)

*Let  $f$  be an excellent discrete Morse function on a surface (distinct critical values) with gradient field  $V$ . Any persistence pair  $(\sigma, \tau)$  can be canceled in  $V$  after all persistence pairs  $(\tilde{\sigma}, \tilde{\tau})$  with*

$$f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$$

*have been canceled.*

# Persistence pairs and Morse cancellations

Theorem (B., Lange, Wardetzky, 2011)

*Let  $f$  be an excellent discrete Morse function on a surface (distinct critical values) with gradient field  $V$ . Any persistence pair  $(\sigma, \tau)$  can be canceled in  $V$  after all persistence pairs  $(\tilde{\sigma}, \tilde{\tau})$  with*

$$f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$$

*have been canceled.*

Corollary

*On a surface, it is possible to cancel just the persistence pairs with persistence  $\leq 2\delta$  (without canceling the other pairs).*

# Persistence pairs and Morse cancellations

Theorem (B., Lange, Wardetzky, 2011)

*Let  $f$  be an excellent discrete Morse function on a surface (distinct critical values) with gradient field  $V$ . Any persistence pair  $(\sigma, \tau)$  can be canceled in  $V$  after all persistence pairs  $(\tilde{\sigma}, \tilde{\tau})$  with*

$$f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$$


*have been canceled.*

Corollary


*On a surface, it is possible to cancel just the persistence pairs with persistence  $\leq 2\delta$  (without canceling the other pairs).*

Does not hold in higher dimensions!


# Cancelling critical points of a function




# Cancelling critical points of a function




# Cancelling critical points of a function




# Cancelling critical points of a function




# Cancelling critical points of a function




# Cancelling critical points of a function



# Cancelling critical points of a function




# Degenerate functions



After cancelation, function is no longer Morse


# Degenerate functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones


# Degenerate functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones
  - ▶  $f$  is *consistent* with  $V$ : if  $\sigma$  is facet of  $\tau$ ,  
 $(\sigma, \tau) \notin V \Rightarrow f(\sigma) \leq f(\tau)$   
 $(\sigma, \tau) \in V \Rightarrow f(\sigma) \geq f(\tau)$


# Degenerate functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones
  - ▶  $f$  is *consistent* with  $V$ : if  $\sigma$  is facet of  $\tau$ ,  
 $(\sigma, \tau) \notin V \Rightarrow f(\sigma) \leq f(\tau)$   
 $(\sigma, \tau) \in V \Rightarrow f(\sigma) \geq f(\tau)$
  - ▶ closure of the set of discrete Morse functions


# Degenerate functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones
  - ▶  $f$  is *consistent* with  $V$ : if  $\sigma$  is facet of  $\tau$ ,  
 $(\sigma, \tau) \notin V \Rightarrow f(\sigma) \leq f(\tau)$   
 $(\sigma, \tau) \in V \Rightarrow f(\sigma) \geq f(\tau)$
  - ▶ closure of the set of discrete Morse functions
- ▶ Gradient vector field is no longer unique in general

# Degenerate functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones
  - ▶  $f$  is *consistent* with  $V$ : if  $\sigma$  is facet of  $\tau$ ,  
 $(\sigma, \tau) \notin V \Rightarrow f(\sigma) \leq f(\tau)$   
 $(\sigma, \tau) \in V \Rightarrow f(\sigma) \geq f(\tau)$
  - ▶ closure of the set of discrete Morse functions
- ▶ Gradient vector field is no longer unique in general

# Symbolic perturbation

- ▶ Use infinitesimal perturbations to resolve degeneracies

## Symbolic perturbation

- ▶ Use infinitesimal perturbations to resolve degeneracies

Let  $f$  pseudo-Morse,  $g$  excellent Morse (distinct critical values), both consistent with gradient vector field  $V$ .

# Symbolic perturbation

- ▶ Use infinitesimal perturbations to resolve degeneracies

Let  $f$  pseudo-Morse,  $g$  excellent Morse (distinct critical values), both consistent with gradient vector field  $V$ .

- ▶ For any  $\epsilon > 0$ ,  
 $f_\epsilon = f + \epsilon g$  is an excellent Morse function consistent with  $V$

# Symbolic perturbation

- ▶ Use infinitesimal perturbations to resolve degeneracies

Let  $f$  pseudo-Morse,  $g$  excellent Morse (distinct critical values), both consistent with gradient vector field  $V$ .

- ▶ For any  $\epsilon > 0$ ,  
 $f_\epsilon = f + \epsilon g$  is an excellent Morse function consistent with  $V$

Assume additionally that the order induced by  $g$  extends the order induced by  $f$ :  $g(\sigma) < g(\tau) \Rightarrow f(\sigma) \leq f(\tau)$

# Symbolic perturbation

- ▶ Use infinitesimal perturbations to resolve degeneracies

Let  $f$  pseudo-Morse,  $g$  excellent Morse (distinct critical values), both consistent with gradient vector field  $V$ .

- ▶ For any  $\epsilon > 0$ ,  
 $f_\epsilon = f + \epsilon g$  is an excellent Morse function consistent with  $V$

Assume additionally that the order induced by  $g$  extends the order induced by  $f$ :  $g(\sigma) < g(\tau) \Rightarrow f(\sigma) \leq f(\tau)$

- ▶  $f_\epsilon$  induces the same order as  $g$ :  $g(\sigma) < g(\tau) \Leftrightarrow f_\epsilon(\sigma) < f_\epsilon(\tau)$

# Symbolic perturbation

- ▶ Use infinitesimal perturbations to resolve degeneracies

Let  $f$  pseudo-Morse,  $g$  excellent Morse (distinct critical values), both consistent with gradient vector field  $V$ .

- ▶ For any  $\epsilon > 0$ ,  
 $f_\epsilon = f + \epsilon g$  is an excellent Morse function consistent with  $V$

Assume additionally that the order induced by  $g$  extends the order induced by  $f$ :  $g(\sigma) < g(\tau) \Rightarrow f(\sigma) \leq f(\tau)$

- ▶  $f_\epsilon$  induces the same order as  $g$ :  $g(\sigma) < g(\tau) \Leftrightarrow f_\epsilon(\sigma) < f_\epsilon(\tau)$
- ▶ the persistence pairs of  $f_\epsilon$  are the persistence pairs of  $g$

# Symbolic perturbation

- ▶ Use infinitesimal perturbations to resolve degeneracies

Let  $f$  pseudo-Morse,  $g$  excellent Morse (distinct critical values), both consistent with gradient vector field  $V$ .

- ▶ For any  $\epsilon > 0$ ,  
 $f_\epsilon = f + \epsilon g$  is an excellent Morse function consistent with  $V$

Assume additionally that the order induced by  $g$  extends the order induced by  $f$ :  $g(\sigma) < g(\tau) \Rightarrow f(\sigma) \leq f(\tau)$

- ▶  $f_\epsilon$  induces the same order as  $g$ :  $g(\sigma) < g(\tau) \Leftrightarrow f_\epsilon(\sigma) < f_\epsilon(\tau)$
- ▶ the persistence pairs of  $f_\epsilon$  are the persistence pairs of  $g$

Most important statements allow passing to the limit  $\epsilon \rightarrow 0!$

# Optimal topological simplification

Theorem (B., Lange, Wardetzky, 2011)

*Let  $f$  be a pseudo-Morse function on a surface and let  $\delta > 0$ .*

# Optimal topological simplification

Theorem (B., Lange, Wardetzky, 2011)

*Let  $f$  be a pseudo-Morse function on a surface and let  $\delta > 0$ .*

*Let  $f_\delta$  be obtained from  $f$  by canceling all persistence pairs with persistence  $\leq 2\delta$ .*

# Optimal topological simplification

Theorem (B., Lange, Wardetzky, 2011)

*Let  $f$  be a pseudo-Morse function on a surface and let  $\delta > 0$ .*

*Let  $f_\delta$  be obtained from  $f$  by canceling all persistence pairs with persistence  $\leq 2\delta$ . Then*

$$\|f_\delta - f\|_\infty \leq \delta.$$

# Optimal topological simplification

Theorem (B., Lange, Wardetzky, 2011)

*Let  $f$  be a pseudo-Morse function on a surface and let  $\delta > 0$ .*

*Let  $f_\delta$  be obtained from  $f$  by canceling all persistence pairs with persistence  $\leq 2\delta$ . Then*

$$\|f_\delta - f\|_\infty \leq \delta.$$

*I.e.,  $f_\delta$  achieves the lower bound on the number of critical points.*

# Optimal topological simplification

Theorem (B., Lange, Wardetzky, 2011)

Let  $f$  be a pseudo-Morse function on a surface and let  $\delta > 0$ .

Let  $f_\delta$  be obtained from  $f$  by canceling all persistence pairs with persistence  $\leq 2\delta$ . Then

$$\|f_\delta - f\|_\infty \leq \delta.$$

I.e.,  $f_\delta$  achieves the lower bound on the number of critical points.

- ▶ Does not hold for non-manifold 2-complexes or higher dimensions (in general, simplification is NP-hard)

# Optimal topological simplification

Theorem (B., Lange, Wardetzky, 2011)

Let  $f$  be a pseudo-Morse function on a surface and let  $\delta > 0$ .

Let  $f_\delta$  be obtained from  $f$  by canceling all persistence pairs with persistence  $\leq 2\delta$ . Then

$$\|f_\delta - f\|_\infty \leq \delta.$$

I.e.,  $f_\delta$  achieves the lower bound on the number of critical points.

- ▶ Does not hold for non-manifold 2-complexes or higher dimensions (in general, simplification is NP-hard)
- ▶ Solution can be found in linear time after computation of persistence pairs

## Combination with energy methods

Recall: simplified vector field  $V_\delta$  imposes inequalities on simplified function consistent with  $V_\delta$

## Combination with energy methods

Recall: simplified vector field  $V_\delta$  imposes inequalities on simplified function consistent with  $V_\delta$

$\|f_\delta - f\|_\infty \leq \delta$ : another set of linear inequalities

## Combination with energy methods

Recall: simplified vector field  $V_\delta$  imposes inequalities on simplified function consistent with  $V_\delta$

$\|f_\delta - f\|_\infty \leq \delta$ : another set of linear inequalities

- ▶ defines convex polytope of solutions:  
any function  $g$  consistent with  $V_\delta$  and  $\|g - f\|_\infty \leq \delta$  is a solution


# Combination with energy methods

Recall: simplified vector field  $V_\delta$  imposes inequalities on simplified function consistent with  $V_\delta$


$\|f_\delta - f\|_\infty \leq \delta$ : another set of linear inequalities

- ▶ defines convex polytope of solutions:  
any function  $g$  consistent with  $V_\delta$  and  $\|g - f\|_\infty \leq \delta$  is a solution
- ▶ find the “best” solution using your favorite energy functional






# Removing local extrema from 3D data



In 3D:

- ▶ simplifying critical points is hard

# Removing local extrema from 3D data



In 3D:

- ▶ simplifying critical points is hard
- ▶ simplifying only extrema is easy

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta_*(G_{\leq t})$ .*

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .
- ▶ For any  $X$  with  $L \subset X \subset K$ , there is  $g$  with  $G_{\leq t} = X$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .
- ▶ For any  $X$  with  $L \subset X \subset K$ , there is  $g$  with  $G_{\leq t} = X$ .

Thus we are looking for  $X$  with  $L \subset X \subset K$  minimizing  $\beta_*(X)$ .

# Homological factorization

## Problem

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

# Homological factorization

## Problem

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological factorization* of  $(K, L)$ .

# Homological factorization

## Problem

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological factorization* of  $(K, L)$ .

- ▶ If  $L \subset X \subset K$  then  $\beta_*(X) \geq \text{rank } H_*(L \hookrightarrow K)$

# Homological factorization

## Problem

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological factorization* of  $(K, L)$ .

- ▶ If  $L \subset X \subset K$  then  $\beta_*(X) \geq \text{rank } H_*(L \hookrightarrow K)$
- ▶ in  $\mathbb{R}^3$ : equality iff  $X$  is a homological factorization


# Homological factorization

## Problem

Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.

Such an  $X$  is called a *homological factorization* of  $(K, L)$ .

- ▶ If  $L \subset X \subset K$  then  $\beta_*(X) \geq \text{rank } H_*(L \hookrightarrow K)$
- ▶ in  $\mathbb{R}^3$ : equality iff  $X$  is a homological factorization



homological factorizations  
do not always exist


# Homological factorization

## Problem


Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.

Such an  $X$  is called a *homological factorization* of  $(K, L)$ .

- If  $L \subset X \subset K$  then  $\beta_*(X) \geq \text{rank } H_*(L \hookrightarrow K)$
- in  $\mathbb{R}^3$ : equality iff  $X$  is a homological factorization



homological factorizations  
do not always exist



a homological factorization may exist,  
but not as a subcomplex of  $K$

# Homological factorizability in $\mathbb{R}^3$ is NP-complete

Theorem (Attali, Lieutier; 2010)

*Deciding whether  $(K, L)$  has a homological factorization as a subcomplex is NP-complete.*

## Homological factorizability in $\mathbb{R}^3$ is NP-complete

Theorem (Attali, Lieutier; 2010)

*Deciding whether  $(K, L)$  has a homological factorization as a subcomplex is NP-complete.*

Theorem (A., A., B., D., G., L.; 2012)

*This holds even for complexes  $K$  embeddable in  $\mathbb{R}^3$ .*

# Homological factorizability in $\mathbb{R}^3$ is NP-complete

Theorem (Attali, Lieutier; 2010)

*Deciding whether  $(K, L)$  has a homological factorization as a subcomplex is NP-complete.*

Theorem (A., A., B., D., G., L.; 2012)

*This holds even for complexes  $K$  embeddable in  $\mathbb{R}^3$ .*

Corollary

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

# Homological factorizability in $\mathbb{R}^3$ is NP-complete

Theorem (Attali, Lieutier; 2010)

*Deciding whether  $(K, L)$  has a homological factorization as a subcomplex is NP-complete.*

Theorem (A., A., B., D., G., L.; 2012)

*This holds even for complexes  $K$  embeddable in  $\mathbb{R}^3$ .*

## Corollary

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Idea of proof: reduction from 3-SAT

- ▶ given a 3-SAT instance, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$

# Homological factorizability in $\mathbb{R}^3$ is NP-complete

Theorem (Attali, Lieutier; 2010)

*Deciding whether  $(K, L)$  has a homological factorization as a subcomplex is NP-complete.*

Theorem (A., A., B., D., G., L.; 2012)

*This holds even for complexes  $K$  embeddable in  $\mathbb{R}^3$ .*


## Corollary

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Idea of proof: reduction from 3-SAT


- ▶ given a 3-SAT instance, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶  $X$  is homological factorization  $\Leftrightarrow X$  is acyclic,  $L \subset X \subset K$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  can contain only one of the edges  $\text{True}_i, \text{False}_i$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  can contain only one of the edges  $\text{True}_i, \text{False}_i$
- ▶ edges  $\text{True}_i, \text{False}_i$  correspond to truth assignment of variable  $x_i$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$

## Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,
- ▶ blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,
- ▶ blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$
- ▶  $X$  must contain one  
of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$
- ▶  $X$  must contain one  
of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$
- ▶  $X$  must contain one  
of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$
- ▶  $X$  must contain one  
of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,
- ▶ blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$
- ▶  $X$  must contain one  
of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,
- ▶ blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ ,
- ▶ blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget



- ▶ red: contained in  $L$ ,
- ▶ blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$


# Reduction from 3-SAT: the clause gadget



- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$
- ▶  $X$  must contain one  
of the edges  $a, b, c$

- ▶ For every clause (e.g.,  $(x_1 \vee \neg x_2 \vee x_4)$ ):  
identify  $a, b, c$  with edges of variable gadgets  
corresponding to the literals  $(\text{True}_1, \text{False}_2, \text{True}_4)$

# Reduction from 3-SAT: the clause gadget



- ▶ red: contained in  $L$ ,  
blue:  $K \setminus L$
- ▶  $X$  must contain one  
of the edges  $a, b, c$

- ▶ For every clause (e.g.,  $(x_1 \vee \neg x_2 \vee x_4)$ ):  
identify  $a, b, c$  with edges of variable gadgets  
corresponding to the literals  $(\text{True}_1, \text{False}_2, \text{True}_4)$
- ▶ We can transform a homological factorization of  $(K, L)$   
into a satisfying assignment and vice versa

# Simplification of level sets

## Theorem

*Level set simplification in  $\mathbb{R}^3$  is NP-hard.*

# Simplification of level sets

## Theorem

*Level set simplification in  $\mathbb{R}^3$  is NP-hard.*

Idea of proof:

Assume  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ . Let  $g$  be a  $t$ -level set simplification of  $f$  (a function  $g$  minimizing  $\beta_*(G_{=t})$ ).

Then  $G_{\leq t}$  is a homological factorization of  $(K, L)$ , if one exists.

# Simplification of level sets

## Theorem

*Level set simplification in  $\mathbb{R}^3$  is NP-hard.*

Idea of proof:

Assume  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ . Let  $g$  be a  $t$ -level set simplification of  $f$  (a function  $g$  minimizing  $\beta_*(G_{=t})$ ).

Then  $G_{\leq t}$  is a homological factorization of  $(K, L)$ , if one exists.

- ▶  $\beta_*(G_{=t}) = \beta_*(G_{$

# Simplification of level sets

## Theorem

*Level set simplification in  $\mathbb{R}^3$  is NP-hard.*

Idea of proof:

Assume  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ . Let  $g$  be a  $t$ -level set simplification of  $f$  (a function  $g$  minimizing  $\beta_*(G_{=t})$ ).

Then  $G_{\leq t}$  is a homological factorization of  $(K, L)$ , if one exists.

- ▶  $\beta_*(G_{=t}) = \beta_*(G_{$
- ▶  $g$  may be assumed to have regular value  $t$ . Hence  
$$\beta_*(G_{=t}) = 2\beta_*(G_{\leq t})$$

# Simplification of level sets

## Theorem

*Level set simplification in  $\mathbb{R}^3$  is NP-hard.*

Idea of proof:

Assume  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ . Let  $g$  be a  $t$ -level set simplification of  $f$  (a function  $g$  minimizing  $\beta_*(G_{=t})$ ).

Then  $G_{\leq t}$  is a homological factorization of  $(K, L)$ , if one exists.

- ▶  $\beta_*(G_{=t}) = \beta_*(G_{)}) + \beta_*(G_{\leq t})$
- ▶  $g$  may be assumed to have regular value  $t$ . Hence  
$$\beta_*(G_{=t}) = 2\beta_*(G_{\leq t})$$
- ▶ If a homological factorization exists, then the lower bound on  $\beta_*(G_{\leq t})$  can be achieved

# Simplification of level sets

## Theorem

*Level set simplification in  $\mathbb{R}^3$  is NP-hard.*

Idea of proof:

Assume  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ . Let  $g$  be a  $t$ -level set simplification of  $f$  (a function  $g$  minimizing  $\beta_*(G_{=t})$ ).

Then  $G_{\leq t}$  is a homological factorization of  $(K, L)$ , if one exists.

- ▶  $\beta_*(G_{=t}) = \beta_*(G_{)}) + \beta_*(G_{\leq t})$
- ▶  $g$  may be assumed to have regular value  $t$ . Hence  
$$\beta_*(G_{=t}) = 2\beta_*(G_{\leq t})$$
- ▶ If a homological factorization exists, then the lower bound on  $\beta_*(G_{\leq t})$  can be achieved
- ▶ since  $g$  minimizes  $\beta_*(G_{=t}) = 2\beta_*(G_{\leq t})$ , it achieves this bound

# Thanks for your attention!

