

Applied Topology

From Theory to Computation

Ulrich Bauer

TUM

February 5, 2016

Inaugural lecture



\approx



?





\approx




?


Persistent homology




























What is homology?




- *Simplicial complex K :* space built from *simplices* (points, edges, triangles, tetrahedra, ...)

What is homology?




- *Simplicial complex K :* space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$:* formal \mathbb{Z}_2 linear combinations of simplices

What is homology?




- *Simplicial complex K :* space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$:* formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary

What is homology?




- *Simplicial complex K :* space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$:* formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$

What is homology?




- *Simplicial complex K :* space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$:* formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$

What is homology?




- *Simplicial complex* K : space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains* $C_*(K)$: formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$
- *Cycles* $Z_*(K) = \ker \partial_*$ have no boundary

What is homology?




- *Simplicial complex K :* space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$:* formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$
- *Cycles $Z_*(K) = \ker \partial_*$* have no boundary
- *Boundaries $B_*(K) = \text{im } \partial_*$* are always cycles

What is homology?




- *Simplicial complex K :* space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$:* formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$
- *Cycles $Z_*(K) = \ker \partial_*$* have no boundary
- *Boundaries $B_*(K) = \text{im } \partial_*$* are always cycles

What is homology?




- *Simplicial complex K* : space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$* : formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$
- *Cycles $Z_*(K) = \ker \partial_*$* have no boundary
- *Boundaries $B_*(K) = \text{im } \partial_*$* are always cycles

What is homology?




- *Simplicial complex K* : space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$* : formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$
- *Cycles $Z_*(K) = \ker \partial_*$* have no boundary
- *Boundaries $B_*(K) = \text{im } \partial_*$* are always cycles

What is homology?



- *Simplicial complex K* : space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$* : formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$
- *Cycles $Z_*(K) = \ker \partial_*$* have no boundary
- *Boundaries $B_*(K) = \text{im } \partial_*$* are always cycles
- Two cycles are *homologous* if they differ by a boundary

What is homology?




- *Simplicial complex K* : space built from *simplices* (points, edges, triangles, tetrahedra, ...)
- *Chains $C_*(K)$* : formal \mathbb{Z}_2 linear combinations of simplices
- Each simplex has a boundary
- Defines linear *boundary map* $\partial_* : C_*(K) \rightarrow C_*(K)$
- *Cycles $Z_*(K) = \ker \partial_*$* have no boundary
- *Boundaries $B_*(K) = \text{im } \partial_*$* are always cycles
- Two cycles are *homologous* if they differ by a boundary
- The equivalence classes form the *homology group*
$$H_*(K) = Z_*(K)/B_*(K)$$

Why is homology algebraic?

Roughly: $\dim H_*(K)$ is the number of holes

- Why not just count holes then?




Homology is a *functor*:

Why is homology algebraic?

Roughly: $\dim H_*(K)$ is the number of holes

- Why not just count holes then?




Homology is a *functor*:

- An inclusion map $L \hookrightarrow K$ induces a linear map $H_*(L) \hookrightarrow H_*(K)$

Why is homology algebraic?

Roughly: $\dim H_*(K)$ is the number of holes


- Why not just count holes then?





Homology is a *functor*:

- An inclusion map $L \hookrightarrow K$ induces a linear map $H_*(L) \hookrightarrow H_*(K)$
- A “hole” might get mapped to a linear combination of two “holes”



What is persistent homology?





What is persistent homology?





What is persistent homology?





What is persistent homology?





What is persistent homology?



What is persistent homology?



What is persistent homology?



Homology inference & reconstruction

Homology inference

Problem (Homology inference)

Determine the homology $H_(\Omega)$ of a shape $\Omega \subset \mathbb{R}^d$ from a finite sample $P \subset \Omega$.*

Homology inference

Problem (Homology inference)

Determine the homology $H_(\Omega)$ of a shape $\Omega \subset \mathbb{R}^d$ from a finite sample $P \subset \Omega$.*

Problem (Homological reconstruction)

Given a finite sample $P \subset \Omega$, construct a shape X that is geometrically close to Ω and satisfies $H_(X) \cong H_*(\Omega)$.*

Homology inference

Problem (Homology inference)

Determine the homology $H_*(\Omega)$ of a shape $\Omega \subset \mathbb{R}^d$ from a finite sample $P \subset \Omega$.

Problem (Homological reconstruction)

Given a finite sample $P \subset \Omega$, construct a shape X that is geometrically close to Ω and satisfies $H_*(X) \cong H_*(\Omega)$.

- Approximate the shape by a thickening $B_\delta(P)$ covering Ω
- Geometrically close means that the isomorphism is induced by the inclusion $\Omega \hookrightarrow B_\delta(P)$
- Replace $B_\delta(P)$ by an equivalent simplicial complex
 - Čech complex $\text{Cech}_\delta(P)$
 - Delaunay complex $\text{Del}_\delta(P)$

Homology reconstruction using union of balls

It is sometimes possible to recover the homology of Ω this way,
but the assumptions are quite strong:

Homology reconstruction using union of balls

It is sometimes possible to recover the homology of Ω this way, but the assumptions are quite strong:

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$ be such that $M \subseteq P^\delta$ for some $\delta < \sqrt{3/20} \text{reach}(M)$. Then

$$H_*(M) \cong H_*(P^{2\delta}).$$

Homology reconstruction using union of balls

It is sometimes possible to recover the homology of Ω this way, but the assumptions are quite strong:

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$ be such that $M \subseteq P^\delta$ for some $\delta < \sqrt{3/20} \text{reach}(M)$. Then

$$H_*(M) \cong H_*(P^{2\delta}).$$

- $P_\delta = B_\delta(P)$: δ -neighborhood (union of balls) around P .

Homology reconstruction using union of balls

It is sometimes possible to recover the homology of Ω this way, but the assumptions are quite strong:

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$ be such that $M \subseteq P^\delta$ for some $\delta < \sqrt{3/20} \text{reach}(M)$. Then

$$H_*(M) \cong H_*(P^{2\delta}).$$

- $P_\delta = B_\delta(P)$: δ -neighborhood (union of balls) around P .
- All points in \mathbb{R}^d with distance $< \text{reach}(M)$ to M have a unique closest point on M

Homology reconstruction using union of balls


It is sometimes possible to recover the homology of Ω this way, but the assumptions are quite strong:


Theorem (Niyogi, Smale, Weinberger 2006)


Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$ be such that $M \subseteq P^\delta$ for some $\delta < \sqrt{3/20} \text{reach}(M)$. Then


$$H_*(M) \cong H_*(P^{2\delta}).$$


- $P_\delta = B_\delta(P)$: δ -neighborhood (union of balls) around P .
- All points in \mathbb{R}^d with distance $< \text{reach}(M)$ to M have a unique closest point on M
- The isomorphism is induced by the inclusion $M \hookrightarrow P^{2\delta}$.























































Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$

Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$

Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$

Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$

Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$

Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$

Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$

Homology inference using persistent homology

For $Q \subseteq \mathbb{R}^d$:

$Q_\delta = B_\delta(Q)$: δ -neighborhood (union of balls) around Q



Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$ be such that

- $\Omega \subseteq P_\delta$ for some $\delta > 0$ and
- both $H_*(\Omega \hookrightarrow \Omega_\delta)$ and $H_*(\Omega_\delta \hookrightarrow \Omega_{2\delta})$ are isomorphisms.

Then

$$H_*(\Omega) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}).$$



Proof.

$$\begin{array}{ccc} H_*(\Omega) & H_*(\Omega_\delta) & H_*(\Omega_{2\delta}) \\ \searrow & \nearrow & \searrow \\ H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \hookrightarrow & H_*(\Omega_\delta) & \hookrightarrow & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \hookrightarrow & H_*(\Omega_\delta) & \hookrightarrow & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \hookrightarrow & H_*(\Omega_\delta) & \hookrightarrow & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \hookrightarrow & H_*(\Omega_\delta) & \hookrightarrow & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & & H_*(P_\delta) & \longrightarrow & H_*(P_{2\delta}) \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \hookrightarrow & H_*(\Omega_\delta) & \hookrightarrow & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & & H_*(P_\delta) & \longrightarrow & H_*(P_{2\delta}) \\ & & \searrow & \nearrow & \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \hookrightarrow & H_*(\Omega_\delta) & \hookrightarrow & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow \uparrow & \swarrow & \nearrow \\ & & H_*(P_\delta) & \xrightarrow{\cong} & H_*(P_{2\delta}) \\ & & \downarrow & & \downarrow \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \overset{\hookleftarrow \cong \rightarrow}{\longrightarrow} & H_*(\Omega_\delta) & \xhookrightarrow{\quad} & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow \nwarrow & \nearrow & \nearrow \\ & & H_*(P_\delta) & \xrightarrow{\cong} & H_*(P_{2\delta}) \\ & & \searrow & \downarrow & \nearrow \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

Proof.

$$\begin{array}{ccccc} H_*(\Omega) & \xhookleftarrow{\cong} & H_*(\Omega_\delta) & \xhookrightarrow{\quad} & H_*(\Omega_{2\delta}) \\ & \searrow & \nearrow \nwarrow & \nearrow & \nearrow \\ & & H_*(P_\delta) & \xrightarrow{\cong} & H_*(P_{2\delta}) \\ & & \searrow & \downarrow & \nearrow \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

□

Homological realization

This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$


Homological realization

This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$




Homological realization

This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$



Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$


This is not always possible:




Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)

The homological realization problem is NP-hard, even in \mathbb{R}^3 .

Simplification





Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

*Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$
minimizing $\dim H_*(G_{\leq t})$.*

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

*Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$
minimizing $\dim H_*(G_{\leq t})$.*

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$
minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$
minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.
- For any X with $L \subseteq X \subseteq K$, there is g with $G_{\leq t} = X$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$
minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.
- For any X with $L \subseteq X \subseteq K$, there is g with $G_{\leq t} = X$.

Thus we seek X with $L \subseteq X \subseteq K$ minimizing $\dim H_*(X)$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$
minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.


- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.
- For any X with $L \subseteq X \subseteq K$, there is g with $G_{\leq t} = X$.

Thus we seek X with $L \subseteq X \subseteq K$ minimizing $\dim H_*(X)$.


Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)

Sublevel set simplification in \mathbb{R}^3 is NP-hard.

Simplification of functions







Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

*Given a function f and a real number $\delta \geq 0$,
find a function f_δ subject to $\|f_\delta - f\|_\infty \leq \delta$
with minimal number of critical points.*

Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

*Given a function f and a real number $\delta \geq 0$,
find a function f_δ subject to $\|f_\delta - f\|_\infty \leq \delta$
with minimal number of critical points.*

(Discrete) Morse theory:

- Relates critical points to homology of sublevel set

Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

*Given a function f and a real number $\delta \geq 0$,
find a function f_δ subject to $\|f_\delta - f\|_\infty \leq \delta$
with minimal number of critical points.*

(Discrete) Morse theory:

- Relates critical points to homology of sublevel set
- Provides method for canceling pairs of critical points

Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

*Given a function f and a real number $\delta \geq 0$,
find a function f_δ subject to $\|f_\delta - f\|_\infty \leq \delta$
with minimal number of critical points.*

(Discrete) Morse theory:

- Relates critical points to homology of sublevel set
- Provides method for canceling pairs of critical points

Persistent homology:

- Relates homology of different sublevel set

Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

*Given a function f and a real number $\delta \geq 0$,
find a function f_δ subject to $\|f_\delta - f\|_\infty \leq \delta$
with minimal number of critical points.*

(Discrete) Morse theory:

- Relates critical points to homology of sublevel set
- Provides method for canceling pairs of critical points

Persistent homology:

- Relates homology of different sublevel set
- Identifies pairs of critical points (birth and death of homological feature) and quantifies their *persistence*

Persistence and discrete Morse theory

By stability of persistence barcodes:

Proposition

The critical points with persistence $> 2\delta$ provide a lower bound on the number of critical points of any function g with $\|g - f\|_\infty \leq \delta$.

Persistence and discrete Morse theory

By stability of persistence barcodes:

Proposition

The critical points with persistence $> 2\delta$ provide a lower bound on the number of critical points of any function g with $\|g - f\|_\infty \leq \delta$.

Theorem (B, Lange, Wardetzky, 2011)

Let f be a function on a surface and let $\delta > 0$.

Construct a function f_δ from f by canceling all persistence pairs with persistence $\leq 2\delta$.

Persistence and discrete Morse theory

By stability of persistence barcodes:

Proposition

The critical points with persistence $> 2\delta$ provide a lower bound on the number of critical points of any function g with $\|g - f\|_\infty \leq \delta$.

Theorem (B, Lange, Wardetzky, 2011)

Let f be a function on a surface and let $\delta > 0$.

Construct a function f_δ from f by canceling all persistence pairs with persistence $\leq 2\delta$. Then f_δ satisfies

$$\|f_\delta - f\|_\infty \leq \delta$$

and achieves the lower bound on the number of critical points.

Persistence and discrete Morse theory

By stability of persistence barcodes:

Proposition

The critical points with persistence $> 2\delta$ provide a lower bound on the number of critical points of any function g with $\|g - f\|_\infty \leq \delta$.

Theorem (B, Lange, Wardetzky, 2011)

Let f be a function on a surface and let $\delta > 0$.

Construct a function f_δ from f by canceling all persistence pairs with persistence $\leq 2\delta$. Then f_δ satisfies

$$\|f_\delta - f\|_\infty \leq \delta$$


and achieves the lower bound on the number of critical points.

- Does not generalize to higher-dimensional manifolds!

Morse theory of geometric complexes

Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)



$\check{\text{C}}$ ech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

- We identify a simplex with its vertex set



$\check{\text{C}}$ ech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

- We identify a simplex with its vertex set



$\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

- We identify a simplex with its vertex set

Two functions on simplices $Q \subseteq X$:

- *$\check{\text{C}}\text{ech function}$* $f_C(Q)$:
radius of smallest enclosing sphere of Q



$\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

- We identify a simplex with its vertex set

Two functions on simplices $Q \subseteq X$:

- *$\check{\text{C}}\text{ech function}$* $f_C(Q)$:
radius of smallest enclosing sphere of Q
- *Delaunay function* $f_D(Q)$:
radius of smallest empty circumsphere of Q



$\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

- We identify a simplex with its vertex set

Two functions on simplices $Q \subseteq X$:

- *$\check{\text{C}}\text{ech function}$* $f_C(Q)$:
radius of smallest enclosing sphere of Q
- *Delaunay function* $f_D(Q)$:
radius of smallest empty circumsphere of Q
 - defined only for the *Delaunay triangulation* $\text{Del}(X)$:
all simplices Q having an empty circumsphere




Čech and Delaunay complexes from functions

For a point set $X \subset \mathbb{R}^d$, define for any radius r :

$\check{\text{C}}\text{ech}$ and Delaunay complexes from functions

For a point set $X \subset \mathbb{R}^d$, define for any radius r :


- $\check{\text{C}}\text{ech complex}: \text{Cech}_r(X) = f_C^{-1}(-\infty, r]$
 - all simplices on X having an enclosing sphere of radius $\leq r$



$\check{\text{C}}\text{ech}$ and Delaunay complexes from functions

For a point set $X \subset \mathbb{R}^d$, define for any radius r :


- *$\check{\text{C}}\text{ech complex}$* : $\text{Cech}_r(X) = f_C^{-1}(-\infty, r]$
 - all simplices on X having an enclosing sphere of radius $\leq r$
- *Delaunay complex (α -shape)*: $\text{Del}_r(X) = f_D^{-1}(-\infty, r]$
 - all simplices having an empty circumsphere of radius $\leq r$



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.




Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex contained in a unique higher-dimensional simplex.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex contained in a unique higher-dimensional simplex.
- We can *collapse* K to a subcomplex L by removing a free face F along with the higher-dimensional simplex.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex contained in a unique higher-dimensional simplex.
- We can *collapse* K to a subcomplex L by removing a free face F along with the higher-dimensional simplex.
- Such a collapse is called *elementary*.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex contained in a unique higher-dimensional simplex.
- We can *collapse* K to a subcomplex L by removing a free face F along with the higher-dimensional simplex.
- Such a collapse is called *elementary*.
- L is homotopy equivalent to K .



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex contained in a unique higher-dimensional simplex.
- We can *collapse* K to a subcomplex L by removing a free face F along with the higher-dimensional simplex.
- Such a collapse is called *elementary*.
- L is homotopy equivalent to K .




If there is a sequence of elementary collapses from K to M , we say that K *collapses* to M (written as $K \searrow M$).

Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014, 2015)

Čech, Delaunay, and Wrap complexes are homotopy equivalent by simplicial collapses:

$$\text{Cech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014, 2015)

Čech, Delaunay, and Wrap complexes are homotopy equivalent by simplicial collapses:

$$\text{Cech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- Wrap complex: commercially used for surface reconstruction (Edelsbrunner 1995, Geomagic Wrap®)

Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014, 2015)

Čech, Delaunay, and Wrap complexes are homotopy equivalent by simplicial collapses:

$$\text{Cech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- Wrap complex: commercially used for surface reconstruction (Edelsbrunner 1995, Geomagic Wrap®)
- All collapses are induced by a single *discrete gradient field*

Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014, 2015)

Čech, Delaunay, and Wrap complexes are homotopy equivalent by simplicial collapses:

$$\text{Cech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- Wrap complex: commercially used for surface reconstruction (Edelsbrunner 1995, Geomagic Wrap®)
- All collapses are induced by a single *discrete gradient field*
- Explicit isomorphisms in homology

Discrete Morse theory

Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a simplicial complex
is a partition of the set of simplices
into singleton sets $\{C\}$ and pairs $\{L, U\}$,
where L is a face of U with codimension 1.



Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the set of simplices into singleton sets $\{C\}$ and pairs $\{L, U\}$, where L is a face of U with codimension 1.




- indicated by an arrow from L to U

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the set of simplices into singleton sets $\{C\}$ and pairs $\{L, U\}$, where L is a face of U with codimension 1.



- indicated by an arrow from L to U


The singletons are called *critical simplices*.

Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes

•
1

Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ partition the set of simplices into a discrete vector field (the *discrete gradient* of f)




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ partition the set of simplices into a discrete vector field (the *discrete gradient* of f)




If $f^{-1}(t) = \{C\}$ is a singleton set, then t is a *critical value*.

Collapses from Morse functions and gradients

Theorem (Forman 1998)

Let f be a discrete Morse function on a simplicial complex K .

*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s .*




Collapses from Morse functions and gradients

Theorem (Forman 1998)

Let f be a discrete Morse function on a simplicial complex K .

*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s .*




Collapses from Morse functions and gradients

Theorem (Forman 1998)

Let f be a discrete Morse function on a simplicial complex K .

If $(s, t]$ contains no critical value of f ,

then K_t collapses to K_s .




Collapses from Morse functions and gradients

Theorem (Forman 1998)

Let f be a discrete Morse function on a simplicial complex K .

*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s .*




Corollary

Let V be a discrete gradient field on a simplicial complex K ,

and let L be a subcomplex of K .

If $K \setminus L$ is a union of pairs of V ,

then K collapses to L .




Collapses from Morse functions and gradients

Theorem (Forman 1998)

Let f be a discrete Morse function on a simplicial complex K .

*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s .*




Corollary

Let V be a discrete gradient field on a simplicial complex K ,

and let L be a subcomplex of K .

If $K \setminus L$ is a union of pairs of V ,

then K collapses to L .




Collapses from Morse functions and gradients

Theorem (Forman 1998)

Let f be a discrete Morse function on a simplicial complex K .


*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s .*



Corollary

*Let V be a discrete gradient field on a simplicial complex K ,
and let L be a subcomplex of K .*

*If $K \setminus L$ is a union of pairs of V ,
then K collapses to L .*



We say that V induces the collapse $K \searrow L$.

Unfortunately...

Neither the Čech nor the Delaunay functions
are discrete Morse functions!

Unfortunately...

Neither the Čech nor the Delaunay functions
are discrete Morse functions!

- Example: two simplices Q, Q' with $f_C(Q) = f_C(Q')$
such that neither is a face of the other:




Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the set of simplices into clusters of simplices containing a simplex L (common face) and contained in a simplex U (common coface):

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the set of simplices into clusters of simplices containing a simplex L (common face) and contained in a simplex U (common coface):

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




- indicated by an arrow from L to U

Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the set of simplices into clusters of simplices containing a simplex L (common face) and contained in a simplex U (common coface):

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$



- indicated by an arrow from L to U


The singletons are called *critical simplices*.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes



The collapsing theorems carry over to generalized discrete Morse functions.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes

•
1

The collapsing theorems carry over to generalized discrete Morse functions.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




The collapsing theorems carry over to generalized discrete Morse functions.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




The collapsing theorems carry over to generalized discrete Morse functions.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




The collapsing theorems carry over to generalized discrete Morse functions.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




The collapsing theorems carry over to generalized discrete Morse functions.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ partition the set of simplices into a generalized discrete vector field (the *discrete gradient* of f)



The collapsing theorems carry over to generalized discrete Morse functions.

Čech and Delaunay intervals

Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)


*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*


*The critical simplices are the centered Delaunay simplices
(containing the circumcenter in the interior)*



Čech intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff




Q

Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff

On $S \subseteq Q'$



Q




On S

Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$\text{On } S \subseteq Q' \subseteq \text{Encl } S.$$



Q



$\text{On } S$




$\text{Encl } S$

Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$Q' \in [\text{On } S, \text{Encl } S].$$



Q



$\text{On } S$



$\text{Encl } S$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $Q = \text{On } S \subseteq X$.

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $Q = \text{On } S \subseteq X$. Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

The front and back faces of a simplex


Let S be the smallest circumsphere of a simplex $Q = \text{On } S \subseteq X$. Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $Q = \text{On } S \subseteq X$. Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$


$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



Delaunay intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff





Delaunay intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff

$$\text{Front } S \subseteq Q'$$




Delaunay intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff


$$\text{Front } S \subseteq Q' \subseteq \text{On } S.$$



Q



Front S




On S

Delaunay intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff


$$Q' \in [\text{Front } S, \text{On } S].$$



Q



Front S



On S

Selective Delaunay complexes

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\underset{r,z}{\text{minimize}}$$

$$r$$

$$\text{subject to}$$

$$\|z - q\| \leq r, \quad q \in Q,$$

$$\|z - e\| \geq r, \quad e \in E.$$

Here r is the radius of the sphere S , and z is the center of S .

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

minimize
 r,z

r

subject to

$$\begin{aligned}\|z - q\| &\leq r, & q \in Q, \\ \|z - e\| &\geq r, & e \in E.\end{aligned}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

$\check{\text{C}}$ ech and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

Proposition

A sphere S enclosing Q and excluding E is minimal iff

S is the smallest circumsphere of $\text{On } S$,

$Q \in [\text{Front } S, \text{Encl } S]$, and

$E \in [\text{Back } S, \text{Excl } S]$.

$\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

Proposition

A sphere S enclosing Q and excluding E is minimal iff

S is the smallest circumsphere of $\text{On } S$,

$Q \in [\text{Front } S, \text{Encl } S]$, and

$E \in [\text{Back } S, \text{Excl } S]$.

Corollary

The $\check{\text{C}}\text{ech}$ intervals are of the form $[\text{On } S, \text{Encl } S]$.

$\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

Proposition

A sphere S enclosing Q and excluding E is minimal iff

S is the smallest circumsphere of $\text{On } S$,

$Q \in [\text{Front } S, \text{Encl } S]$, and

$E \in [\text{Back } S, \text{Excl } S]$.

Corollary

The $\check{\text{C}}\text{ech}$ intervals are of the form $[\text{On } S, \text{Encl } S]$.

The Delaunay intervals are of the form $[\text{Front } S, \text{On } S]$.

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $Q \subseteq X \mapsto f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if $Q \subseteq X$ has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$ (*selective Delaunay simplices*)

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $Q \subseteq X \mapsto f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if $Q \subseteq X$ has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$ (*selective Delaunay simplices*)
- *Selective Delaunay complex* $\text{Del}_r(X, E) = f_D^{-1}(-\infty, r]$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $Q \subseteq X \mapsto f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if $Q \subseteq X$ has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$ (*selective Delaunay simplices*)
- *Selective Delaunay complex* $\text{Del}_r(X, E) = f_D^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2015)

Let $E \subseteq F \subseteq X$. Then

$$\text{Del}_r(X, E) \setminus \text{Del}_r(X, E) \cap \text{Del}(X, F) \setminus \text{Del}_r(X, F).$$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $Q \subseteq X \mapsto f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if $Q \subseteq X$ has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$ (*selective Delaunay simplices*)
- *Selective Delaunay complex* $\text{Del}_r(X, E) = f_D^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2015)

Let $E \subseteq F \subseteq X$. Then


$$\text{Del}_r(X, E) \setminus \text{Del}_r(X, E) \cap \text{Del}(X, F) \setminus \text{Del}_r(X, F).$$

Note: choosing $E = \emptyset$ and $F = X$ yields


$$\text{Cech}_r(X) \setminus \text{Del}(X) \cap \text{Cech}_r(X) \setminus \text{Del}_r(X).$$

Connecting different Delaunay complexes

X




Y



Connecting different Delaunay complexes

$B_r(X)$




$B_r(Y)$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



$\text{Del}_r(X)$





$\text{Del}_r(X \cup Y)$



$\text{Del}_r(Y)$



Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$




Stability

The pipeline of topological data analysis




The pipeline of topological data analysis



Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.



Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- matching $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- matching $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$
- δ -matching of barcodes:

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- matching $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$
- δ -matching of barcodes:
 - matched intervals have endpoints within distance $\leq \delta$

Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- matching $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$
- δ -matching of barcodes:
 - matched intervals have endpoints within distance $\leq \delta$
 - unmatched intervals have length $\leq 2\delta$

Stability revisited



Stability revisited



Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

If $\|f - g\|_\infty \leq \delta$ then $F_t \subseteq G_{t+\delta}$ and $G_t \subseteq F_{t+\delta}$.


Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

If $\|f - g\|_\infty \leq \delta$ then $F_t \subseteq G_{t+\delta}$ and $G_t \subseteq F_{t+\delta}$.

So the sublevel sets are δ -interleaved:



Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

If $\|f - g\|_\infty \leq \delta$ then $F_t \subseteq G_{t+\delta}$ and $G_t \subseteq F_{t+\delta}$.

So the sublevel sets are δ -interleaved:


$$\begin{array}{ccccc} H_*(F_t) & \longrightarrow & H_*(F_{t+2\delta}) & & \\ \searrow & & \nearrow & & \searrow \\ & & H_*(G_{t+\delta}) & \longrightarrow & H_*(G_{t+3\delta}) \end{array}$$

Homology is a *functor*: homology groups are interleaved too.


Geometric interleavings




Geometric interleavings




Geometric interleavings




Geometric interleavings



Geometric interleavings



Geometric interleavings



Barcodes: the structure of persistence

Theorem (Crawley-Boevey 2012)

Let $M = (M_t)_{t \in \mathbb{R}}$ be a diagram of vector spaces ($\dim M_t < \infty$) indexed over the reals (with linear maps $M_s \rightarrow M_t$ for each $s \leq t$).

Barcodes: the structure of persistence

Theorem (Crawley-Boevey 2012)

Let $M = (M_t)_{t \in \mathbb{R}}$ be a diagram of vector spaces ($\dim M_t < \infty$) indexed over the reals (with linear maps $M_s \rightarrow M_t$ for each $s \leq t$).

Then M is uniquely (up to isomorphism) described by a collection of intervals $B(M)$ (the barcode of M).

Barcodes: the structure of persistence

Theorem (Crawley-Boevey 2012)

Let $M = (M_t)_{t \in \mathbb{R}}$ be a diagram of vector spaces ($\dim M_t < \infty$) indexed over the reals (with linear maps $M_s \rightarrow M_t$ for each $s \leq t$).

Then M is uniquely (up to isomorphism) described by a collection of intervals $B(M)$ (the barcode of M).

- Motivates use of homology with field coefficients

Algebraic stability of persistence barcodes


Theorem (Chazal et al. 2009, 2012; B, Lesnick 2014)

*If two \mathbb{R} -indexed diagrams of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.*

Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2014)


If two \mathbb{R} -indexed diagrams of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.



Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2014)

If two \mathbb{R} -indexed diagrams of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.




- converse statement also holds (isometry theorem)

Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2014)

If two \mathbb{R} -indexed diagrams of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.




- converse statement also holds (isometry theorem)
- original proof: indirect, elaborate construction (80 page paper)

Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2014)

If two \mathbb{R} -indexed diagrams of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.



- converse statement also holds (isometry theorem)
- original proof: indirect, elaborate construction (80 page paper)
- our proof: constructs matching directly from interleaving maps

Stability via induced matchings




Stability via induced matchings




Stability via induced matchings




Stability via induced matchings



Stability via induced matchings



Stability via induced matchings



Stability via induced matchings

