

Delaunay complexes, persistence diagrams, and induced matchings

Ulrich Bauer

TUM

September 17, 2015

Shape Up, TU Berlin

Joint work with
Herbert Edelsbrunner (IST Austria)
and
Michael Lesnick (Columbia, IMA)



112



?





≈





?
































What is persistent homology?





What is persistent homology?





What is persistent homology?





What is persistent homology?




What is persistent homology?




What is persistent homology?






T. Nakamura, Y. Hiraoka, A. Hirata, E. G. Escolar, and Y. Nishiura. Persistent Homology and Many-Body Atomic Structure for Medium-Range Order in the Glass. *Nanotechnology* 26 (2015) 304001.




T. Nakamura, Y. Hiraoka, A. Hirata, E. G. Escolar, and Y. Nishiura. Persistent Homology and Many-Body Atomic Structure for Medium-Range Order in the Glass. *Nanotechnology* 26 (2015) 304001.



T. Nakamura, Y. Hiraoka, A. Hirata, E. G. Escolar, and Y. Nishiura. Persistent Homology and Many-Body Atomic Structure for Medium-Range Order in the Glass. *Nanotechnology* 26 (2015) 304001.


The pipeline of topological data analysis



Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)


If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.



Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- matching $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- *matching* $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$
- δ -*matching of barcodes*:

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- *matching* $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$
- *δ -matching of barcodes*:
 - matched intervals have endpoints within distance $\leq \delta$

Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$,
then there exists a δ -matching of their barcodes.




- matching $A \rightarrow B$: bijection of subsets $A' \subseteq A, B' \subseteq B$
- δ -matching of barcodes:
 - matched intervals have endpoints within distance $\leq \delta$
 - unmatched intervals have length $\leq 2\delta$


Stability for functions in the big picture




Stability for functions in the big picture



Stability for functions in the big picture



Stability for functions in the big picture



Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

If $\|f - g\|_\infty \leq \delta$ then $F_t \subseteq G_{t+\delta}$ and $G_t \subseteq F_{t+\delta}$.


Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

If $\|f - g\|_\infty \leq \delta$ then $F_t \subseteq G_{t+\delta}$ and $G_t \subseteq F_{t+\delta}$.

So the sublevel sets are *δ -interleaved*:



Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$,
- $G_t = g^{-1}(-\infty, t]$.

If $\|f - g\|_\infty \leq \delta$ then $F_t \subseteq G_{t+\delta}$ and $G_t \subseteq F_{t+\delta}$.

So the sublevel sets are *δ -interleaved*:


$$\begin{array}{ccccc} H_*(F_t) & \longrightarrow & H_*(F_{t+2\delta}) & & \\ \searrow & & \nearrow & & \searrow \\ & & H_*(G_{t+\delta}) & \longrightarrow & H_*(G_{t+3\delta}) \end{array}$$

Homology is a *functor*: homology groups are interleaved too.


Geometric interleavings




Geometric interleavings




Geometric interleavings




Geometric interleavings



Geometric interleavings



Geometric interleavings



Algebraic stability of persistence barcodes


Theorem (Chazal et al. 2009, 2012)

*If two 1-parameter families of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.*

Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012)


*If two 1-parameter families of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.*



Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012)

*If two 1-parameter families of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.*




Theorem (B, Lesnick 2014)

Such a δ -matching is directly induced by a δ -interleaving.

Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012)

*If two 1-parameter families of vector spaces are δ -interleaved,
then there exists a δ -matching of their barcodes.*



Theorem (B, Lesnick 2014)

Such a δ -matching is directly induced by a δ -interleaving.

Specifically: by any of the two 1-parameter families of linear maps in a δ -interleaving (though the barcode of their images).

Stability via induced matchings



Stability via induced matchings




$B(M)$




$B(N(\delta))$


Stability via induced matchings




Stability via induced matchings




Stability via induced matchings




Stability via induced matchings




Stability via induced matchings




Connect the dots: topology from geometry




Connect the dots: topology from geometry




Connect the dots: topology from geometry




Connect the dots: topology from geometry



Connect the dots: topology from geometry




Connect the dots: topology from geometry



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X



$\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

Two functions on simplices $Q \subseteq X$:

$\check{\text{C}}\text{ech}$ function $f_C(Q)$:

radius of smallest enclosing sphere of Q



$\check{\text{C}}\text{ech}$ and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X


Two functions on simplices $Q \subseteq X$:

$\check{\text{C}}\text{ech function}$ $f_C(Q)$:

radius of smallest enclosing sphere of Q

Delaunay function $f_D(Q)$:

radius of smallest empty circumsphere



$\check{\text{C}}\text{ech}$ and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

Two functions on simplices $Q \subseteq X$:


$\check{\text{C}}\text{ech function}$ $f_C(Q)$:

radius of smallest enclosing sphere of Q

Delaunay function $f_D(Q)$:

radius of smallest empty circumsphere

- defined only if Q has an empty circumsphere: $Q \in \text{Del}(X)$



\check{C} ech and Delaunay complexes from functions

Define for any radius r :

$\check{\text{C}}\text{ech}$ and Delaunay complexes from functions

Define for any radius r :

- $\check{\text{C}}\text{ech complex } \text{Cech}_r = f_C^{-1}(-\infty, r]$

Čech and Delaunay complexes from functions


Define for any radius r :

- Čech complex $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$

$\check{\text{C}}\text{ech}$ and Delaunay complexes from functions

Define for any radius r :


- $\check{\text{C}}\text{ech complex}$ $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- $\text{Delaunay}-\check{\text{C}}\text{ech complex}$ $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$



$\check{\text{C}}\text{ech}$ and Delaunay complexes from functions

Define for any radius r :


- $\check{\text{C}}\text{ech complex}$ $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- $\text{Delaunay}-\check{\text{C}}\text{ech complex}$ $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$
 - restriction of $\check{\text{C}}\text{ech}$ complex to Delaunay simplices



Čech and Delaunay complexes from functions

Define for any radius r :


- Čech complex $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- Delaunay–Čech complex $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$
 - restriction of Čech complex to Delaunay simplices
- Delaunay complex (α -shape, for $\alpha = r$) $\text{Del}_r = f_D^{-1}(-\infty, r]$



Čech and Delaunay complexes from functions

Define for any radius r :


- Čech complex $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- Delaunay–Čech complex $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$
 - restriction of Čech complex to Delaunay simplices
- Delaunay complex (α -shape, for $\alpha = r$) $\text{Del}_r = f_D^{-1}(-\infty, r]$
 - all simplices having an empty circumsphere of radius $\leq r$



Čech and Delaunay complexes from functions

Define for any radius r :

- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$
 - restriction of Čech complex to Delaunay simplices
- *Delaunay complex (α -shape, for $\alpha = r$)* $\text{Del}_r = f_D^{-1}(-\infty, r]$
 - all simplices having an empty circumsphere of radius $\leq r$




A sequence of collapses

Theorem (B, Edelsbrunner 2015)

Čech, Delaunay–Čech, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




A sequence of collapses

Theorem (B, Edelsbrunner 2015)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*

A sequence of collapses

Theorem (B, Edelsbrunner 2015)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*
- Explicit chain maps inducing isomorphisms in homology

A sequence of collapses

Theorem (B, Edelsbrunner 2015)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$



- All collapses are induced by a single *discrete gradient field*
- Explicit chain maps inducing isomorphisms in homology
- Also works for weighted point sets

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

minimize
 $r_{,z}$

r

subject to

$$\|z - q\| \leq r, \quad q \in Q,$$

$$\|z - e\| \geq r, \quad e \in E.$$

Here r is the radius of the sphere S , and z is the center of S .

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

minimize
 r, z

subject to

r

$$\|z - q\| \leq r, \quad q \in Q,$$

$$\|z - e\| \geq r, \quad e \in E.$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex* $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex* $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2015)

Let $E \subseteq F \subseteq X$. Then

$$\text{Del}_r(X, E) \setminus \text{Del}_r(X, E) \cap \text{Del}(X, F) \setminus \text{Del}_r(X, F).$$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex* $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2015)

Let $E \subseteq F \subseteq X$. Then


$$\text{Del}_r(X, E) \setminus \text{Del}_r(X, E) \cap \text{Del}(X, F) \setminus \text{Del}_r(X, F).$$

Note: choosing $E = \emptyset$ and $F = X$ yields


$$\text{Cech}_r(X) \setminus \text{DelCech}_r(X) \setminus \text{Del}_r(X).$$

Connecting different Delaunay complexes

X




Y



Connecting different Delaunay complexes

$B_r(X)$




$B_r(Y)$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



$\text{Del}_r(X)$





$\text{Del}_r(X \cup Y)$



$\text{Del}_r(Y)$



Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



Wrapping up

- Persistent homology can describe connectedness of shapes at different scales
- Delaunay complexes enable efficient computations
- Induced matchings provide an identification of topological features for similar shapes