

# The Morse theory of Čech and Delaunay complexes

3<sup>rd</sup> ERC SDModels workshop

*Discrete Models in Geometry and Topology*


Ulrich Bauer

TUM

March 23, 2015

Joint work with Herbert Edelsbrunner (IST Austria)

# Connect the dots: topology from geometry




# Connect the dots: topology from geometry




# Connect the dots: topology from geometry




# Connect the dots: topology from geometry




# Connect the dots: topology from geometry



# Connect the dots: topology from geometry



# Connect the dots: topology from geometry



# Čech and Delaunay functions


$X \subset \mathbb{R}^d$ : finite point set (in general position)



# Čech and Delaunay functions

$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$



# $\check{\text{C}}$ ech and Delaunay functions

$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$



# Čech and Delaunay functions


$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$

Two functions on simplices  $Q \subseteq X$ :

*Čech function*  $f_C(Q)$ :

radius of smallest enclosing sphere of  $Q$



# Čech and Delaunay functions

$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$


Two functions on simplices  $Q \subseteq X$ :

*Čech function*  $f_C(Q)$ :

radius of smallest enclosing sphere of  $Q$

*Delaunay function*  $f_D(Q)$ :

radius of smallest empty circumsphere



# $\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$

Two functions on simplices  $Q \subseteq X$ :

*$\check{\text{C}}\text{ech function}$*   $f_C(Q)$ :

radius of smallest enclosing sphere of  $Q$



*Delaunay function*  $f_D(Q)$ :

radius of smallest empty circumsphere

- defined only if  $Q$  has an empty circumsphere:  $Q \in \text{Del}(X)$

# Čech and Delaunay complexes from functions

Define for any radius  $r$ :

# Čech and Delaunay complexes from functions

Define for any radius  $r$ :

- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$

# Čech and Delaunay complexes from functions


Define for any radius  $r$ :

- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$

# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- Delaunay–Čech complex  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- Delaunay–Čech complex  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- Delaunay–Čech complex  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices
- Delaunay complex ( $\alpha$ -shape, for  $\alpha = r$ )  $\text{Del}_r = f_D^{-1}(-\infty, r]$



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- Delaunay–Čech complex  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices
- Delaunay complex ( $\alpha$ -shape, for  $\alpha = r$ )  $\text{Del}_r = f_D^{-1}(-\infty, r]$ 
  - all simplices having an empty circumsphere of radius  $\leq r$



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :

- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- Delaunay–Čech complex  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices
- Delaunay complex ( $\alpha$ -shape, for  $\alpha = r$ )  $\text{Del}_r = f_D^{-1}(-\infty, r]$ 
  - all simplices having an empty circumsphere of radius  $\leq r$



# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?

# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$


- Are all three complexes homotopy equivalent?
- Are they related by a sequence of simplicial collapses?

# Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014)

$\check{\text{C}}\text{ech}$ , Delaunay– $\check{\text{C}}\text{ech}$ , Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




# Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014)

$\check{\text{C}}\text{ech}$ , Delaunay– $\check{\text{C}}\text{ech}$ , Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*

# Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014)

$\check{\text{C}}\text{ech}$ , Delaunay– $\check{\text{C}}\text{ech}$ , Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*
- Explicit chain maps inducing isomorphisms in homology

# Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014)

$\check{\text{C}}\text{ech}$ , Delaunay– $\check{\text{C}}\text{ech}$ , Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*
- Explicit chain maps inducing isomorphisms in homology
- Also works for weighted point sets

# Discrete Morse theory

# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs  $\{L, U\}$ , where  $L$  is a face of  $U$  with codimension 1.




# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs  $\{L, U\}$ , where  $L$  is a face of  $U$  with codimension 1.

- indicated by an arrow from  $L$  to  $U$




# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs  $\{L, U\}$ , where  $L$  is a face of  $U$  with codimension 1.

- indicated by an arrow from  $L$  to  $U$

The singletons are called *critical simplices*.




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes



# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes

•  
1

# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes
- the level sets  $f^{-1}(t)$  form a discrete vector field (the *discrete gradient* of  $f$ )




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :


- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes
- the level sets  $f^{-1}(t)$  form a discrete vector field (the *discrete gradient* of  $f$ )

If  $f^{-1}(t) = \{Q\}$  then  $t$  is a *critical value*.



# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**

*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**

*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

Theorem (Forman 1998)

If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**


*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*



Let  $V$  be a discrete gradient field on a simplicial complex  $K$ ,  
and let  $L$  be a subcomplex of  $K$ .

**Corollary**

*If  $K \setminus L$  is the union of some pairs of  $V$ ,  
then  $K \searrow L$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**


*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*



Let  $V$  be a discrete gradient field on a simplicial complex  $K$ ,  
and let  $L$  be a subcomplex of  $K$ .

**Corollary**

*If  $K \setminus L$  is the union of some pairs of  $V$ ,  
then  $K \searrow L$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**


*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*



Let  $V$  be a discrete gradient field on a simplicial complex  $K$ ,  
and let  $L$  be a subcomplex of  $K$ .

**Corollary**

*If  $K \setminus L$  is the union of some pairs of  $V$ ,  
then  $K \searrow L$ .*



We say that  $V$  induces the collapse  $K \searrow L$ .


## Unfortunately...

Neither the Čech nor the Delaunay functions  
are discrete Morse functions!


# Unfortunately...

Neither the Čech nor the Delaunay functions are discrete Morse functions!

- Example: two simplices  $Q, Q'$  with  $f_C(Q) = f_C(Q')$  such that neither is a face of the other:



$Q$




$Q'$

# Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




# Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




- indicated by an arrow from  $L$  to  $U$

# Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$



- indicated by an arrow from  $L$  to  $U$


The singletons are called *critical simplices*.

# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes



# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex  
is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$   
are subcomplexes

•  
1

# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :


- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes
- the level sets  $f^{-1}(t)$  form a generalized vector field (the *discrete gradient* of  $f$ )



## Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :




# Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :

- choose an arbitrary vertex  $x \in U \setminus L$



# Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :

- choose an arbitrary vertex  $x \in U \setminus L$
- partition  $[L, U]$  into pairs  $(Q \setminus \{x\}, Q \cup \{x\})$  for all  $Q \in [L, U]$ .




# Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :

- choose an arbitrary vertex  $x \in U \setminus L$
- partition  $[L, U]$  into pairs  $(Q \setminus \{x\}, Q \cup \{x\})$  for all  $Q \in [L, U]$ .



Therefore the collapsing theorems also hold for generalized discrete Morse functions.

# Čech and Delaunay intervals

# Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

# Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*

- *$Q$  is a critical simplex of  $f_C$*

# Morse theory of Čech and Delaunay complexes

## Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*

- *$Q$  is a critical simplex of  $f_C$*
- *$Q$  is a critical simplex of  $f_D$*

# Morse theory of Čech and Delaunay complexes

## Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*

- *$Q$  is a critical simplex of  $f_C$*
- *$Q$  is a critical simplex of  $f_D$*
- $f_D(Q) = f_C(Q)$


# Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*


- *$Q$  is a critical simplex of  $f_C$*
- *$Q$  is a critical simplex of  $f_D$*
- $f_D(Q) = f_C(Q)$
- *$Q$  is a centered Delaunay simplex  
(containing the circumcenter in the interior)*



# Čech intervals

## Lemma

Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff




$Q$

# Čech intervals


## Lemma

Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff

On  $S \subseteq Q'$



$Q$




On  $S$

# Čech intervals


## Lemma

Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff


$\text{On } S \subseteq Q' \subseteq \text{Encl } S$ .



$Q$



$\text{On } S$




$\text{Encl } S$

# Čech intervals


## Lemma

Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff


$$Q' \in [\text{On } S, \text{Encl } S].$$



$Q$



$\text{On } S$



$\text{Encl } S$

## The front and back faces of a simplex

Let  $S$  be the smallest circumsphere of a simplex On  $S \subseteq X$ .

## The front and back faces of a simplex

Let  $S$  be the smallest circumsphere of a simplex  $\text{On } S \subseteq X$ .

Then the center  $z$  of  $S$  is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

# The front and back faces of a simplex

Let  $S$  be the smallest circumsphere of a simplex. On  $S \subseteq X$ .


Then the center  $z$  of  $S$  is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



# The front and back faces of a simplex



Let  $S$  be the smallest circumsphere of a simplex. On  $S \subseteq X$ .  
Then the center  $z$  of  $S$  is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$






# Delaunay intervals

## Lemma

Let  $Q \subseteq X$  be a simplex with smallest empty circumsphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest empty circumsphere  $S$  iff

$$Q' \in [\text{Front } S, \text{On } S].$$



# Selective Delaunay complexes

## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

minimize  
 $r, z$

$r$

subject to

$$\|z - q\| \leq r, \quad q \in Q,$$

$$\|z - e\| \geq r, \quad e \in E.$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

minimize  
 $r, z$

$r$

subject to

$$\begin{aligned}\|z - q\| \leq r, & \quad q \in Q, \\ \|z - e\| \geq r, & \quad e \in E.\end{aligned}$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

- Čech function: choose  $E = \emptyset$
- Delaunay function: choose  $E = X$

# $\check{\text{C}}$ ech and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

## Proposition

*A sphere  $S$  enclosing  $Q$  and excluding  $E$  is minimal iff*

*$S$  is the smallest circumsphere of  $\text{On } S$ ,*

*$Q \in [\text{Front } S, \text{Encl } S]$ , and*

*$E \in [\text{Back } S, \text{Excl } S]$ .*

# $\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

## Proposition

*A sphere  $S$  enclosing  $Q$  and excluding  $E$  is minimal iff*

*$S$  is the smallest circumsphere of  $\text{On } S$ ,*

*$Q \in [\text{Front } S, \text{Encl } S]$ , and*

*$E \in [\text{Back } S, \text{Excl } S]$ .*

## Corollary

*The  $\check{\text{C}}\text{ech}$  intervals are of the form  $[\text{On } S, \text{Encl } S]$ .*

# $\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

## Proposition

*A sphere  $S$  enclosing  $Q$  and excluding  $E$  is minimal iff*

*$S$  is the smallest circumsphere of  $\text{On } S$ ,*

*$Q \in [\text{Front } S, \text{Encl } S]$ , and*

*$E \in [\text{Back } S, \text{Excl } S]$ .*

## Corollary

*The  $\check{\text{C}}\text{ech}$  intervals are of the form  $[\text{On } S, \text{Encl } S]$ .*

*The Delaunay intervals are of the form  $[\text{Front } S, \text{On } S]$ .*

## Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$

## Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$

# Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex*  $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

# Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex*  $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

## Theorem (B, Edelsbrunner 2014)

Let  $E \subseteq F \subseteq X$ . Then

$$\text{Del}_r(X, E) \downarrow \text{Del}_r(X, E) \cap \text{Del}(X, F) \downarrow \text{Del}_r(X, F).$$

# Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex*  $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

## Theorem (B, Edelsbrunner 2014)

Let  $E \subseteq F \subseteq X$ . Then


$$\text{Del}_r(X, E) \setminus \text{Del}_r(X, E) \cap \text{Del}(X, F) \setminus \text{Del}_r(X, F).$$

Note: choosing  $E = \emptyset$  and  $F = X$  yields


$$\text{Cech}_r(X) \setminus \text{DelCech}_r(X) \setminus \text{Del}_r(X).$$

# Connecting different Delaunay complexes

$X$



$Y$



# Connecting different Delaunay complexes

$B_r(X)$





$B_r(Y)$




# Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$




# Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



$\text{Del}_r(X)$





$\text{Del}_r(X \cup Y)$



$\text{Del}_r(Y)$



# Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



# Connecting different Delaunay complexes


$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



# Wrap complexes


# Wrap complexes

Generalizes and greatly simplifies the surface reconstruction algorithm *Wrap* (Edelsbrunner 1995, Geomagic)



# Wrap complexes


Consider the Delaunay function  $f_D$  of  $X$ .



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order




# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$




# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$




# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$  in that partial order

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$



The Delaunay intervals induce a collapse  $\text{Del}_r \searrow \text{Wrap}_r$ .

# Wrapping up

- Čech and Delaunay complexes from Morse functions
- Explicit homotopy equivalence by simplicial collapses
- Simple definition and generalization of *Wrap* complexes