

Criticality and simplification

Ulrich Bauer

IST Austria

March 5, 2012

What is a critical point/value?

Classical definition:

For a smooth function $f : M \rightarrow \mathbb{R}$:

- ▶ x is a *critical point* of f : differential $D_x f \equiv 0$
- ▶ $f(x)$ is a *critical value*

What is a critical point/value?

Classical definition:

For a smooth function $f : M \rightarrow \mathbb{R}$:

- ▶ x is a *critical point* of f : differential $D_x f \equiv 0$
- ▶ $f(x)$ is a *critical value*

For a general smooth map $f : M \rightarrow N$:

- ▶ x is a *critical point* of f : differential $D_x f$ is not surjective

Homological critical values

$M_t = f^{-1}(-\infty, t]$: sublevel set of $f : M \rightarrow \mathbb{R}$

Homological critical values

$M_t = f^{-1}(-\infty, t]$: sublevel set of $f : M \rightarrow \mathbb{R}$

Homology of sublevel sets changes only at critical values:

- ▶ If $[t - \epsilon, t]$ contains no critical value, then
 $M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Homological critical values

$M_t = f^{-1}(-\infty, t]$: sublevel set of $f : M \rightarrow \mathbb{R}$

Homology of sublevel sets changes only at critical values:

- ▶ If $[t - \epsilon, t]$ contains no critical value, then
 $M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Turn this property into a definition [Cohen-Steiner et al. 2005]:

- ▶ t is a *homological critical value*:
 $\nexists \epsilon > 0: M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Homological critical values

$M_t = f^{-1}(-\infty, t]$: sublevel set of $f : M \rightarrow \mathbb{R}$

Homology of sublevel sets changes only at critical values:

- ▶ If $[t - \epsilon, t]$ contains no critical value, then
 $M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Turn this property into a definition [Cohen-Steiner et al. 2005]:

- ▶ t is a *homological critical value*:
 $\nexists \epsilon > 0: M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Homological critical values

$M_t = f^{-1}(-\infty, t]$: sublevel set of $f : M \rightarrow \mathbb{R}$

Homology of sublevel sets changes only at critical values:

- ▶ If $[t - \epsilon, t]$ contains no critical value, then
 $M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Turn this property into a definition [Cohen-Steiner et al. 2005]:

- ▶ t is a *homological critical value*:
 $\nexists \epsilon > 0: M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology
- ▶ also applicable to non-smooth functions

Homological critical values

$M_t = f^{-1}(-\infty, t]$: sublevel set of $f : M \rightarrow \mathbb{R}$

Homology of sublevel sets changes only at critical values:

- ▶ If $[t - \epsilon, t]$ contains no critical value, then
 $M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Turn this property into a definition [Cohen-Steiner et al. 2005]:

- ▶ t is a *homological critical value*:
 $\nexists \epsilon > 0: M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology
- ▶ also applicable to non-smooth functions
- ▶ homological critical values are stable

Homological critical values

$M_t = f^{-1}(-\infty, t]$: sublevel set of $f : M \rightarrow \mathbb{R}$


Homology of sublevel sets changes only at critical values:

- ▶ If $[t - \epsilon, t]$ contains no critical value, then
 $M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology

Turn this property into a definition [Cohen-Steiner et al. 2005]:


- ▶ t is a *homological critical value*:
 $\nexists \epsilon > 0: M_{t-\epsilon} \hookrightarrow M_t$ induces isomorphism in homology
- ▶ also applicable to non-smooth functions
- ▶ homological critical values are stable
- ▶ a homological regular value might be critical in the classical sense!

Topological denoising of a function




Noise (even small) can create lots of critical points

Topological denoising of a function



Noise (even small) can create lots of critical points

Topological denoising of a function




Noise (even small) can create lots of critical points

Topological denoising by simplification of critical set:

- ▶ removing critical points caused by noise

Topological denoising of a function



Noise (even small) can create lots of critical points

Topological denoising by simplification of critical set:

- ▶ removing critical points caused by noise
- ▶ given a function f on a surface and $\delta > 0$,
find a function f_δ that:
 - ▶ minimizes number of critical points
 - ▶ stays close to input function: $\|f_\delta - f\|_\infty < \delta$


Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets

Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D




Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets



Example: connected components in 1D



Persistence diagrams [Cohen-Steiner et al., 2005]



Persistence diagrams [Cohen-Steiner et al., 2005]




Stability of persistence diagrams

Theorem (Cohen-Steiner et al., 2005)

Let $\|f - g\|_\infty \leq \delta$.

The persistence pairs of f that have persistence $> 2\delta$ can be mapped injectively to the persistence pairs of g .




Stability of persistence diagrams

Theorem (Cohen-Steiner et al., 2005)

Let $\|f - g\|_\infty \leq \delta$.

The persistence pairs of f that have persistence $> 2\delta$ can be mapped injectively to the persistence pairs of g .

Corresponding points p_f, p_g in the persistence diagrams have distance $\|p_f - p_g\|_\infty \leq \delta$.



A bound on number of critical points

Corollary

Let f be a discrete Morse function on a surface and let $\delta > 0$.

Then for every function f_δ with $\|f_\delta - f\|_\infty < \delta$ we have:

$$\begin{aligned} & \# \text{ critical points of } f_\delta \\ & \geq \# \text{ critical points of } f \text{ with persistence } \geq 2\delta. \end{aligned}$$

Persistence-guided simplification

Persistence is a quantity that

- ▶ bounds the stability of critical points

Persistence-guided simplification

Persistence is a quantity that

- ▶ bounds the stability of critical points
- ▶ can help to decide whether a critical point may be caused by noise

Persistence-guided simplification

Persistence is a quantity that

- ▶ bounds the stability of critical points
- ▶ can help to decide whether a critical point may be caused by noise

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:

- ▶ remove all persistence pairs of f with $\text{persistence} < 2\delta$

Persistence-guided simplification


Persistence is a quantity that

- ▶ bounds the stability of critical points
- ▶ can help to decide whether a critical point may be caused by noise


Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:

- ▶ remove all persistence pairs of f with $\text{persistence} < 2\delta$
- ▶ leave all other persistence pairs unmodified


Side-effects of elimination



Side-effects of elimination




Side-effects of elimination




For an optimal solution, we must allow the critical values to change!


Canceling critical points of a gradient field




Canceling critical points of a gradient field




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function



Cancelling critical points of a function



Cancelling critical points of a function



Results on topological simplification

For functions on surfaces:

- ▶ Every persistence pair can eventually be canceled in the Morse-theoretic sense;

Results on topological simplification

For functions on surfaces:

- ▶ Every persistence pair can eventually be canceled in the Morse-theoretic sense;
- ▶ On surfaces, the lower bound can always be achieved (in time $O(n \log n)$) [B., Lange, Wardetzky 2011]

Results on topological simplification

For functions on surfaces:

- ▶ Every persistence pair can eventually be canceled in the Morse-theoretic sense;
- ▶ On surfaces, the lower bound can always be achieved (in time $O(n \log n)$) [B., Lange, Wardetzky 2011]
- ▶ When restricting to simplexwise linear functions, the problem becomes NP-hard [Gray et al. 2010, B. 2011]

Results on topological simplification

For functions on surfaces:

- ▶ Every persistence pair can eventually be canceled in the Morse-theoretic sense;
- ▶ On surfaces, the lower bound can always be achieved (in time $O(n \log n)$) [B., Lange, Wardetzky 2011]
- ▶ When restricting to simplexwise linear functions, the problem becomes NP-hard [Gray et al. 2010, B. 2011]

Does not hold anymore in 3D. Moreover:

Results on topological simplification

For functions on surfaces:

- ▶ Every persistence pair can eventually be canceled in the Morse-theoretic sense;
- ▶ On surfaces, the lower bound can always be achieved (in time $O(n \log n)$) [B., Lange, Wardetzky 2011]
- ▶ When restricting to simplexwise linear functions, the problem becomes NP-hard [Gray et al. 2010, B. 2011]

Does not hold anymore in 3D. Moreover:

- ▶ Topological function simplification is NP-hard [Egecioglu, Gonzalez 1995]

Results on topological simplification


For functions on surfaces:

- ▶ Every persistence pair can eventually be canceled in the Morse-theoretic sense;
- ▶ On surfaces, the lower bound can always be achieved (in time $O(n \log n)$) [B., Lange, Wardetzky 2011]
- ▶ When restricting to simplexwise linear functions, the problem becomes NP-hard [Gray et al. 2010, B. 2011]

Does not hold anymore in 3D. Moreover:

- ▶ Topological function simplification is NP-hard [Egecioglu, Gonzalez 1995]
- ▶ Even simplification of a single isosurface is NP-hard (ongoing joint work with Amenta, Attali, Devillers, Glisse, Lieutier)





Robustness

What about other tolerance norms than $\|\cdot\|_\infty$?

- ▶ Persistence is not robust to outliers!

(ongoing joint work with Schönlieb, Munk, Wardetzky)

A robust denoising method

A well-known denoising method (Rudin, Osher, Fatemi, 1992):

Given $f \in L^1([a, b])$, $\alpha \geq 0$. Find u minimizing

$$\|f - u\|_2 + 2\alpha \text{TV}(u) \quad (\text{ROF})$$

A robust denoising method

A well-known denoising method (Rudin, Osher, Fatemi, 1992):

Given $f \in L^1([a, b])$, $\alpha \geq 0$. Find u minimizing

$$\|f - u\|_2 + 2\alpha \text{TV}(u) \quad (\text{ROF})$$

Recall the definition of TV (*total variation*):

- ▶ For differentiable u : $\text{TV}(u) = \int_a^b |u'(t)| dt$
- ▶ Can be continuously extended to L^1

Equivalence with the taut string problem

Theorem (Grasmair, 2006)

Let u_α be a minimizer of the ROF functional. Then $u_\alpha = U'_\alpha$, where U_α minimizes



$$\int_a^b \sqrt{1 + U'(t)^2} dt$$

subject to


$$\|U - F\|_\infty \leq \alpha, \quad U(a) = F(a), \quad U(b) = F(b),$$

where $F(t) = \int_a^t f(\tau) d\tau$.


An example




An example



Properties:




An example



Properties:

- ▶ u_α also minimizes the number of (homological) critical points


An example



Properties:

- ▶ u_α also minimizes the number of (homological) critical points
- ▶ u_α coincides with f apart from some intervals, on which it is constant

An example



Properties:

- ▶ u_{α} also minimizes the number of (homological) critical points
- ▶ u_{α} coincides with f apart from some intervals, on which it is constant
- ▶ Some but not all constant intervals cancel critical points

A stable descriptor for critical points

Consider the values α at which the number of critical points of u_α changes

- ▶ Decreasing sequence s_i of “stability” values
- ▶ $s_i(f) := \inf_G \|G - F\|_\infty$: $g = G'$ has $\leq i$ critical points

A stable descriptor for critical points

Consider the values α at which the number of critical points of u_α changes

- ▶ Decreasing sequence s_i of “stability” values
- ▶ $s_i(f) := \inf_G \|G - F\|_\infty$: $g = G'$ has $\leq i$ critical points

This sequence is stable under perturbations of F :

$$|s_i(f) - s_i(\tilde{f})| \leq \|F - \tilde{F}\|_\infty$$

Reason: s_i is a distance function
(to the set of functions with $\leq i$ critical points)

Estimating the number of critical points

Consider:

- ▶ 1D signal, n measurements with Gaussian noise $\sigma \sim \frac{1}{n}$
- ▶ Estimate number of critical points by thresholding stability values

Which one performs better? Persistence or TV/taut string?

- ▶ Experimental evidence in favor of TV/taut string
- ▶ Behavior for fixed n and different functions?
Asymptotic behavior $n \rightarrow \infty$?

Simplifying plane maps



Simplifying plane maps

Any map from an orientable surface to the plane is arbitrarily close to

- ▶ (in C^∞) a map with only fold or cusp singularities
(Whitney)

Simplifying plane maps

Any map from an orientable surface to the plane is arbitrarily close to

- ▶ (in C^∞) a map with only fold or cusp singularities (Whitney)
- ▶ (in C^0) a map with a connected singular curve (Millet)

Simplifying plane maps

Any map from an orientable surface to the plane is arbitrarily close to

- ▶ (in C^∞) a map with only fold or cusp singularities (Whitney)
- ▶ (in C^0) a map with a connected singular curve (Millet)
- ▶ (in C^0) a map with only fold singularities (Eliashberg, Millet)

Simplifying plane maps

Any map from an orientable surface to the plane is arbitrarily close to

- ▶ (in C^∞) a map with only fold or cusp singularities (Whitney)
- ▶ (in C^0) a map with a connected singular curve (Millet)
- ▶ (in C^0) a map with only fold singularities (Eliashberg, Millet)

A possible topological simplification problem:

- ▶ Find a fold map with minimum number of components within a given C^0 tolerance

Simplifying plane maps

Any map from an orientable surface to the plane is arbitrarily close to

- ▶ (in C^∞) a map with only fold or cusp singularities (Whitney)
- ▶ (in C^0) a map with a connected singular curve (Millett)
- ▶ (in C^0) a map with only fold singularities (Eliashberg, Millett)

A possible topological simplification problem:

- ▶ Find a fold map with minimum number of components within a given C^0 tolerance

Fold curves can also be eliminated in pairs (bounding an annulus; ongoing joint work with Zorin)