

Gromov-hyperbolicity, geodesic defect, and persistent homology of Vietoris–Rips complexes

Ulrich Bauer

TUM

Jul 28, 2022

Bavarian Geometry & Topology Meeting X

Universität Augsburg



112



?





112





?













Geometry and topology of biomolecules




Gramicidin (an antibiotic functioning as an ion channel)

Geometry and topology of biomolecules




Gramicidin (an antibiotic functioning as an ion channel)


Čech complexes




Čech complexes




Čech complexes




Čech complexes




Čech complexes



Čech complexes



Čech complexes



Definition

Let X be a topological space, and let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of X .
The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \left\{ J \subseteq I \mid |J| < \infty \text{ and } \bigcap_{i \in J} U_i \neq \emptyset \right\}.$$

Definition

Let X be a topological space, and let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of X .

The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid |J| < \infty \text{ and } \bigcap_{i \in J} U_i \neq \emptyset\}.$$

Theorem (Borsuk 1948, and many more)

Let \mathcal{U} be a good cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Definition

Let X be a topological space, and let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of X .
The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid |J| < \infty \text{ and } \bigcap_{i \in J} U_i \neq \emptyset\}.$$

Theorem (Borsuk 1948, and many more)

Let \mathcal{U} be a good cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Here *good* can mean different things:

- open, numerable cover, contractible intersections
- closed, compact, and convex finite cover

Definition

Let X be a topological space, and let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of X .
The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid |J| < \infty \text{ and } \bigcap_{i \in J} U_i \neq \emptyset\}.$$

Theorem (Borsuk 1948, and many more)

Let \mathcal{U} be a good cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Here *good* can mean different things:

- open, numerable cover, contractible intersections
- closed, compact, and convex finite cover




U. Bauer, M. Kerber, F. Roll, and A. Rolle


A Unified View on the Functorial Nerve Theorem and its Variations

Preprint, arXiv:2203.03571, 2022


Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes



Discrete Morse theory




Discrete Morse theory



Discrete Morse theory

Theorem (Forman 1998)


A simplicial complex with a discrete Morse function f is homotopy equivalent to a CW complex build from the critical simplices off.



Discrete Morse theory

Theorem (Forman 1998)

A simplicial complex with a discrete Morse function f is homotopy equivalent to a CW complex build from the critical simplices off.




Discrete Morse functions – and their gradients – encode collapses of sublevel sets:




Discrete Morse theory

Theorem (Forman 1998)

A simplicial complex with a discrete Morse function f is homotopy equivalent to a CW complex build from the critical simplices off.




Discrete Morse functions – and their gradients – encode collapses of sublevel sets:



Generalizing discrete Morse theory

Generalized discrete Morse functions/gradients:



Morse theory for Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

The Čech complexes and the Delaunay complexes (alpha shapes) are sublevel sets of (generalized) discrete Morse functions.

Morse theory for Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

The Čech complexes and the Delaunay complexes (alpha shapes) are sublevel sets of (generalized) discrete Morse functions.

Theorem (B., Edelsbrunner 2017)

Čech, Delaunay, and Wrap complexes are related by collapses

$$\text{Cech}_r X \searrow \text{Del}_r X \searrow \text{Wrap}_r X,$$


encoded by a single discrete gradient field.



Delaunay and Wrap complexes



Delaunay and Wrap complexes



Homology inference

Inferring homology from samples

Given: finite sample $P \subset X$ of unknown shape $X \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(X)$.*

Inferring homology from samples

Given: finite sample $P \subset X$ of unknown shape $X \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(X)$.*

Problem (Homological reconstruction)

Construct a shape R with $H_(R) \cong H_*(X)$.*

Inferring homology from samples

Given: finite sample $P \subset X$ of unknown shape $X \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(X)$.*

Problem (Homological reconstruction)

Construct a shape R with $H_(R) \cong H_*(X)$.*

Approach:

- approximate X by a thickening $P_\delta = \bigcup_{p \in P} B_\delta(p)$ that covers X

Inferring homology from samples

Given: finite sample $P \subset X$ of unknown shape $X \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(X)$.*

Problem (Homological reconstruction)

Construct a shape R with $H_(R) \cong H_*(X)$.*

Approach:

- approximate X by a thickening $P_\delta = \bigcup_{p \in P} B_\delta(p)$ that covers X

This can work, but requires strong assumptions:

Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.


Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.



Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.



Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.



Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.



Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let X be a submanifold of \mathbb{R}^d . Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X , and
- $\delta < \sqrt{3/20} \text{reach}(X)$.

Then $H_*(X) \cong H_*(P_{2\delta})$.

Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.


Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.



Homology inference using persistence


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.

Proof.



Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.

Proof.

$$\begin{array}{ccc} H_*(X) & H_*(X_\delta) & H_*(X_{2\delta}) \\ \searrow & \nearrow & \searrow \\ H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.

Proof.

$$\begin{array}{ccccc} H_*(X) & \xleftarrow{\quad} & H_*(X_\delta) & \xrightarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & H_*(P_\delta) & & H_*(P_{2\delta}) & \end{array}$$

Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.

Proof.

$$\begin{array}{ccccc} H_*(X) & \xleftarrow{\quad} & H_*(X_\delta) & \xrightarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & H_*(P_\delta) & & H_*(P_{2\delta}) & \end{array}$$

Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.

Proof.

$$\begin{array}{ccccc} H_*(X) & \xleftarrow{\quad} & H_*(X_\delta) & \xrightarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & H_*(P_\delta) & & H_*(P_{2\delta}) & \end{array}$$

Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.

Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.

Proof.

$$\begin{array}{ccccc} H_*(X) & \xleftarrow{\quad} & H_*(X_\delta) & \xrightarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \downarrow \cong & \swarrow \nearrow \\ H_*(P_\delta) & & & H_*(P_{2\delta}) & \\ & \searrow & \downarrow & \swarrow & \nearrow \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $X \subset \mathbb{R}^d$. Let $P \subset X$ and $\delta > 0$ be such that

- P_δ covers X ,
- the inclusions $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$ of thickenings induce isomorphisms in homology.


Then $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$.

Proof.



$$\begin{array}{ccccc} H_*(X) & \xhookleftarrow{\quad \cong \quad} & H_*(X_\delta) & \xhookrightarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \downarrow \cong & \swarrow \nearrow \\ H_*(P_\delta) & & & H_*(P_{2\delta}) & \\ & \searrow & \downarrow & \nearrow & \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

□



What is persistent homology?





What is persistent homology?





What is persistent homology?





What is persistent homology?





What is persistent homology?



What is persistent homology?



What is persistent homology?



What is persistent homology?

What is persistent homology?

Persistent homology is the homology of a filtration.

What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram $K : \mathbf{R} \rightarrow \mathbf{Top}$ of topological spaces, indexed over the poset of real numbers $\mathbf{R} := (\mathbb{R}, \leq)$

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space K_t for each $t \in \mathbb{R}$
- an inclusion map $K_s \hookrightarrow K_t$ for each $s \leq t \in \mathbb{R}$

What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram $K : \mathbf{R} \rightarrow \mathbf{Top}$ of topological spaces, indexed over the poset of real numbers $\mathbf{R} := (\mathbb{R}, \leq)$

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space K_t for each $t \in \mathbb{R}$
- an inclusion map $K_s \hookrightarrow K_t$ for each $s \leq t \in \mathbb{R}$
- Apply homology $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$

What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram $K : \mathbf{R} \rightarrow \mathbf{Top}$ of topological spaces, indexed over the poset of real numbers $\mathbf{R} := (\mathbb{R}, \leq)$

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space K_t for each $t \in \mathbb{R}$
- an inclusion map $K_s \hookrightarrow K_t$ for each $s \leq t \in \mathbb{R}$
- Apply homology $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- Persistent homology is a diagram $M = H_* \circ K : \mathbf{R} \rightarrow \mathbf{Vect}$ (*persistence module*):

$$\dots \rightarrow M_s \longrightarrow M_t \rightarrow \dots$$

Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module $M : \mathbf{R} \rightarrow \mathbf{vect}$ (of finite dim. vector spaces over some field \mathbb{F}) decomposes as a direct sum of interval modules

$$\rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow$$

(in an essentially unique way).

Barcodes: the structure of persistence modules


Theorem (Crawley-Boevey 2015)

Any persistence module $M : \mathbf{R} \rightarrow \mathbf{vect}$ (of finite dim. vector spaces over some field \mathbb{F}) decomposes as a direct sum of interval modules

$$\rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow$$

(in an essentially unique way).

- The supporting intervals form the *persistence barcode*.



Stability


Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $f, g : X \rightarrow \mathbb{R}$ with $\|f - g\|_\infty = \delta$ (and some regularity assumptions).

- Consider the sublevel set filtrations $f^{-1}(\infty, t]$ and $g^{-1}(\infty, t]$, and
- take the resulting persistence barcodes.

Then there exists a δ -matching between the barcodes, meaning that:



Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $f, g : X \rightarrow \mathbb{R}$ with $\|f - g\|_\infty = \delta$ (and some regularity assumptions).

- Consider the sublevel set filtrations $f^{-1}(\infty, t]$ and $g^{-1}(\infty, t]$, and
- take the resulting persistence barcodes.

Then there exists a δ -matching between the barcodes, meaning that:

- matched intervals have endpoints within distance $\leq \delta$, and



Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $f, g : X \rightarrow \mathbb{R}$ with $\|f - g\|_\infty = \delta$ (and some regularity assumptions).

- Consider the sublevel set filtrations $f^{-1}(\infty, t]$ and $g^{-1}(\infty, t]$, and
- take the resulting persistence barcodes.

Then there exists a δ -matching between the barcodes, meaning that:

- matched intervals have endpoints within distance $\leq \delta$, and
- unmatched intervals have length $\leq 2\delta$.




Persistence and stability: the big picture

Data


point cloud

$$P \subset \mathbb{R}^d$$


Persistence and stability: the big picture




Persistence and stability: the big picture




Persistence and stability: the big picture




Persistence and stability: the big picture



Persistence and stability: the big picture



Persistence and stability: the big picture




Interleavings

Let $\delta = \|f - g\|_\infty$. Write $F_t = f^{-1}(-\infty, t]$ for the t -sublevel set of f .

Interleavings

Let $\delta = \|f - g\|_\infty$. Write $F_t = f^{-1}(-\infty, t]$ for the t -sublevel set of f .


Then the sublevel set filtrations $F, G : \mathbf{R} \rightarrow \mathbf{Top}$ are δ -interleaved:



Interleavings


Let $\delta = \|f - g\|_\infty$. Write $F_t = f^{-1}(-\infty, t]$ for the t -sublevel set of f .

Then the sublevel set filtrations $F, G : \mathbf{R} \rightarrow \mathbf{Top}$ are δ -interleaved:



Applying homology, the persistence modules


$H_*(F), H_*(G) : \mathbf{R} \rightarrow \mathbf{Vect}$ are δ -interleaved:



Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

*If two persistence modules are δ -interleaved,
then there exists a δ -matching of their barcodes.*



Induced matchings

For $f : M \rightarrow N$ a morphism (natural transformation) of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$


gives an *induced matching* between their barcodes:

Induced matchings

For $f : M \rightarrow N$ a morphism (natural transformation) of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

gives an *induced matching* between their barcodes:



Induced matchings

For $f : M \rightarrow N$ a morphism (natural transformation) of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

gives an *induced matching* between their barcodes:




Theorem (Induced Matching theorem; B, Lesnick 2015)


If f is a δ -interleaving morphism, then this is a δ -matching.

Persistence of Vietoris–Rips complexes


Vietoris–Rips complexes




Vietoris–Rips complexes




Vietoris–Rips complexes




Vietoris–Rips complexes




Vietoris–Rips complexes




Vietoris–Rips complexes




Vietoris–Rips complexes



Vietoris–Rips complexes



Vietoris–Rips complexes



Vietoris–Rips complexes

For a metric space X , the *Vietoris–Rips complex* at scale $t > 0$ is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$

An example computation

Example data set:

- 192 points on \mathbb{S}^2
- homology up to dimension 2: over 56 mio. simplices in 3-skeleton

An example computation

Example data set:

- 192 points on \mathbb{S}^2
- homology up to dimension 2: over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

An example computation

Example data set:

- 192 points on \mathbb{S}^2
- homology up to dimension 2: over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

Current state of the art:

- Ripser: 0.72 seconds, 83 MB

An example computation

Example data set:


- 192 points on \mathbb{S}^2
- homology up to dimension 2: over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

Current state of the art:

- Ripser: 0.72 seconds, 83 MB
- try live.ripser.org



Apparent pairs:

Ripser uses the following construction for a computational shortcut:

Definition

In a simplexwise filtration $(K_i = \{\sigma_1, \dots, \sigma_i\})_i$, a pair of simplices (σ, τ) is an *apparent pair* if

- σ is the latest proper face of τ , and
- τ is the earliest proper coface of σ .

Apparent pairs:

Ripser uses the following construction for a computational shortcut:

Definition

In a simplexwise filtration $(K_i = \{\sigma_1, \dots, \sigma_i\})_i$, a pair of simplices (σ, τ) is an *apparent pair* if

- σ is the latest proper face of τ , and
- τ is the earliest proper coface of σ .

Proposition

The apparent pairs form a discrete gradient.

Proposition


The apparent pairs form persistence pairs: the cycle $\partial\tau$ gives rise to an interval $[f(\sigma), f(\tau)]$ in the persistence barcode.

The diameter-lexicographic filtration

We use the *lexicographic refinement* of the Vietoris–Rips filtration:


- Choose a total order on the vertices
- Order simplices by diameter
- Order simplices with same diameter by lexicographic vertex order

Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Topology of viral evolution




Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Observation: Ripser runs unusually fast on genetic distance data

- SARS-CoV2 RNA sequences (spike protein)

Topology of viral evolution




Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Observation: Ripser runs unusually fast on genetic distance data

- SARS-CoV2 RNA sequences (spike protein)
- 25556 data points (2.8×10^{12} simplices in 2-skeleton)

Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Observation: Ripser runs unusually fast on genetic distance data

- SARS-CoV2 RNA sequences (spike protein)
- 25556 data points (2.8×10^{12} simplices in 2-skeleton)
- 120 s computation time (with data points ordered appropriately)

The Rips Contractibility Lemma

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.

The Rips Contractibility Lemma

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.


- What about non-geodesic spaces?
- In particular, finite metric spaces?
- Connection to apparent pairs/Ripser?

Gromov-hyperbolicity

Definition (Gromov 1988)

A metric space X is δ -hyperbolic if for all $w, x, y, z \in X$ we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$




Gromov-hyperbolicity

Definition (Gromov 1988)

A metric space X is δ -hyperbolic if for all $w, x, y, z \in X$ we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$



- The hyperbolic plane is $\ln 2$ -hyperbolic




Gromov-hyperbolicity

Definition (Gromov 1988)

A metric space X is δ -hyperbolic if for all $w, x, y, z \in X$ we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$



- The hyperbolic plane is $\ln 2$ -hyperbolic



- 0-hyperbolic spaces are subspaces of trees

Rips contractibility for non-geodesic spaces


Theorem (B, Roll 2022)

Let X be a finite δ -hyperbolic space. Then there is a discrete gradient encoding the collapses


$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow \{\ast\}$$

for all $u > t \geq 4\delta + 2\nu$, where ν is the geodesic defect of X .


Geodesic defect



Geodesic defect



Geodesic defect



Definition (Bonk, Schramm 2000)

A metric space X is ν -geodesic if for all points $x, y \in X$ and all $r, s \geq 0$ with $r + s = d(x, y)$ we have

$$B_{r+\nu}(x) \cap B_{r+\nu}(y) \neq \emptyset.$$

The infimum of all such ν is the *geodesic defect* of X .

Bounds on the geodesic defect

ν : geodesic defect of finite (X, d)

- lower bound:

$$\nu \geq \frac{1}{2} \min_{x \neq y \in X} d(x, y)$$

Bounds on the geodesic defect

ν : geodesic defect of finite (X, d)

- lower bound:

$$\nu \geq \frac{1}{2} \min_{x \neq y \in X} d(x, y)$$

- upper bound:

$$\nu \leq d_H(X, Z)$$

for $X \subseteq Z$ an ambient geodesic space

Bounds on the geodesic defect

ν : geodesic defect of finite (X, d)

- lower bound:

$$\nu \geq \frac{1}{2} \min_{x \neq y \in X} d(x, y)$$

- upper bound:

$$\nu \leq d_H(X, Z)$$

for $X \subseteq Z$ an ambient geodesic space

Bounds on the geodesic defect

ν : geodesic defect of finite (X, d)

- lower bound:

$$\nu \geq \frac{1}{2} \min_{x \neq y \in X} d(x, y)$$

- upper bound:

$$\nu \leq d_H(X, Z)$$

for $X \subseteq Z$ an ambient geodesic space

Possible ambient spaces for X :

- $\ell^\infty(X)$ (functions $X \rightarrow \mathbb{R}$)

Bounds on the geodesic defect

ν : geodesic defect of finite (X, d)

- lower bound:

$$\nu \geq \frac{1}{2} \min_{x \neq y \in X} d(x, y)$$

- upper bound:

$$\nu \leq d_H(X, Z)$$

for $X \subseteq Z$ an ambient geodesic space

Possible ambient spaces for X :

- $\ell^\infty(X)$ (functions $X \rightarrow \mathbb{R}$)
 - Kuratowski embedding $e: X \rightarrow \ell^\infty(X)$, $x \mapsto (d_x = d(x, -)): X \rightarrow \mathbb{R}$

Bounds on the geodesic defect

ν : geodesic defect of finite (X, d)

- lower bound:

$$\nu \geq \frac{1}{2} \min_{x \neq y \in X} d(x, y)$$

- upper bound:

$$\nu \leq d_H(X, Z)$$

for $X \subseteq Z$ an ambient geodesic space

Possible ambient spaces for X :

- $\ell^\infty(X)$ (functions $X \rightarrow \mathbb{R}$)
 - Kuratowski embedding $e: X \rightarrow \ell^\infty(X)$, $x \mapsto (d_x = d(x, -)): X \rightarrow \mathbb{R}$
- injective hull (tight span): subspace of $\ell^\infty(X)$

The tight span of a metric space

Given a metric space (X, d) , the *injective hull* is

$$E(X) = \{f: X \rightarrow \mathbb{R} \mid f(x) + f(y) \geq d(x, y), f(x) = \sup_{y \in X} (d(x, y) - f(y))\},$$

equipped with the metric induced by the sup-norm.

The tight span of a metric space

Given a metric space (X, d) , the *injective hull* is

$$E(X) = \{f: X \rightarrow \mathbb{R} \mid f(x) + f(y) \geq d(x, y), f(x) = \sup_{y \in X} (d(x, y) - f(y))\},$$

equipped with the metric induced by the sup-norm.

Some properties:

- geodesic

The tight span of a metric space

Given a metric space (X, d) , the *injective hull* is

$$E(X) = \{f: X \rightarrow \mathbb{R} \mid f(x) + f(y) \geq d(x, y), f(x) = \sup_{y \in X} (d(x, y) - f(y))\},$$

equipped with the metric induced by the sup-norm.

Some properties:

- geodesic
- hyperconvex (metric balls have the Helly property)

The tight span of a metric space

Given a metric space (X, d) , the *injective hull* is

$$E(X) = \{f: X \rightarrow \mathbb{R} \mid f(x) + f(y) \geq d(x, y), f(x) = \sup_{y \in X} (d(x, y) - f(y))\},$$

equipped with the metric induced by the sup-norm.

Some properties:

- geodesic
- hyperconvex (metric balls have the Helly property)
- contains the Kuratowski embedding $e(X): x \mapsto d_x$

The tight span of a metric space

Given a metric space (X, d) , the *injective hull* is

$$E(X) = \{f: X \rightarrow \mathbb{R} \mid f(x) + f(y) \geq d(x, y), f(x) = \sup_{y \in X} (d(x, y) - f(y))\},$$

equipped with the metric induced by the sup-norm.

Some properties:

- geodesic
- hyperconvex (metric balls have the Helly property)
- contains the Kuratowski embedding $e(X): x \mapsto d_x$
- contractible [Isbell 1962]

Density of Kuratowski embedding

Theorem (Lang 2013 + ϵ)

Let X be a δ -hyperbolic v -geodesic metric space. Then the injective hull $E(X)$ is δ -hyperbolic, and every point in $E(X)$ has distance at most $2\delta + v$ to $e(X)$.

Density of Kuratowski embedding

Theorem (Lang 2013 + ϵ)

Let X be a δ -hyperbolic v -geodesic metric space. Then the injective hull $E(X)$ is δ -hyperbolic, and every point in $E(X)$ has distance at most $2\delta + v$ to $e(X)$.

Corollary (Lim, Mémoli, Okutan 2020 + ϵ)

$\text{Rips}_t(X)$ is contractible for all $t > 4\delta + 2v$.

Density of Kuratowski embedding

Theorem (Lang 2013 + ϵ)

Let X be a δ -hyperbolic ν -geodesic metric space. Then the injective hull $E(X)$ is δ -hyperbolic, and every point in $E(X)$ has distance at most $2\delta + \nu$ to $e(X)$.

Corollary (Lim, Mémoli, Okutan 2020 + ϵ)

$\text{Rips}_t(X)$ is contractible for all $t > 4\delta + 2\nu$.

Proof outline.

- $\text{Rips}_t(X) = \text{Cech}_r(e(X), E(X))$ (hyperconvexity)

Density of Kuratowski embedding

Theorem (Lang 2013 + ϵ)

Let X be a δ -hyperbolic v -geodesic metric space. Then the injective hull $E(X)$ is δ -hyperbolic, and every point in $E(X)$ has distance at most $2\delta + v$ to $e(X)$.

Corollary (Lim, Mémoli, Okutan 2020 + ϵ)

$\text{Rips}_t(X)$ is contractible for all $t > 4\delta + 2v$.

Proof outline.

- $\text{Rips}_t(X) = \text{Cech}_r(e(X), E(X))$ (hyperconvexity)
- For $r > 2\delta + v$: $(B_r(x))_{x \in e(X)}$ is a good cover of $E(X)$ (density)

Density of Kuratowski embedding

Theorem (Lang 2013 + ϵ)

Let X be a δ -hyperbolic v -geodesic metric space. Then the injective hull $E(X)$ is δ -hyperbolic, and every point in $E(X)$ has distance at most $2\delta + v$ to $e(X)$.

Corollary (Lim, Mémoli, Okutan 2020 + ϵ)

$\text{Rips}_t(X)$ is contractible for all $t > 4\delta + 2v$.

Proof outline.

- $\text{Rips}_t(X) = \text{Cech}_r(e(X), E(X))$ (hyperconvexity)
- For $r > 2\delta + v$: $(B_r(x))_{x \in e(X)}$ is a good cover of $E(X)$ (density)
- $\text{Cech}_r(e(X), E(X)) \simeq E(X)$ (nerve theorem)

Density of Kuratowski embedding

Theorem (Lang 2013 + ϵ)

Let X be a δ -hyperbolic v -geodesic metric space. Then the injective hull $E(X)$ is δ -hyperbolic, and every point in $E(X)$ has distance at most $2\delta + v$ to $e(X)$.

Corollary (Lim, Mémoli, Okutan 2020 + ϵ)

$\text{Rips}_t(X)$ is contractible for all $t > 4\delta + 2v$.

Proof outline.

- $\text{Rips}_t(X) = \text{Cech}_r(e(X), E(X))$ (hyperconvexity)
- For $r > 2\delta + v$: $(B_r(x))_{x \in e(X)}$ is a good cover of $E(X)$ (density)
- $\text{Cech}_r(e(X), E(X)) \simeq E(X)$ (nerve theorem)
- $E(X) \simeq \{\ast\}$ (contractibility)

□

Collapsing Vietoris–Rips complexes of generic trees

Consider a *generic* finite metric tree $T = (V, E)$ (distinct distances).

- $\text{diam}: 2^V \rightarrow \mathbb{R}$ is a generalized discrete Morse function.

Collapsing Vietoris–Rips complexes of generic trees

Consider a *generic* finite metric tree $T = (V, E)$ (distinct distances).

- $\text{diam}: 2^V \rightarrow \mathbb{R}$ is a generalized discrete Morse function.
- Its discrete gradient has the non-critical intervals $[e, \Delta_e]$, where

Collapsing Vietoris–Rips complexes of generic trees

Consider a *generic* finite metric tree $T = (V, E)$ (distinct distances).

- $\text{diam}: 2^V \rightarrow \mathbb{R}$ is a generalized discrete Morse function.
- Its discrete gradient has the non-critical intervals $[e, \Delta_e]$, where
 - $e \in \binom{V}{2} \setminus E$ is any non-tree edge

Collapsing Vietoris–Rips complexes of generic trees

Consider a *generic* finite metric tree $T = (V, E)$ (distinct distances).

- $\text{diam}: 2^V \rightarrow \mathbb{R}$ is a generalized discrete Morse function.
- Its discrete gradient has the non-critical intervals $[e, \Delta_e]$, where
 - $e \in \binom{V}{2} \setminus E$ is any non-tree edge
 - Δ_e is the unique maximal cofaces with $\text{diam } \Delta_e = l(e)$

Collapsing Vietoris–Rips complexes of generic trees

Consider a *generic* finite metric tree $T = (V, E)$ (distinct distances).

- $\text{diam}: 2^V \rightarrow \mathbb{R}$ is a generalized discrete Morse function.
- Its discrete gradient has the non-critical intervals $[e, \Delta_e]$, where
 - $e \in \binom{V}{2} \setminus E$ is any non-tree edge
 - Δ_e is the unique maximal cofaces with $\text{diam } \Delta_e = l(e)$
- Only vertices V and edges E are critical

Collapsing Vietoris–Rips complexes of generic trees

Consider a *generic* finite metric tree $T = (V, E)$ (distinct distances).

- $\text{diam}: 2^V \rightarrow \mathbb{R}$ is a generalized discrete Morse function.
- Its discrete gradient has the non-critical intervals $[e, \Delta_e]$, where
 - $e \in \binom{V}{2} \setminus E$ is any non-tree edge
 - Δ_e is the unique maximal cofaces with $\text{diam } \Delta_e = l(e)$
- Only vertices V and edges E are critical

Corollary (B, Roll 2022)

The discrete gradient induces collapses

$$\text{Rips}_t(X) \searrow T_t \quad \text{for all } t \in \mathbb{R}, \text{ with}$$

$$\text{Rips}_t(X) \searrow T \searrow \{\ast\} \quad \text{for all } t \geq \max l(E), \text{ and}$$

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \quad \text{for all } (t, u] \cap l(E) = \emptyset.$$

In particular, the persistent homology is trivial in degrees > 0 .

Arbitrary tree metrics

Example: phylogenetic trees

- non-generic tree metric
- diam is not a generalized discrete Morse function

Arbitrary tree metrics

Example: phylogenetic trees

- non-generic tree metric
- diam is not a generalized discrete Morse function

There is still a compatible gradient:

- independent of choices (*canonical gradient*)
- for generic trees: equals the diam gradient
- only V, E critical
- induces the same collapses

Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order

Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order


This results in a gradient structure like a generic tree metric

Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order

This results in a gradient structure like a generic tree metric




Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order

This results in a gradient structure like a generic tree metric




Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order

This results in a gradient structure like a generic tree metric




Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order

This results in a gradient structure like a generic tree metric




Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order

This results in a gradient structure like a generic tree metric




- The canonical gradient (from the previous slide) refines the perturbed gradient (for any choice of total order)

Symbolic perturbation

Tie breaking for non-distinct pairwise distances:

- Choose total order on vertices
- Order edges lexicographically
- Perturb metric symbolically w.r.t. edge order

This results in a gradient structure like a generic tree metric



- The canonical gradient (from the previous slide) refines the perturbed gradient (for any choice of total order)
- Hence, the perturbed gradient induces the same collapses

Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.

- Choose a root and extend the tree order to a total order.

Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.


- Choose a root and extend the tree order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for the resulting filtration order has critical simplices only on the tree T . It induces the collapses

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow T_t$$

for all $u > t$ such that no tree edge $e \in E$ has length $l(e) \in (t, u]$.



Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.


- Choose a root and extend the tree order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for the resulting filtration order has critical simplices only on the tree T . It induces the collapses

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow T_t$$

for all $u > t$ such that no tree edge $e \in E$ has length $l(e) \in (t, u]$.



Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.


- Choose a root and extend the tree order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for the resulting filtration order has critical simplices only on the tree T . It induces the collapses

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow T_t$$

for all $u > t$ such that no tree edge $e \in E$ has length $l(e) \in (t, u]$.



Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.


- Choose a root and extend the tree order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for the resulting filtration order has critical simplices only on the tree T . It induces the collapses

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow T_t$$

for all $u > t$ such that no tree edge $e \in E$ has length $l(e) \in (t, u]$.



Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.


- Choose a root and extend the tree order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for the resulting filtration order has critical simplices only on the tree T . It induces the collapses

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow T_t$$

for all $u > t$ such that no tree edge $e \in E$ has length $l(e) \in (t, u]$.



Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.


- Choose a root and extend the tree order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for the resulting filtration order has critical simplices only on the tree T . It induces the collapses

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow T_t$$

for all $u > t$ such that no tree edge $e \in E$ has length $l(e) \in (t, u]$.



Collapsing Rips complexes of trees with apparent pairs

Let X be the path length metric space for a weighted tree $T = (X, E)$.


- Choose a root and extend the tree order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for the resulting filtration order has critical simplices only on the tree T . It induces the collapses

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow T_t$$

for all $u > t$ such that no tree edge $e \in E$ has length $l(e) \in (t, u]$.



- Explains why Ripser is very fast on genetic distances (tree-like)



U. Bauer, M. Kerber, F. Roll, and A. Rolle

A Unified View on the Functorial Nerve Theorem and its Variations

Preprint, arXiv:2203.03571, 2022



U. Bauer, H. Edelsbrunner

The Morse Theory of Čech and Delaunay Complexes

Transactions of the AMS, 2017.



U. Bauer, M. Lesnick

Induced matchings and the algebraic stability of persistence barcodes

Journal of Computational Geometry, 2015.



U. Bauer, F. Roll

Gromov hyperbolicity, geodesic defect, and apparent pairs in
Vietoris-Rips filtrations

Symposium on Computational Geometry, 2022.



U. Bauer

Ripser: efficient computation of Vietoris–Rips persistence barcodes

Journal of Applied and Computational Topology, 2021