

# Geometric complexes and topological persistence

## A panoramic view

Ulrich Bauer

Technical University of Munich (TUM)

Aug 1, 2025

TSVP Thematic Program  
TDA PARTI: Topological Data Analysis, Persistence And Representation Theory Intertwined  
OIST, Okinawa

Funded by



Deutsche  
Forschungsgemeinschaft  
German Research Foundation



SFB  
TRR  
109


Discretization  
In Geometry  
and DynamIcs

Technical  
University  
of Munich




mcmL  
Munich Center for Machine Learning

# Biogeometry




Gramicidin (an antibiotic functioning as an ion channel)


# Biogeometry




Gramicidin (an antibiotic functioning as an ion channel)













# Geometric complexes


## Čech complexes




# Čech complexes




# Čech complexes




## Čech complexes



# Čech complexes



# Čech complexes



# Nerves

## Definition (Alexandrov 1928)

Let  $\mathcal{U} = (U_i)_{i \in I}$  be a cover of a space  $X$ . The *nerve* of  $\mathcal{U}$  is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}.$$

# Nerves

## Definition (Alexandrov 1928)

Let  $\mathcal{U} = (U_i)_{i \in I}$  be a cover of a space  $X$ . The *nerve* of  $\mathcal{U}$  is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}.$$

## Theorem (Borsuk 1948, and many more)

Let  $\mathcal{U}$  be a good cover of  $X$ . Then  $\text{Nrv}(\mathcal{U})$  is homotopy equivalent to  $X$ .

# Nerves

## Definition (Alexandrov 1928)

Let  $\mathcal{U} = (U_i)_{i \in I}$  be a cover of a space  $X$ . The *nerve* of  $\mathcal{U}$  is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}.$$

## Theorem (Borsuk 1948, and many more)

Let  $\mathcal{U}$  be a good cover of  $X$ . Then  $\text{Nrv}(\mathcal{U})$  is homotopy equivalent to  $X$ .

Here *good* can mean different things:

- open, numerable cover, contractible intersections
- closed, convex finite cover

# Nerves

## Definition (Alexandrov 1928)

Let  $\mathcal{U} = (U_i)_{i \in I}$  be a cover of a space  $X$ . The *nerve* of  $\mathcal{U}$  is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}.$$

## Theorem (Borsuk 1948, and many more)

Let  $\mathcal{U}$  be a good cover of  $X$ . Then  $\text{Nrv}(\mathcal{U})$  is homotopy equivalent to  $X$ .

Here *good* can mean different things:

- open, numerable cover, contractible intersections
- closed, convex finite cover




U. Bauer, M. Kerber, F. Roll, and A. Rolle


A Unified View on the Functorial Nerve Theorem and its Variations

Expositiones Mathematicae, 2023. doi:10.1016/j.exmath.2023.04.005


## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes



## Delaunay complexes




## Delaunay complexes



# Simplicial collapses

Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.




# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.



# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.
- We can *collapse*  $K$  to a subcomplex  $L$ :



# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.
- We can *collapse*  $K$  to a subcomplex  $L$ :
  - remove a free face and its proper coface.



# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.
- We can *collapse*  $K$  to a subcomplex  $L$ :
  - remove a free face and its proper coface.
  - $L$  is homotopy equivalent to  $K$ .




# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.
- We can *collapse*  $K$  to a subcomplex  $L$ :
  - remove a free face and its proper coface.
  - $L$  is homotopy equivalent to  $K$ .

If there is a sequence of such elementary collapses from  $K$  to  $M$ , we say that  $K$  *collapses* to  $M$  (written as  $K \searrow M$ ).




# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .




# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .



A function  $f : K \rightarrow \mathbb{R}$  on a cell complex is a *discrete Morse function* if

- the sublevel sets  $f^{-1}(-\infty, t]$  are subcomplexes, and


•<sub>1</sub>

# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .



A function  $f : K \rightarrow \mathbb{R}$  on a cell complex is a *discrete Morse function* if

- the sublevel sets  $f^{-1}(-\infty, t]$  are subcomplexes, and




# Discrete Morse theory

## Definition (Forman 1998)


A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .



A function  $f : K \rightarrow \mathbb{R}$  on a cell complex is a *discrete Morse function* if

- the sublevel sets  $f^{-1}(-\infty, t]$  are subcomplexes, and



# Discrete Morse theory



## Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .

A function  $f : K \rightarrow \mathbb{R}$  on a cell complex is a *discrete Morse function* if

- the sublevel sets  $f^{-1}(-\infty, t]$  are subcomplexes, and



# Discrete Morse theory



## Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .

A function  $f : K \rightarrow \mathbb{R}$  on a cell complex is a *discrete Morse function* if

- the sublevel sets  $f^{-1}(-\infty, t]$  are subcomplexes, and



# Discrete Morse theory



## Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .

A function  $f : K \rightarrow \mathbb{R}$  on a cell complex is a *discrete Morse function* if

- the sublevel sets  $f^{-1}(-\infty, t]$  are subcomplexes, and




# Discrete Morse theory

## Definition (Forman 1998)


A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets  $\{\phi\}$  (*critical cells*), and
- pairs  $\{\sigma, \tau\}$ , where  $\sigma$  is a facet of  $\tau$ .




A function  $f : K \rightarrow \mathbb{R}$  on a cell complex is a *discrete Morse function* if


- the sublevel sets  $f^{-1}(-\infty, t]$  are subcomplexes, and
- the level sets  $f^{-1}(t)$  form a discrete vector field.



# Fundamental theorem of discrete Morse theory




# Fundamental theorem of discrete Morse theory



# Fundamental theorem of discrete Morse theory

## Theorem (Forman 1998)


A simplicial complex with a discrete Morse function  $f$  is homotopy equivalent to a space (a CW complex) built from the critical simplices off.




# Fundamental theorem of discrete Morse theory

## Theorem (Forman 1998)

A simplicial complex with a discrete Morse function  $f$  is homotopy equivalent to a space (a CW complex) built from the critical simplices off.




Discrete Morse functions – and their gradients – encode collapses of sublevel sets:




# Fundamental theorem of discrete Morse theory

## Theorem (Forman 1998)

A simplicial complex with a discrete Morse function  $f$  is homotopy equivalent to a space (a CW complex) built from the critical simplices off.



Discrete Morse functions – and their gradients – encode collapses of sublevel sets:




# Generalized discrete Morse theory

## Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex  $K$  is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$

- indicated by an arrow from  $L$  to  $U$



# Generalized discrete Morse theory

## Definition (Chari 2000, Freij 2009)



A *generalized discrete vector field* on a simplicial complex  $K$  is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$

- indicated by an arrow from  $L$  to  $U$

A *generalized discrete Morse function*  $f : K \rightarrow \mathbb{R}$  satisfies:

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes (for all  $t \in \mathbb{R}$ )
- the level sets  $f^{-1}(t)$  form a generalized vector field (the *discrete gradient* of  $f$ )



# Morse theory for Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech complexes and the Delaunay complexes (alpha shapes) are sublevel sets of (generalized) discrete Morse functions.*

# Morse theory for Čech and Delaunay complexes


Proposition (B, Edelsbrunner 2014)

*The Čech complexes and the Delaunay complexes (alpha shapes) are sublevel sets of (generalized) discrete Morse functions.*

Theorem (B, Edelsbrunner 2017)


*Čech, Delaunay, and Wrap complexes (at any scale  $r$ ) of a point set  $X \subset \mathbb{R}^d$  in general position are related by collapses encoded by a single discrete gradient field:*

$$\text{Cech}_r X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$




## Wrap complexes

Foundation of the surface reconstruction software *Wrap* (Edelsbrunner 1995, Geomagic)




## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .



## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .



**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**


$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ :

- the smallest subcomplex of  $\text{Del}_r X$  that
  - contains all critical simplices
  - is compatible with  $V$  (a union of intervals of  $V$ ).

## Delaunay and Wrap complexes




## Delaunay and Wrap complexes




# Persistent homology


## Persistence barcodes




## Persistence barcodes




## Persistence barcodes




## Persistence barcodes




## Persistence barcodes



## Persistence barcodes



## Persistence barcodes



# Computing persistent homology via matrix reduction

Algorithm (matrix reduction; a variant of Gauss elimination)

**Require:**  $D: m \times n$  matrix

**Ensure:**  $V$  is full rank upper triangular,  $R = D \cdot V$  has unique column pivots

**function** Reduce( $D$ )

$R = D$

$V = I(n)$

**while** there exist  $i < j$  such that pivot  $R_i =$  pivot  $R_j$  **do**

        add column  $R_i$  to column  $R_j$     ▷ eliminate the nonzero entry in row pivot  $R_i$

        add column  $V_i$  to column  $V_j$

**return**  $R, V$

# Computing persistent homology via matrix reduction

Algorithm (exhaustive matrix reduction; a variant of Gauss–Jordan elimination)

**Require:**  $D: m \times n$  matrix

**Ensure:**  $V$  is full rank upper triangular,  $R = D \cdot V$  has unique column pivots

**function** Reduce( $D$ )

$R = D$

$V = I(n)$

**while** there exist  $i < j$  such that  $R_j$  has a nonzero entry in row pivot  $R_i$  **do**

add column  $R_i$  to column  $R_j$  ▷ eliminate the nonzero entry in row pivot  $R_i$   
add column  $V_i$  to column  $V_j$

**return**  $R, V$

# Computing persistent homology via matrix reduction

Algorithm (**exhaustive** matrix reduction; a variant of Gauss–Jordan elimination)

**Require:**  $D: m \times n$  matrix

**Ensure:**  $V$  is full rank upper triangular,  $R = D \cdot V$  has unique column pivots

**function** Reduce( $D$ )

$R = D$

$V = I(n)$

**while** there exist  $i < j$  such that  $R_j$  has a nonzero entry in row pivot  $R_i$  **do**

    add column  $R_i$  to column  $R_j$     ▷ eliminate the nonzero entry in row pivot  $R_i$


    add column  $V_i$  to column  $V_j$

**return**  $R, V$


## Proposition

The resulting columns  $R_j$  are minimal (in a lexicographic order) within their homology class (in  $K_{j-1}$ ).


## Wrap complexes and lexicographically minimal cycles




# Wrap complexes and lexicographically minimal cycles




# Wrap complexes and lexicographically minimal cycles



# Wrap complexes and lexicographically minimal cycles




# Wrap complexes and lexicographically minimal cycles




# Wrap complexes and lexicographically minimal cycles



## Wrap complexes and lexicographically minimal cycles




## Wrap complexes and lexicographically minimal cycles






## Standard reduction and exhaustive reduction



## Standard reduction and exhaustive reduction



## Point cloud reconstruction with minimal cycles




# Wrap complexes support minimal cycles

## Theorem (B, Roll 2024)

Let  $X \subset \mathbb{R}$  be a finite subset in general position and let  $r \in \mathbb{R}$ .

- Exhaustive matrix reduction computes the minimal cycles homologous to a simplex boundary.
- Any lexicographically minimal cycle of  $\text{Del}_r(X)$  is supported on  $\text{Wrap}_r(X)$ .




# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:

# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:


- Forman (1998) defines a flow  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)



# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:


- Forman (1998) defines a flow  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)



# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:


- Forman (1998) defines a flow  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)



# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:


- Forman (1998) defines a flow  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)



# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:

- Forman (1998) defines a *flow*  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)




- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)

# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:

- Forman (1998) defines a *flow*  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)




- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)

We interpret persistent homology in terms of algebraic Morse theory:

# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:

- Forman (1998) defines a *flow*  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)



- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)


We interpret persistent homology in terms of algebraic Morse theory:

- Persistence pairs correspond to algebraic gradient pairs

# Algebraic gradient flows and persistent homology

Ingredients for relating the two approaches:

- Forman (1998) defines a *flow*  $C_*(K) \rightarrow C_*(K)$  associated to a gradient (discrete Morse theory)



- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)

We interpret persistent homology in terms of algebraic Morse theory:


- Persistence pairs correspond to algebraic gradient pairs
- Matrix reduction (exhaustive) corresponds to gradient flow

# Vietoris–Rips persistence

## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



# Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite, } \text{diam } S \leq t\}.$$



## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite, } \text{diam } S \leq t\}.$$



## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



## Vietoris–Rips complexes



For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



# Ripser: software for computing Vietoris–Rips persistence barcodes



Open source software ([ripser.org](http://ripser.org))



Ripser users across the globe

# Ripser: software for computing Vietoris–Rips persistence barcodes

Open source software ([ripser.org](http://ripser.org))



Ripser users across the globe

Computational improvements based on

- *implicit matrix representations*
- *apparent pairs*, connecting persistence to discrete Morse theory

# Apparent pairs

Ripser uses the following pairing of simplices (breaking ties in the filtration lexicographically):

## Definition (B 2016, 2021)

In a simplexwise filtration ( $K_i = \{\sigma_1, \dots, \sigma_i\}_i$ ), two simplices  $(\sigma_i, \sigma_j)$  form an *apparent pair* if

- $\sigma_i$  is the latest proper face of  $\sigma_j$ , and
- $\sigma_j$  is the earliest proper coface of  $\sigma_i$ .

## Apparent pairs

Ripser uses the following pairing of simplices (breaking ties in the filtration lexicographically):

### Definition (B 2016, 2021)

In a simplexwise filtration ( $K_i = \{\sigma_1, \dots, \sigma_i\}_i$ ), two simplices  $(\sigma_i, \sigma_j)$  form an *apparent pair* if

- $\sigma_i$  is the latest proper face of  $\sigma_j$ , and
- $\sigma_j$  is the earliest proper coface of  $\sigma_i$ .

Special cases and equivalent definitions have been (re)discovered independently multiple times

- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

# Apparent pairs

Ripser uses the following pairing of simplices (breaking ties in the filtration lexicographically):

## Definition (B 2016, 2021)

In a simplexwise filtration ( $K_i = \{\sigma_1, \dots, \sigma_i\}_i$ ), two simplices  $(\sigma_i, \sigma_j)$  form an *apparent pair* if

- $\sigma_i$  is the latest proper face of  $\sigma_j$ , and
- $\sigma_j$  is the earliest proper coface of  $\sigma_i$ .

Special cases and equivalent definitions have been (re)discovered independently multiple times


- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

## Proposition (B 2021)


*The apparent pairs are both*

- *persistence pairs (creating/destroying a feature in homology) and*
- *gradient pairs (in the sense of discrete Morse theory).*

## Apparent pairs of the diameter-lexicographic filtration




# Apparent pairs of the diameter-lexicographic filtration



$$\partial_1 = \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} \begin{matrix} (a) \\ (b) \\ (c) \\ (d) \end{matrix}$$

The matrix has 5 columns corresponding to the vertices a, b, c, d, and a row vector. The first column has a circled '1' in the top-left position. The second column has a circled '1' in the middle position. The third column has a circled '1' in the middle position. The fourth column has a circled '1' in the bottom-right position. The fifth column has a circled '1' in the bottom-left position.


# Apparent pairs of the diameter-lexicographic filtration



$$\partial_1 = \begin{pmatrix} 1 & & (a,b):3 & & \\ & 1 & (c,d):3 & & \\ & & 1 & (a,d):4 & \\ & & & 1 & (b,c):5 \\ 1 & & & & 1 \\ & 1 & & & 1 \\ & & 1 & (a,c):5 & \\ & & & 1 & (b,d):5 \\ & & & & 1 \end{pmatrix} \begin{matrix} (a) \\ (b) \\ (c) \\ (d) \end{matrix}$$

$$\partial_2 = \begin{pmatrix} (a,b,c):5 & & & & \\ & (a,b,d):5 & & & \\ & & (a,c,d):5 & & \\ & & & (b,c,d):5 & \\ 1 & 1 & 1 & 1 & (a,b):3 \\ & 1 & 1 & 1 & (c,d):3 \\ 1 & & 1 & 1 & (a,d):4 \\ & 1 & & 1 & (b,c):4 \\ & & 1 & 1 & (a,c):5 \\ & & & 1 & (b,d):5 \end{pmatrix}$$

# Apparent pairs of the diameter-lexicographic filtration



$$\partial_1 = \begin{pmatrix} & (a,b):3 & (c,d):3 & & \\ & 1 & & 1 & (a,d):4 \\ & & 1 & & (b,c):5 \\ 1 & & & 1 & (a,c):5 \\ & 1 & & 1 & (b,d):5 \\ & & 1 & & \\ & & & 1 & \end{pmatrix} \begin{matrix} (a) \\ (b) \\ (c) \\ (d) \end{matrix}$$

$$\partial_2 = \begin{pmatrix} & (a,b,c):5 & (a,b,d):5 & (a,c,d):5 & (b,c,d):5 \\ & 1 & 1 & 1 & 1 \\ & & 1 & 1 & 1 \\ 1 & & & 1 & 1 \\ & 1 & & 1 & 1 \\ & & 1 & & 1 \\ & & & 1 & \end{pmatrix} \begin{matrix} (a,b):3 \\ (c,d):3 \\ (a,d):4 \\ (b,c):4 \\ (a,c):5 \\ (b,d):5 \end{matrix}$$

$$\partial_3 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} (a,b,c) \\ (a,b,d) \\ (a,c,d) \\ (b,c,d) \end{matrix}$$

# A shortcut for finding pivots

$$\begin{array}{c} (a,b,c) \\ \left( \begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array} \right) \begin{array}{l} (a,b) \\ (a,c) \\ (a,d) \\ (b,c) \\ (b,d) \\ (c,d) \end{array} \end{array} \quad \begin{array}{c} (a,b,c):5 \\ \left( \begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array} \right) \begin{array}{l} (a,b):3 \\ (c,d):3 \\ (a,d):4 \\ (b,c):4 \\ (a,c):5 \\ (b,d):5 \end{array} \end{array}$$

Shortcut for finding the pivot (latest) facet of a simplex  $\tau$ :


# A shortcut for finding pivots

$$\begin{array}{c} (a,b,c) \\ \left( \begin{array}{l} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array} \right) \begin{array}{l} (a,b) \\ (a,c) \\ (a,d) \\ (b,c) \\ (b,d) \\ (c,d) \end{array} \end{array} \xrightarrow{\hspace{1cm}} \begin{array}{c} (a,b,c):5 \\ \left( \begin{array}{l} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array} \right) \begin{array}{l} (a,b):3 \\ (c,d):3 \\ (a,d):4 \\ (b,c):4 \\ (a,c):5 \\ (b,d):5 \end{array} \end{array}$$

Shortcut for finding the pivot (latest) facet of a simplex  $\tau$ :

- Enumerate facets  $\sigma$  of  $\tau$  in (reverse) lexicographic order


# A shortcut for finding pivots



Shortcut for finding the pivot (latest) facet of a simplex  $\tau$ :

- Enumerate facets  $\sigma$  of  $\tau$  in (reverse) lexicographic order
- The first facet  $\sigma$  with  $\text{diam } \sigma = \text{diam } \tau$  is the pivot of  $\tau$

# A shortcut for finding pivots



Shortcut for finding the pivot (latest) facet of a simplex  $\tau$ :


- Enumerate facets  $\sigma$  of  $\tau$  in (reverse) lexicographic order
- The first facet  $\sigma$  with  $\text{diam } \sigma = \text{diam } \tau$  is the pivot of  $\tau$

# Gromov-hyperbolicity

## Definition (Gromov 1988)

A metric space  $X$  is  $\delta$ -hyperbolic (for  $\delta \geq 0$ ) if for all  $w, x, y, z \in X$  we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$




# Gromov-hyperbolicity

## Definition (Gromov 1988)

A metric space  $X$  is  $\delta$ -hyperbolic (for  $\delta \geq 0$ ) if for all  $w, x, y, z \in X$  we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$

- The hyperbolic plane is  $(\ln 2)$ -hyperbolic.



# Gromov-hyperbolicity

## Definition (Gromov 1988)



A metric space  $X$  is  $\delta$ -hyperbolic (for  $\delta \geq 0$ ) if for all  $w, x, y, z \in X$  we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$


- The hyperbolic plane is  $(\ln 2)$ -hyperbolic.



- Metric trees and their subspaces are precisely the 0-hyperbolic spaces.




# Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

# Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Observation: Ripser runs unusually fast on genetic distance data

- SARS-CoV2 RNA sequences (spike protein)
- 25556 data points ( $2.8 \times 10^{12}$  simplices in 2-skeleton)
- 120 s computation time (with data points ordered appropriately)

# Rips complexes of hyperbolic spaces

## Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

# Rips complexes of hyperbolic spaces

## Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

- Generalization to non-geodesic spaces? In particular, finite metric spaces?

# Rips complexes of hyperbolic spaces

## Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

- Generalization to non-geodesic spaces? In particular, finite metric spaces?
- Collapsibility? By a discrete gradient compatible with the filtration?

# Rips complexes of hyperbolic spaces

## Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

- Generalization to non-geodesic spaces? In particular, finite metric spaces?
- Collapsibility? By a discrete gradient compatible with the filtration?
- Connection to computation of persistent homology?

# Rips complexes of hyperbolic spaces

## Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

- Generalization to non-geodesic spaces? In particular, finite metric spaces?
- Collapsibility? By a discrete gradient compatible with the filtration?
- Connection to computation of persistent homology?


## Theorem (B, Roll 2022)

*Let  $X$  be a finite  $\delta$ -hyperbolic space. Then there is a single discrete gradient encoding the collapses*


$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow \{\ast\}$$

*for all  $u > t \geq 4\delta + 2\nu$ , where  $\nu$  is the geodesic defect of  $X$ .*


## Geodesic defect




## Geodesic defect



## Geodesic defect



# Geodesic defect



## Definition (Bonk, Schramm 2000)

A metric space  $X$  is  $\nu$ -geodesic if for all points  $x, y \in X$  and all  $r, s \geq 0$  with  $r + s = d(x, y)$  we have

$$B_{r+\nu}(x) \cap B_{s+\nu}(y) \neq \emptyset.$$

The infimum of all such  $\nu$  is the *geodesic defect* of  $X$ .

# The diameter function of generic trees

Proposition (B, Roll 2022)

*Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.*

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The critical simplices are the tree simplices,  $V \cup E$ .

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The critical simplices are the tree simplices,  $V \cup E$ .
- The apparent pairs (lexicographic refinement, based on arbitrary vertex order) refine this gradient.

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The critical simplices are the tree simplices,  $V \cup E$ .
- The apparent pairs (lexicographic refinement, based on arbitrary vertex order) refine this gradient.

## Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space  $X$  induces the collapses

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The critical simplices are the tree simplices,  $V \cup E$ .
- The apparent pairs (lexicographic refinement, based on arbitrary vertex order) refine this gradient.

## Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space  $X$  induces the collapses

- $\text{Rips}_t(X) \searrow T_t$  for all  $t \in \mathbb{R}$ ,

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The critical simplices are the tree simplices,  $V \cup E$ .
- The apparent pairs (lexicographic refinement, based on arbitrary vertex order) refine this gradient.

## Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space  $X$  induces the collapses

- $\text{Rips}_t(X) \searrow T_t$  for all  $t \in \mathbb{R}$ ,
- $\text{Rips}_t(X) \searrow T \searrow \{\ast\}$  for  $t \geq \max d(E)$ , and

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The critical simplices are the tree simplices,  $V \cup E$ .
- The apparent pairs (lexicographic refinement, based on arbitrary vertex order) refine this gradient.

## Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space  $X$  induces the collapses

- $\text{Rips}_t(X) \searrow T_t$  for all  $t \in \mathbb{R}$ ,
- $\text{Rips}_t(X) \searrow T \searrow \{\ast\}$  for  $t \geq \max d(E)$ , and
- $\text{Rips}_u(X) \searrow \text{Rips}_t(X)$  whenever no pairwise distance lies in the interval  $(t, u]$ .

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The critical simplices are the tree simplices,  $V \cup E$ .
- The apparent pairs (lexicographic refinement, based on arbitrary vertex order) refine this gradient.

## Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space  $X$  induces the collapses

- $\text{Rips}_t(X) \searrow T_t$  for all  $t \in \mathbb{R}$ ,
- $\text{Rips}_t(X) \searrow T \searrow \{\ast\}$  for  $t \geq \max d(E)$ , and
- $\text{Rips}_u(X) \searrow \text{Rips}_t(X)$  whenever no pairwise distance lies in the interval  $(t, u]$ .

In particular, the persistent homology is trivial in degrees  $> 0$ .

## Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

## Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .

## Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.


# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case.*




# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case.*




# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case.*




# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case.*




# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case.*




# Stability

## Stability of persistence barcodes

If two point clouds are close, then their barcodes are also close:

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X, Y \subset \mathbb{R}^d$  be finite point sets with Hausdorff distance  $d_H(X, Y) = \delta$ .




# Stability of persistence barcodes

If two point clouds are close, then their barcodes are also close:

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X, Y \subset \mathbb{R}^d$  be finite point sets with Hausdorff distance  $d_H(X, Y) = \delta$ .

Then there exists a  $\delta$ -matching between the intervals in the ( $\check{\text{C}}$ ech or Delaunay) persistence barcodes for  $X$  and  $Y$ :



# Stability of persistence barcodes



If two point clouds are close, then their barcodes are also close:

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X, Y \subset \mathbb{R}^d$  be finite point sets with Hausdorff distance  $d_H(X, Y) = \delta$ .

Then there exists a  $\delta$ -matching between the intervals in the ( $\check{\text{C}}$ ech or Delaunay) persistence barcodes for  $X$  and  $Y$ :

- matched intervals have endpoints within distance  $\leq \delta$ , and



# Stability of persistence barcodes



If two point clouds are close, then their barcodes are also close:

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X, Y \subset \mathbb{R}^d$  be finite point sets with Hausdorff distance  $d_H(X, Y) = \delta$ .

Then there exists a  $\delta$ -matching between the intervals in the ( $\check{\text{C}}$ ech or Delaunay) persistence barcodes for  $X$  and  $Y$ :

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .




# Persistence and stability: the big picture

Data


point cloud

$$P \subset \mathbb{R}^d$$


# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




## Geometric interleavings




## Geometric interleavings




## Geometric interleavings



## Geometric interleavings



## Geometric interleavings




## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ , and similarly  $G_t = g^{-1}(-\infty, t]$ .

# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ , and similarly  $G_t = g^{-1}(-\infty, t]$ .


Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:




# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ , and similarly  $G_t = g^{-1}(-\infty, t]$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:




Applying homology, the persistence modules  $H_*(F), H_*(G) : \mathbf{R} \rightarrow \mathbf{Vect}$  are  $\delta$ -interleaved:



# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2016; B, Lesnick 2015)

*If two persistence modules are  $\delta$ -interleaved,  
then their barcodes admit a  $\delta$ -matching:*




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2016; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then their barcodes admit a  $\delta$ -matching:

- matched intervals have endpoints within distance  $\leq \delta$ , and




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2016; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then their barcodes admit a  $\delta$ -matching:

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .



# Structure of persistence sub-/quotient modules

## Proposition (B, Lesnick 2015)

Let  $M \twoheadrightarrow N$  be an epimorphism of persistence modules.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that

if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial (compatible with composition).



# Structure of persistence sub-/quotient modules

## Proposition (B, Lesnick 2015)

Let  $M \twoheadrightarrow N$  be an epimorphism of persistence modules.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that

if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial (compatible with composition).

Dually, for a monomorphism  $N \hookrightarrow O$  there is an injection  $B(N) \hookrightarrow B(O)$ .



## Induced matchings

Any morphism of pfd persistence modules  $f : M \rightarrow N$  factors through its image as an epimorphism followed by a monomorphism:

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

This gives an *induced matching* between their barcodes:

## Induced matchings

Any morphism of pfd persistence modules  $f : M \rightarrow N$  factors through its image as an epimorphism followed by a monomorphism:

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

This gives an *induced matching* between their barcodes:

- compose the injections  $B(M) \hookleftarrow B(\text{im } f) \hookrightarrow B(N)$  from before to a matching  $B(M) \leftrightarrow B(N)$ .



## Induced matchings

Any morphism of pfd persistence modules  $f : M \rightarrow N$  factors through its image as an epimorphism followed by a monomorphism:

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

This gives an *induced matching* between their barcodes:

- compose the injections  $B(M) \hookleftarrow B(\text{im } f) \hookrightarrow B(N)$  from before to a matching  $B(M) \leftrightarrow B(N)$ .



Warning:

- The corresponding morphism of persistence modules may not be isomorphic to  $f$

# The induced matching theorem



# The induced matching theorem



# The induced matching theorem



# The induced matching theorem




# The induced matching theorem



# The induced matching theorem

Theorem (B, Lesnick, 2015, 2020)


If  $f : M \rightarrow N$  has  $\delta$ -small (co)kernel, then so does the induced matching  $\chi(f) : B(M) \rightarrow B(N)$ .



# The induced matching theorem

Theorem (B, Lesnick, 2015, 2020)

If  $f : M \rightarrow N$  has  $\delta$ -small (co)kernel, then so does the induced matching  $\chi(f) : B(M) \rightarrow B(N)$ .




Corollary

If  $f : M \rightarrow N(\delta)$  is a  $\delta$ -interleaving morphism, then this induced matching is a  $\delta$ -matching.


# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $t \in \mathbb{R}$ ):



# Algebraic stability via induced matchings


Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $t \in \mathbb{R}$ ):



Construct an induced  $\delta$ -matching of barcodes:

# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $t \in \mathbb{R}$ ):



Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  of length  $> 2\delta$  to a similar interval in  $B(M)$



$B(N)$

# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \gg & \text{im } N_{t-\delta, t+\delta} & \longleftarrow \ll & N_{t+\delta} \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  of length  $> 2\delta$  to a similar interval in  $B(M)$



# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \gg & \text{im } N_{t-\delta, t+\delta} & \longleftarrow \ll & N_{t+\delta} \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  of length  $> 2\delta$  to a similar interval in  $B(M)$




# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \twoheadrightarrow & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow & N_{t+\delta} \\ & & \curvearrowright & & \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  of length  $> 2\delta$  to a similar interval in  $B(M)$




# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \twoheadrightarrow & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow & N_{t+\delta} \\ & & \nearrow & \searrow & \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  of length  $> 2\delta$  to a similar interval in  $B(M)$



# Instability

## Indecomposables abound in multi-parameter persistence

Theorem (B, Scoccola 2022)

*For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.*

# Indecomposables abound in multi-parameter persistence

Theorem (B, Scoccola 2022)

*For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.*

- *Given a persistence module  $M$ , any neighborhood contains an indecomposable:*

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

*For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.*

- *Given a persistence module  $M$ , any neighborhood contains an indecomposable:*
  - *For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .*

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

*For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.*

- *Given a persistence module  $M$ , any neighborhood contains an indecomposable:*
  - *For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .*
- *Any indecomposable  $N$  has a neighborhood of almost indecomposables:*

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.

- Given a persistence module  $M$ , any neighborhood contains an indecomposable:
  - For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .
- Any indecomposable  $N$  has a neighborhood of almost indecomposables:
  - For every  $\epsilon > 0$ , there is  $\delta > 0$  such that any persistence module  $M$  within distance  $\delta$  ( $d_I(M, N) < \epsilon$ ) decomposes into

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.

- Given a persistence module  $M$ , any neighborhood contains an indecomposable:
  - For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .
- Any indecomposable  $N$  has a neighborhood of almost indecomposables:
  - For every  $\epsilon > 0$ , there is  $\delta > 0$  such that any persistence module  $M$  within distance  $\delta$  ( $d_I(M, N) < \epsilon$ ) decomposes into
    - an indecomposable summand and

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.

- Given a persistence module  $M$ , any neighborhood contains an indecomposable:
  - For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .
- Any indecomposable  $N$  has a neighborhood of almost indecomposables:
  - For every  $\epsilon > 0$ , there is  $\delta > 0$  such that any persistence module  $M$  within distance  $\delta$  ( $d_I(M, N) < \epsilon$ ) decomposes into
    - an indecomposable summand and
    - an  $\epsilon$ -small summand (distance  $< \epsilon$  to zero module).

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.

- Given a persistence module  $M$ , any neighborhood contains an indecomposable:
  - For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .
- Any indecomposable  $N$  has a neighborhood of almost indecomposables:
  - For every  $\epsilon > 0$ , there is  $\delta > 0$  such that any persistence module  $M$  within distance  $\delta$  ( $d_I(M, N) < \epsilon$ ) decomposes into
    - an indecomposable summand and
    - an  $\epsilon$ -small summand (distance  $< \epsilon$  to zero module).

In other words:

- Indecomposables form a dense subset.

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.

- Given a persistence module  $M$ , any neighborhood contains an indecomposable:
  - For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .
- Any indecomposable  $N$  has a neighborhood of almost indecomposables:
  - For every  $\epsilon > 0$ , there is  $\delta > 0$  such that any persistence module  $M$  within distance  $\delta$  ( $d_I(M, N) < \epsilon$ ) decomposes into
    - an indecomposable summand and
    - an  $\epsilon$ -small summand (distance  $< \epsilon$  to zero module).

In other words:

- Indecomposables form a dense subset.
- $\epsilon$ -almost indecomposables form an open and dense subset.

# Indecomposables abound in multi-parameter persistence

## Theorem (B, Scoccola 2022)

For  $n > 1$ , almost every finitely presentable  $n$ -parameter persistence module is indecomposable.

- Given a persistence module  $M$ , any neighborhood contains an indecomposable:
  - For every  $\epsilon > 0$ , there is an indecomposable  $N$  with  $d_I(M, N) < \epsilon$ .
- Any indecomposable  $N$  has a neighborhood of almost indecomposables:
  - For every  $\epsilon > 0$ , there is  $\delta > 0$  such that any persistence module  $M$  within distance  $\delta$  ( $d_I(M, N) < \epsilon$ ) decomposes into
    - an indecomposable summand and
    - an  $\epsilon$ -small summand (distance  $< \epsilon$  to zero module).

In other words:

- Indecomposables form a dense subset.
- $\epsilon$ -almost indecomposables form an open and dense subset.
- Indecomposability is generic (in the closure of the finitely presented persistence modules).

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

- An  $\delta$ -*matching* between  $M$  and  $N$  consists of decompositions  $M \cong \bigoplus_{j \in J} M_j$  and  $N \cong \bigoplus_{j \in J} N_j$

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

- An  $\delta$ -*matching* between  $M$  and  $N$  consists of decompositions  $M \cong \bigoplus_{j \in J} M_j$  and  $N \cong \bigoplus_{j \in J} N_j$ 
  - indexed by the same set  $J$ ,

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

- An  $\delta$ -*matching* between  $M$  and  $N$  consists of decompositions  $M \cong \bigoplus_{j \in J} M_j$  and  $N \cong \bigoplus_{j \in J} N_j$ 
  - indexed by the same set  $J$ ,
  - where each summand is either indecomposable or zero, and

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

- An  $\delta$ -*matching* between  $M$  and  $N$  consists of decompositions  $M \cong \bigoplus_{j \in J} M_j$  and  $N \cong \bigoplus_{j \in J} N_j$ 
  - indexed by the same set  $J$ ,
  - where each summand is either indecomposable or zero, and
  - such that  $M_j$  and  $N_j$  are  $\delta$ -interleaved for every  $j \in J$ .

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

- An  $\delta$ -*matching* between  $M$  and  $N$  consists of decompositions  $M \cong \bigoplus_{j \in J} M_j$  and  $N \cong \bigoplus_{j \in J} N_j$ 
  - indexed by the same set  $J$ ,
  - where each summand is either indecomposable or zero, and
  - such that  $M_j$  and  $N_j$  are  $\delta$ -interleaved for every  $j \in J$ .
- The *bottleneck distance* between  $M$  and  $N$  is

$$d_B(M, N) = \inf\{\delta \geq 0 \mid \text{there exists a } \delta\text{-matching between } M \text{ and } N\}.$$

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

- An  $\delta$ -*matching* between  $M$  and  $N$  consists of decompositions  $M \cong \bigoplus_{j \in J} M_j$  and  $N \cong \bigoplus_{j \in J} N_j$ 
  - indexed by the same set  $J$ ,
  - where each summand is either indecomposable or zero, and
  - such that  $M_j$  and  $N_j$  are  $\delta$ -interleaved for every  $j \in J$ .
- The *bottleneck distance* between  $M$  and  $N$  is

$$d_B(M, N) = \inf\{\delta \geq 0 \mid \text{there exists a } \delta\text{-matching between } M \text{ and } N\}.$$

- $M$  is *structurally stable* if passing from interleaving to bottleneck distance is continuous at  $M$ .

# Bottleneck distance and structural stability

The definition of bottleneck (or matching) distance extends to multi-parameter persistence.

## Definition

Let  $\delta \geq 0$ , and let  $M, N: \mathbf{R}^n \rightarrow \mathbf{vect}$ .

- An  $\delta$ -*matching* between  $M$  and  $N$  consists of decompositions  $M \cong \bigoplus_{j \in J} M_j$  and  $N \cong \bigoplus_{j \in J} N_j$ 
  - indexed by the same set  $J$ ,
  - where each summand is either indecomposable or zero, and
  - such that  $M_j$  and  $N_j$  are  $\delta$ -interleaved for every  $j \in J$ .
- The *bottleneck distance* between  $M$  and  $N$  is

$$d_B(M, N) = \inf\{\delta \geq 0 \mid \text{there exists a } \delta\text{-matching between } M \text{ and } N\}.$$

- $M$  is *structurally stable* if passing from interleaving to bottleneck distance is continuous at  $M$ .
  - Nearby persistence modules (in the interleaving distance) have a similar structure.

# Bottleneck distance and structural stability

We can reinterpret stability of barcodes and density of indecomposables:

## Corollary

*1-parameter and multi-parameter persistence modules differ in their different structural stability:*

# Bottleneck distance and structural stability

We can reinterpret stability of barcodes and density of indecomposables:

## Corollary

*1-parameter and multi-parameter persistence modules differ in their different structural stability:*

- *Every 1-parameter pfd persistence module is structurally stable.*

# Bottleneck distance and structural stability

We can reinterpret stability of barcodes and density of indecomposables:

## Corollary

*1-parameter and multi-parameter persistence modules differ in their different structural stability:*

- *Every 1-parameter pfd persistence module is structurally stable.*
- *A finitely presentable  $n$ -parameter persistence module (with  $n > 1$ ) is structurally stable if and only if it is indecomposable.*

# Origins

# When was persistent homology discovered first?

# When was persistent homology discovered first?

ANNALS OF MATHEMATICS  
Vol. 41, No. 2, April, 1940

## RANK AND SPAN IN FUNCTIONAL TOPOLOGY

BY MARSTON MORSE

(Received August 9, 1939)

### 1. Introduction.

The analysis of functions  $F$  on metric spaces  $M$  of the type which appear in variational theories is made difficult by the fact that the critical limits, such as absolute minima, relative minima, minimax values etc., are in general infinite in number. These limits are associated with relative  $k$ -cycles of various dimensions and are classified as 0-limits, 1-limits etc. The number of  $k$ -limits suitably counted is called the  $k^{\text{th}}$  type number  $m_k$  of  $F$ . The theory seeks to establish relations between the numbers  $m_k$  and the connectivities  $p_k$  of  $M$ . The numbers  $p_k$  are finite in the most important applications. It is otherwise with the numbers  $m_k$ .

The theory has been able to proceed provided one of the following hypotheses is satisfied. The critical limits cluster at most at  $p_k$ ; the critical points are

# When was persistent homology discovered first?

Web Images More... Sign in

Google

Scholar 9 results (0.02 sec) My Citations ▾

All citations

Articles

Rank and span in functional topology

Search within citing articles

Case law

Exact homomorphism sequences in homology theory

[ed.ac.uk \[PDF\]](#)

My library

JL Kelley, E Pitcher - *Annals of Mathematics*, 1947 - JSTOR

Any time

The developments of this paper stem from the attempts of one of the authors to deduce relations between homology groups of a complex and homology groups of a complex which is its image under a simplicial map. Certain relations were deduced (see [EP 1] and [EP 2] ...)

Since 2016

Cited by 46 Related articles All 3 versions Cite Save More

Since 2015

Since 2012

Marston Morse and his mathematical works

[ams.org \[PDF\]](#)

R Bott - *Bulletin of the American Mathematical Society*, 1980 - ams.org

American Mathematical Society. Thus Morse grew to maturity just at the time when the subject of Analysis Situs was being shaped by such masters<sup>2</sup> as Poincaré, Veblen, LEJ Brouwer, GD Birkhoff, Lefschetz and Alexander, and it was Morse's genius and destiny to ...

Sort by relevance

Cited by 24 Related articles All 4 versions Cite Save More

Sort by date

Unstable minimal surfaces of higher topological structure

include citations

M Morse, CB Tompkins - *Duke Math. J.* 1941 - projecteuclid.org

1. Introduction. We are concerned with extending the calculus of variations in the large to multiple integrals. The problem of the existence of minimal surfaces of unstable type contains many of the typical difficulties, especially those of a topological nature. Having studied this ...

Create alert

Cited by 19 Related articles All 2 versions Cite Save

[PDF] Persistence in discrete Morse theory

[psu.edu \[PDF\]](#)

# When was persistent homology discovered first?

BULLETIN (New Series) OF THE  
AMERICAN MATHEMATICAL SOCIETY  
Volume 3, Number 3, November 1980

## MARSTON MORSE AND HIS MATHEMATICAL WORKS

BY RAOUL BOTT<sup>1</sup>

**1. Introduction.** Marston Morse was born in 1892, so that he was 33 years old when in 1925 his paper *Relations between the critical points of a real-valued function of n independent variables* appeared in the Transactions of the American Mathematical Society. Thus Morse grew to maturity just at the time when the subject of Analysis Situs was being shaped by such masters<sup>2</sup> as Poincaré, Veblen, L. E. J. Brouwer, G. D. Birkhoff, Lefschetz and Alexander, and it was Morse's genius and destiny to discover one of the most beautiful and far-reaching relations between this fledgling and Analysis; a relation which is now known as *Morse Theory*.


In retrospect all great ideas take on a certain simplicity and inevitability, partly because they shape the whole subsequent development of the subject. And so to us, today, Morse Theory seems natural and inevitable. However one only has to glance at these early papers to see what a tour de force it was

## When was persistent homology discovered first?

*inequalities pertain between the dimensions of the  $A_i$  and those of  $H(A_i)$ .* Thus the Morse inequalities already reflect a certain part of the “Spectral Sequence magic”, and a modern and tremendously general account of Morse’s work on rank and span in the framework of Leray’s theory was developed by Deheuvels [D] in the 50’s.

Unfortunately both Morse’s and Deheuvel’s papers are not easy reading. On the other hand there is no question in my mind that the papers [36] and [44] constitute another tour de force by Morse. Let me therefore illustrate rather than explain some of the ideas of the rank and span theory in a very simple and tame example.


In the figure which follows I have drawn a homeomorph of  $M = S^1$  in the plane, and I will be studying the height function  $F = y$  on  $M$ .



# Motivation and application: minimal surfaces

## Problem (Plateau's problem)

*Find an immersed disk of least area spanned by a given closed Jordan curve.*




(from Dierkes et al.: *Minimal Surfaces*, 2010)

# Motivation and application: minimal surfaces

## Problem (Plateau's problem)

*Find an immersed disk of least area spanned by a given closed Jordan curve.*



(from Dierkes et al.: *Minimal Surfaces*, 2010)


Solution by Douglas (1930; first Fields Medal 1936):

- identifies minimal surfaces with critical points of the *Douglas functional*

# Existence of unstable minimal surfaces

Theorem (Morse, Tompkins 1939; Shiffman 1939)

*Assume that a given curve bounds two separate stable minimal surfaces.*




# Existence of unstable minimal surfaces

Theorem (Morse, Tompkins 1939; Shiffman 1939)

Assume that a given curve bounds two separate stable minimal surfaces.

Then that curve also bounds an unstable minimal surface (a critical point that is not a local minimum).



# Whatever happened to functional topology?

# Whatever happened to functional topology?

PLATEAU'S PROBLEM  
AND THE  
CALCULUS OF VARIATIONS

BY

MICHAEL STRUWE

# Whatever happened to functional topology?

82

## A. The classical Plateau Problem for disc - type minimal surfaces.

The technical complexity and the use of a sophisticated topological machinery (which is not shadowed in our presentation) moreover tend to make Morse-Tompkins' original paper unreadable and inaccessible for the non-specialist, cf. Hildebrandt [4, p. 324].

Confronting Morse-Tompkins' and Shiffman's approach with that given in Chapter 4 we see how much can be gained in simplicity and strength by merely replacing the  $C^0$ -topology by the  $H^{1/2,2}$ -topology and verifying the Palais - Smale - type condition stated in Lemma 2.10.

However, in 1964/65 when Palais and Smale introduced this condition in the calculus of variations it was not clear that it could be meaningful for analyzing the geometry of surfaces, cf. Hildebrandt [4, p. 323 f.].

## Q-tame persistence modules

### Definition (Chazal et al. 2009)

A persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  is *q-tame* if all structure maps  $M_s \rightarrow M_t$  ( $s < t$ ) have finite rank.

# Q-tame persistence modules

## Definition (Chazal et al. 2009)

A persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  is *q-tame* if all structure maps  $M_s \rightarrow M_t$  ( $s < t$ ) have finite rank.

- A q-tame persistence module still admits a barcode (disregarding the endpoints).

# Q-tame persistence modules

## Definition (Chazal et al. 2009)

A persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  is *q-tame* if all structure maps  $M_s \rightarrow M_t$  ( $s < t$ ) have finite rank.

- A q-tame persistence module still admits a barcode (disregarding the endpoints).

Morse's main technical result, in modern language:

- The minimal surface functional has q-tame persistent homology.

## Structure of q-tame persistence modules

### Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*The radical of a q-tame persistence module  $M$ ,  $(\text{rad } M)_t = \sum_{s < t} \text{im } M_{s,t}$ ,*

- *admits a barcode decomposition, and*

## Structure of q-tame persistence modules

### Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*The radical of a q-tame persistence module  $M$ ,  $(\text{rad } M)_t = \sum_{s < t} \text{im } M_{s,t}$ ,*

- *admits a barcode decomposition, and*
- *the canonical inclusion  $\text{rad } M \hookrightarrow M$  has an ephemeral cokernel:*

## Structure of q-tame persistence modules

### Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*The radical of a q-tame persistence module  $M$ ,  $(\text{rad } M)_t = \sum_{s < t} \text{im } M_{s,t}$ ,*

- *admits a barcode decomposition, and*
- *the canonical inclusion  $\text{rad } M \hookrightarrow M$  has an ephemeral cokernel:*
  - *In the kernel, all maps between different indices  $s < t$  are zero.*

# Structure of q-tame persistence modules

## Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*The radical of a q-tame persistence module  $M$ ,  $(\text{rad } M)_t = \sum_{s < t} \text{im } M_{s,t}$ ,*

- *admits a barcode decomposition, and*
- *the canonical inclusion  $\text{rad } M \hookrightarrow M$  has an ephemeral cokernel:*
  - *In the kernel, all maps between different indices  $s < t$  are zero.*

## Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*Every q-tame persistence module has a unique persistence diagram that completely describes its isomorphism type in the observable category.*

# Structure of q-tame persistence modules

## Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*The radical of a q-tame persistence module  $M$ ,  $(\text{rad } M)_t = \sum_{s < t} \text{im } M_{s,t}$ ,*

- *admits a barcode decomposition, and*
- *the canonical inclusion  $\text{rad } M \hookrightarrow M$  has an ephemeral cokernel:*
  - *In the kernel, all maps between different indices  $s < t$  are zero.*

## Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*Every q-tame persistence module has a unique persistence diagram that completely describes its isomorphism type in the observable category.*

- *A persistence diagram describes the intervals in a barcode, modulo the endpoints.*

# Structure of q-tame persistence modules

## Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*The radical of a q-tame persistence module  $M$ ,  $(\text{rad } M)_t = \sum_{s < t} \text{im } M_{s,t}$ ,*

- *admits a barcode decomposition, and*
- *the canonical inclusion  $\text{rad } M \hookrightarrow M$  has an ephemeral cokernel:*
  - *In the kernel, all maps between different indices  $s < t$  are zero.*

## Theorem (Chazal, Crawley-Boevey, de Silva 2016)

*Every q-tame persistence module has a unique persistence diagram that completely describes its isomorphism type in the observable category.*

- *A persistence diagram describes the intervals in a barcode, modulo the endpoints.*
- *The observable category is the category of persistence modules, modulo ephemeral modules.*

## Generalized Morse inequalities

Assume that the sublevel sets of a bounded function  $f : X \rightarrow \mathbb{R}$  have q-tame persistent homology.

## Generalized Morse inequalities

Assume that the sublevel sets of a bounded function  $f : X \rightarrow \mathbb{R}$  have q-tame persistent homology.

- The persistence diagram of  $f$  has finitely many bars of length  $> \epsilon$ , for every  $\epsilon > 0$ .

## Generalized Morse inequalities

Assume that the sublevel sets of a bounded function  $f : X \rightarrow \mathbb{R}$  have q-tame persistent homology.

- The persistence diagram of  $f$  has finitely many bars of length  $> \epsilon$ , for every  $\epsilon > 0$ .
- This gives generalized Morse inequalities

$$\sum_{i=0}^d (-1)^{d-i} (m_i^\epsilon - \beta_i) \geq 0.$$

- $m_i^\epsilon$  counts endpoints in intervals with length  $> \epsilon$

## Generalized Morse inequalities

Assume that the sublevel sets of a bounded function  $f : X \rightarrow \mathbb{R}$  have q-tame persistent homology.

- The persistence diagram of  $f$  has finitely many bars of length  $> \epsilon$ , for every  $\epsilon > 0$ .
- This gives generalized Morse inequalities

$$\sum_{i=0}^d (-1)^{d-i} (m_i^\epsilon - \beta_i) \geq 0.$$

- $m_i^\epsilon$  counts endpoints in intervals with length  $> \epsilon$
- Morse and Tompkins used this idea to show the existence of an unstable minimal surface.

## Vietoris vs singular homology

## Vietoris vs singular homology

- A compact sublevel set filtration  $(f_{\leq t})_{t \in \mathbb{R}}$  is continuous from above:

$$f_{\leq t} = \bigcap_{u > t} f_{\leq u} = \lim_{u > t} f_{\leq u}$$

## Vietoris vs singular homology

- A compact sublevel set filtration  $(f_{\leq t})_{t \in \mathbb{R}}$  is continuous from above:

$$f_{\leq t} = \bigcap_{u > t} f_{\leq u} = \lim_{u > t} f_{\leq u}$$

- Vietoris (equivalently, Čech) homology (with field coefficients) preserves limits:

$$\lim \circ \check{H}_* = \check{H}_* \circ \lim$$

## Vietoris vs singular homology

- A compact sublevel set filtration  $(f_{\leq t})_{t \in \mathbb{R}}$  is continuous from above:

$$f_{\leq t} = \bigcap_{u > t} f_{\leq u} = \lim_{u > t} f_{\leq u}$$

- Vietoris (equivalently, Čech) homology (with field coefficients) preserves limits:

$$\lim \circ \check{H}_* = \check{H}_* \circ \lim$$

- Therefore, persistent Vietoris/Čech homology is also continuous from above:

$$\check{H}_*(f_{\leq t}) = \lim_{u > t} \check{H}_*(f_{\leq u})$$

## Weakly LC filtrations

### Definition (Morse 1937; paraphrased)

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is said to be *weakly locally connected* if for

# Weakly LC filtrations

## Definition (Morse 1937; paraphrased)

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is said to be *weakly locally connected* if for

- any point  $x \in X$ ,

# Weakly LC filtrations

## Definition (Morse 1937; paraphrased)

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is said to be *weakly locally connected* if for

- any point  $x \in X$ ,
- any value  $t > f(x)$ , and

# Weakly LC filtrations

## Definition (Morse 1937; paraphrased)

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is said to be *weakly locally connected* if for

- any point  $x \in X$ ,
- any value  $t > f(x)$ , and
- any neighborhood  $V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, t]$

there is

# Weakly LC filtrations

## Definition (Morse 1937; paraphrased)

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is said to be *weakly locally connected* if for

- any point  $x \in X$ ,
- any value  $t > f(x)$ , and
- any neighborhood  $V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, t]$

there is

- a value  $s$  with  $f(x) < s < t$  and

# Weakly LC filtrations

## Definition (Morse 1937; paraphrased)

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is said to be *weakly locally connected* if for

- any point  $x \in X$ ,
- any value  $t > f(x)$ , and
- any neighborhood  $V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, t]$

there is

- a value  $s$  with  $f(x) < s < t$  and
- a neighborhood  $U \subseteq V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, s]$

such that the inclusion  $U \rightarrow V$  is homotopic to a constant map.

## Q-tameness from local connectivity


### Theorem (Morse, 1937)

*If the sublevel set filtration of  $f: X \rightarrow \mathbb{R}$  is compact and weakly locally connected, then it has  $q$ -tame persistent Vietoris homology.*

# Q-tameness from local connectivity

Theorem (Morse, 1937; incorrect)

*If the sublevel set filtration of  $f: X \rightarrow \mathbb{R}$  is compact and weakly locally connected, then it has  $q$ -tame persistent Vietoris homology.*



# Homologically locally small filtrations

## Definition

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is *homologically locally connected (HLC)* if

- for any point  $x \in X$ ,
- any values  $f(x) < s < t$ , and
- any neighborhood  $V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, t]$ ,

there is

- a neighborhood  $U \subseteq V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, s]$

such that the inclusion  $U \hookrightarrow V$  induces a zero map on homology.

# Homologically locally small filtrations

## Definition

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is *homologically locally connected (HLC)* if

- for any point  $x \in X$ ,
- any values  $f(x) < s < t$ , and
- any neighborhood  $V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, t]$ ,

there is

- a neighborhood  $U \subseteq V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, s]$

such that the inclusion  $U \hookrightarrow V$  induces a zero map on homology.

## Theorem (B, Medina-Mardones, Schmahl 2021)

If the sublevel set filtration of  $f: X \rightarrow \mathbb{R}$  is compact and homologically locally connected, then their persistent homology is  $q$ -tame.

- $f$  is not required to be continuous

# Homologically locally small filtrations

## Definition

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is *homologically locally connected (HLC)* if

- for any point  $x \in X$ ,
- any values  $f(x) < s < t$ , and
- any neighborhood  $V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, t]$ ,

there is

- a neighborhood  $U \subseteq V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, s]$

such that the inclusion  $U \hookrightarrow V$  induces a zero map on homology.

## Theorem (B, Medina-Mardones, Schmahl 2021)

If the sublevel set filtration of  $f: X \rightarrow \mathbb{R}$  is compact and homologically locally connected, then their persistent homology is  $q$ -tame.

- $f$  is not required to be continuous
- Conditions are satisfied by the Douglas functional

# Homologically locally small filtrations

## Definition

The sublevel set filtration of a function  $f: X \rightarrow \mathbb{R}$  is *homologically locally connected (HLC)* if

- for any point  $x \in X$ ,
- any values  $f(x) < s < t$ , and
- any neighborhood  $V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, t]$ ,

there is

- a neighborhood  $U \subseteq V$  of  $x$  in the sublevel set  $f^{-1}(-\infty, s]$

such that the inclusion  $U \hookrightarrow V$  induces a zero map on homology.

## Theorem (B, Medina-Mardones, Schmahl 2021)

If the sublevel set filtration of  $f: X \rightarrow \mathbb{R}$  is compact and homologically locally connected, then their persistent homology is  $q$ -tame.

- $f$  is not required to be continuous
- Conditions are satisfied by the Douglas functional
- Fixes the gap in Morse/Tompkins' proof

# Simplification

# Topological simplification of functions

Consider the following problem:

## Problem (Topological simplification)


*Given a function  $f: X \rightarrow \mathbb{R}$  and a real number  $\delta \geq 0$ , find a function  $f_\delta$*

- *with the minimal number of critical points*
- *subject to  $\|f_\delta - f\|_\infty \leq \delta$ .*

# Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points




(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

# Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points



(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

Persistent homology of a Morse function:

- critical points correspond to endpoints of barcode intervals

## Canceling persistence pairs

By stability of persistence barcodes:

### Proposition

*The critical points of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

# Canceling persistence pairs

By stability of persistence barcodes:


## Proposition

*The critical points of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*


## Theorem (B, Lange, Wardetzky, 2011)


*This lower bound can be achieved for a function on a surface*


- by canceling all critical points with persistence  $\leq 2\delta$ .














Thanks for your attention!