

Discrete Morse theory and persistence of geometric complexes

Elements of computational geometry & topology

Ulrich Bauer

Technical University of Munich (TUM)

February 26, 2024

Dagstuhl-Seminar 24092 *Applied and Combinatorial Topology*

Funded by



Deutsche
Forschungsgemeinschaft
German Research Foundation



SFB
TRR
109


Discretization
in Geometry
and Dynamics


Technical
University
of Munich





MCMCL
Munich Center for Machine Learning













Geometric complexes


Čech complexes




Čech complexes




Čech complexes




Čech complexes



Čech complexes



Čech complexes



Nerves

Definition (Alexandrov 1928)

Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a space X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}$$

recording the nonempty intersections of cover elements.

- Any subset of \mathcal{U} with nonempty intersection corresponds to a simplex in $\text{Nrv}(\mathcal{U})$

Nerves

Definition (Alexandrov 1928)

Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a space X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}$$

recording the nonempty intersections of cover elements.

- Any subset of \mathcal{U} with nonempty intersection corresponds to a simplex in $\text{Nrv}(\mathcal{U})$

Theorem (Borsuk 1948, and many more)

Let \mathcal{U} be a good cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Nerves

Definition (Alexandrov 1928)

Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a space X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}$$

recording the nonempty intersections of cover elements.

- Any subset of \mathcal{U} with nonempty intersection corresponds to a simplex in $\text{Nrv}(\mathcal{U})$

Theorem (Borsuk 1948, and many more)

Let \mathcal{U} be a good cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Here *good* can mean different things:

- open, numerable cover, contractible intersections
- closed, convex finite cover

Nerves

Definition (Alexandrov 1928)

Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a space X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid 0 < |J| < \infty \text{ and } \bigcap_{j \in J} U_j \neq \emptyset\}$$

recording the nonempty intersections of cover elements.

- Any subset of \mathcal{U} with nonempty intersection corresponds to a simplex in $\text{Nrv}(\mathcal{U})$

Theorem (Borsuk 1948, and many more)

Let \mathcal{U} be a good cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Here *good* can mean different things:

- open, numerable cover, contractible intersections
- closed, convex finite cover




U. Bauer, M. Kerber, F. Roll, and A. Rolle

A Unified View on the Functorial Nerve Theorem and its Variations

Nerve theorems, almost true

Nerve theorems, almost true



3. Nerve of a Covering


Let Y be a topological space, $U = (U_\alpha)$ be its covering (either by open or by closed sets), α running over some set A . Let

$$X_n = \{(\alpha_0, \dots, \alpha_n) \mid U_{\alpha_0} \cap \dots \cap U_{\alpha_n} \neq \emptyset\},$$

$$X(f)(\alpha_0, \dots, \alpha_n) = (\alpha_{f(0)}, \dots, \alpha_{f(m)}) \quad \text{for } f : [m] \rightarrow [n].$$

This simplicial set reflects the combinatorial structure of a covering. One can show that if the covering U is locally finite and all nonempty finite intersections $U_{\alpha_0} \cap \dots \cap U_{\alpha_n}$ are contractible, then the geometric realization $|X|$ of X is homotopically equivalent to Y , so that the topology can be efficiently encoded into combinatorical data.


Nerve theorems, almost true



Nerve theorems, almost true


Algebraic Topology

Allen Hatcher




Nerve theorems, almost true

Since X is paracompact there is a partition of unity subordinate to the cover \mathcal{U} . This is a family of maps $\varphi_\alpha : X \rightarrow [0, 1]$ satisfying three conditions: The support of each φ_α is contained in some $X_{i(\alpha)}$, only finitely many φ_α 's are nonzero near each point of X , and $\sum_\alpha \varphi_\alpha = 1$. Define a section $s : X \rightarrow \Delta X_{\mathcal{U}}$ of p by setting $s(x) = \sum_\alpha \varphi_\alpha(x) x_{i(\alpha)}$. The figure shows the case $X = S^1$ with a cover by two arcs, the heavy line indicating the image of s . In the general case the section s embeds X as a retract of $\Delta X_{\mathcal{U}}$, and it is a deformation retract since points in fibers $p^{-1}(x)$ can move linearly along line segments to $s(x)$. \square




Corollary 4G.3. *If \mathcal{U} is an open cover of a paracompact space X such that every nonempty intersection of finitely many sets in \mathcal{U} is contractible, then X is homotopy equivalent to the nerve $N\mathcal{U}$.*


Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes




Delaunay complexes



Discrete Morse theory




Discrete Morse theory



Discrete Morse theory

Theorem (Forman 1998)


A simplicial complex with a discrete Morse function f is homotopy equivalent to a space (a CW complex) built from the critical simplices of f .



Discrete Morse theory

Theorem (Forman 1998)

A simplicial complex with a discrete Morse function f is homotopy equivalent to a space (a CW complex) built from the critical simplices of f .




Discrete Morse functions – and their gradients – encode *collapses* of sublevel sets:




Discrete Morse theory

Theorem (Forman 1998)

A simplicial complex with a discrete Morse function f is homotopy equivalent to a space (a CW complex) built from the critical simplices of f .




Discrete Morse functions – and their gradients – encode *collapses* of sublevel sets:



Generalizing discrete Morse theory

Generalized gradients are a partition into intervals in the face poset (instead of just facet pairs):




Morse theory for Čech and Delaunay complexes

Theorem (B, Edelsbrunner 2017)

Čech, Delaunay, and Wrap complexes (at any scale r) of a point set $X \subset \mathbb{R}^d$ in general position are related by collapses encoded by a single discrete gradient field:

$$\text{Cech}_r X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$



Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

The Čech function and the Delaunay function of a point set $X \subset \mathbb{R}^d$ in general position are generalized discrete Morse functions.

Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

The Čech function and the Delaunay function of a point set $X \subset \mathbb{R}^d$ in general position are generalized discrete Morse functions.

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$

Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

The Čech function and the Delaunay function of a point set $X \subset \mathbb{R}^d$ in general position are generalized discrete Morse functions.

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$
- Q is a critical simplex off f_C

Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

The Čech function and the Delaunay function of a point set $X \subset \mathbb{R}^d$ in general position are generalized discrete Morse functions.

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$
- Q is a critical simplex off f_C
- Q is a critical simplex off f_D


Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

The Čech function and the Delaunay function of a point set $X \subset \mathbb{R}^d$ in general position are generalized discrete Morse functions.

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$
- Q is a critical simplex of f_C
- Q is a critical simplex of f_D
- Q is a centered Delaunay simplex
(containing the circumcenter in the interior)




Čech intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff




Čech intervals

Lemma


Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff

On $S \subseteq Q'$



Q



On S


Čech intervals

Lemma


Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$\text{On } S \subseteq Q' \subseteq \text{Encl } S.$$



Q



$\text{On } S$



$\text{Encl } S$


Čech intervals

Lemma


Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$Q' \in [\text{On } S, \text{Encl } S].$$



Q



$\text{On } S$



$\text{Encl } S$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex On $S \subseteq X$.

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.

Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.


Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex X . On $S \subseteq X$.



Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$






Delaunay intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff

$$Q' \in [\text{Front } S, \text{On } S].$$



Čech and Delaunay intervals from Karush–Kuhn–Tucker conditions

Proposition

A sphere S is the smallest sphere enclosing Q and excluding E iff

- S is the smallest circumsphere of some simplex $P \subseteq X$,
- $Q \in [\text{Front } S, \text{Encl } S]$, and
- $E \in [\text{Back } S, \text{Excl } S]$.

$\check{\text{C}}$ ech and Delaunay intervals from Karush–Kuhn–Tucker conditions

Proposition

A sphere S is the smallest sphere enclosing Q and excluding E iff

- S is the smallest circumsphere of some simplex $P \subseteq X$,
- $Q \in [\text{Front } S, \text{Encl } S]$, and
- $E \in [\text{Back } S, \text{Excl } S]$.

Corollary

The $\check{\text{C}}$ ech gradient is given by the intervals of the form $[\text{On } S, \text{Encl } S]$.

$\check{\text{C}}$ ech and Delaunay intervals from Karush–Kuhn–Tucker conditions

Proposition

A sphere S is the smallest sphere enclosing Q and excluding E iff

- S is the smallest circumsphere of some simplex $P \subseteq X$,
- $Q \in [\text{Front } S, \text{Encl } S]$, and
- $E \in [\text{Back } S, \text{Excl } S]$.


Corollary

The $\check{\text{C}}$ ech gradient is given by the intervals of the form $[\text{On } S, \text{Encl } S]$.

The Delaunay intervals is given by the intervals of the form $[\text{Front } S, \text{On } S]$.


Wrap complexes

Foundation of the surface reconstruction software *Wrap* (Edelsbrunner 1995, Geomagic)




Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.



Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.



Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)


$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r X$:

- the smallest subcomplex of $\text{Del}_r X$ that
 - contains all critical simplices
 - is compatible with V (a union of intervals of V).

Delaunay and Wrap complexes



Delaunay and Wrap complexes



Persistence of Delaunay filtrations

Algorithm (matrix reduction)

Require: $D: m \times n$ matrix

Ensure: $R = D \cdot V$ is reduced, V is full rank upper triangular

function Reduce(D)

$R = D, V = I(n)$

while there exist $i < j$ such that pivot $R_i =$ pivot R_j **do**

 add column R_i to column R_j

 add column V_i to column V_j

return R, V

Algorithm (exhaustive matrix reduction)

Require: $D: m \times n$ matrix

Ensure: $R = D \cdot V$ is reduced, V is full rank upper triangular

function Reduce(D)

$R = D, V = I(n)$

while there exist $i < j$ such that R_j has a nonzero entry in row pivot R_i **do**

 add column R_i to column R_j

 add column V_i to column V_j

return R, V

Algorithm (matrix reduction)

Require: $D: m \times n$ matrix

Ensure: $R = D \cdot V$ is reduced, V is full rank upper triangular

function Reduce(D)

$R = D, V = I(n)$

while there exist $i < j$ such that pivot $R_i =$ pivot R_j **do**

 add column R_i to column R_j

 add column V_i to column V_j

return R, V


Proposition

The resulting columns R_j are minimal (in a lexicographic order) within their homology class (in K_{j-1}).


Standard reduction and exhaustive reduction




Standard reduction and exhaustive reduction




Wrap complexes and lexicographically minimal cycles



Wrap complexes and lexicographically minimal cycles



Wrap complexes and lexicographically minimal cycles




Wrap complexes and lexicographically minimal cycles




Wrap complexes and lexicographically minimal cycles




Wrap complexes and lexicographically minimal cycles






Wrap complexes and lexicographically minimal cycles



Wrap complexes and lexicographically minimal cycles



Point cloud reconstruction with minimal cycles




Wrap complexes support minimal cycles

Theorem (B, Roll 2022)

Let $X \subset \mathbb{R}$ be a finite subset in general position and let $r \in \mathbb{R}$.

- Exhaustive matrix reduction computes the minimal cycles homologous to a simplex boundary.
- Any lexicographically minimal cycle of $\text{Del}_r(X)$ is supported on $\text{Wrap}_r(X)$.




Vietoris–Rips persistence

Vietoris–Rips complexes

For a metric space X , the *Vietoris–Rips complex* at $t > 0$ is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



Vietoris–Rips complexes

For a metric space X , the *Vietoris–Rips complex* at $t > 0$ is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



Vietoris–Rips complexes

For a metric space X , the *Vietoris–Rips complex* at $t > 0$ is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



Vietoris–Rips complexes

For a metric space X , the *Vietoris–Rips complex* at $t > 0$ is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



Vietoris–Rips complexes



For a metric space X , the *Vietoris–Rips complex* at $t > 0$ is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



Ripser: software for computing Vietoris–Rips persistence barcodes



Open source software (ripser.org)



Ripser users across the globe

Ripser: software for computing Vietoris–Rips persistence barcodes

Open source software (ripser.org)



Ripser users across the globe

Computational improvements based on

- *implicit matrix representations*
- *apparent pairs*, connecting persistence to discrete Morse theory

Apparent pairs

Ripser uses the following pairing of simplices (breaking ties in the filtration lexicographically):

Definition (B 2016, 2021)

In a simplexwise filtration ($K_i = \{\sigma_1, \dots, \sigma_i\}$)_i, two simplices (σ_i, σ_j) form an *apparent pair* if

- σ_i is the latest proper face of σ_j , and
- σ_j is the earliest proper coface of σ_i .

Apparent pairs

Ripser uses the following pairing of simplices (breaking ties in the filtration lexicographically):

Definition (B 2016, 2021)

In a simplexwise filtration ($K_i = \{\sigma_1, \dots, \sigma_i\}$)_i, two simplices (σ_i, σ_j) form an *apparent pair* if

- σ_i is the latest proper face of σ_j , and
- σ_j is the earliest proper coface of σ_i .

Special cases and equivalent definitions have been (re)discovered independently multiple times

- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

Apparent pairs

Ripser uses the following pairing of simplices (breaking ties in the filtration lexicographically):

Definition (B 2016, 2021)

In a simplexwise filtration ($K_i = \{\sigma_1, \dots, \sigma_i\}_i$), two simplices (σ_i, σ_j) form an *apparent pair* if

- σ_i is the latest proper face of σ_j , and
- σ_j is the earliest proper coface of σ_i .

Special cases and equivalent definitions have been (re)discovered independently multiple times

- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

Proposition (B 2021)

The apparent pairs are both

- gradient pairs (in the sense of discrete Morse theory) and
- persistence pairs (creating/destroying a feature in homology).

Apparent pairs

Ripser uses the following pairing of simplices (breaking ties in the filtration lexicographically):

Definition (B 2016, 2021)

In a simplexwise filtration ($K_i = \{\sigma_1, \dots, \sigma_i\}\}_i$, two simplices (σ_i, σ_j) form an *apparent pair* if

- σ_i is the latest proper face of σ_j , and
- σ_j is the earliest proper coface of σ_i .

Special cases and equivalent definitions have been (re)discovered independently multiple times

- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

Proposition (B 2021)


The apparent pairs are both

- gradient pairs (in the sense of discrete Morse theory) and
- persistence pairs (creating/destroying a feature in homology).

Proposition (B, Roll 2022)


Any generalized discrete Morse function is refined by apparent pairs.

Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadañ, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Observation: Ripser runs unusually fast on genetic distance data


- SARS-CoV2 RNA sequences (spike protein)
- 25556 data points (2.8×10^{12} simplices in 2-skeleton)
- 120 s computation time (with data points ordered appropriately)

Gromov-hyperbolicity

Definition (Gromov 1988)

A metric space X is δ -hyperbolic (for $\delta \geq 0$) if for all $w, x, y, z \in X$ we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$




Gromov-hyperbolicity

Definition (Gromov 1988)

A metric space X is δ -hyperbolic (for $\delta \geq 0$) if for all $w, x, y, z \in X$ we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$



- The hyperbolic plane is $(\ln 2)$ -hyperbolic.




Gromov-hyperbolicity

Definition (Gromov 1988)

A metric space X is δ -hyperbolic (for $\delta \geq 0$) if for all $w, x, y, z \in X$ we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$



- The hyperbolic plane is $(\ln 2)$ -hyperbolic.



- The 0-hyperbolic spaces are precisely the metric trees and their subspaces.



Rips Contractibility

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.

Rips Contractibility

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.

What about

- non-geodesic spaces? In particular, finite metric spaces?

Rips Contractibility

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.

What about

- non-geodesic spaces? In particular, finite metric spaces?
- collapsibility?

Rips Contractibility

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.

What about

- non-geodesic spaces? In particular, finite metric spaces?
- collapsibility?
- the filtration?

Rips Contractibility

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.

What about

- non-geodesic spaces? In particular, finite metric spaces?
- collapsibility?
- the filtration?
- the connection to computation of persistent homology?

Rips Contractibility

Theorem (Rips; Gromov 1988)

Let X be a δ -hyperbolic geodesic metric space. Then $\text{Rips}_t(X)$ is contractible for all $t \geq 4\delta$.

What about

- non-geodesic spaces? In particular, finite metric spaces?
- collapsibility?
- the filtration?
- the connection to computation of persistent homology?


Theorem (B, Roll 2022)

Let X be a finite δ -hyperbolic space. Then there is a single discrete gradient encoding the collapses


$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow \{\ast\}$$

for all $u > t \geq 4\delta + 2\nu$, where ν is the geodesic defect of X .


Geodesic defect




Geodesic defect



Geodesic defect



Geodesic defect



Definition (Bonk, Schramm 2000)

A metric space X is ν -geodesic if for all points $x, y \in X$ and all $r, s \geq 0$ with $r + s = d(x, y)$ we have

$$B_{r+\nu}(x) \cap B_{s+\nu}(y) \neq \emptyset.$$

The infimum of all such ν is the *geodesic defect* of X .

The diameter function of generic trees

Proposition (B, Roll 2022)

Consider a finite weighted tree (V, E) with a generic path length metric (distinct pairwise distances). Then the diameter function $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$ is a generalized discrete Morse function.

The diameter function of generic trees

Proposition (B, Roll 2022)

Consider a finite weighted tree (V, E) with a generic path length metric (distinct pairwise distances). Then the diameter function $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$ is a generalized discrete Morse function.

- The apparent pairs refine this gradient.

The diameter function of generic trees

Proposition (B, Roll 2022)

Consider a finite weighted tree (V, E) with a generic path length metric (distinct pairwise distances). Then the diameter function $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$ is a generalized discrete Morse function.

- The apparent pairs refine this gradient.

Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space X induces the collapses

- $\text{Rips}_t(X) \searrow T_t$ for all $t \in \mathbb{R}$,

The diameter function of generic trees

Proposition (B, Roll 2022)

Consider a finite weighted tree (V, E) with a generic path length metric (distinct pairwise distances). Then the diameter function $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$ is a generalized discrete Morse function.

- The apparent pairs refine this gradient.

Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space X induces the collapses

- $\text{Rips}_t(X) \searrow T_t$ for all $t \in \mathbb{R}$,
- $\text{Rips}_t(X) \searrow T \searrow \{\ast\}$ for $t \geq \max d(E)$, and

The diameter function of generic trees

Proposition (B, Roll 2022)

Consider a finite weighted tree (V, E) with a generic path length metric (distinct pairwise distances). Then the diameter function $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$ is a generalized discrete Morse function.

- The apparent pairs refine this gradient.

Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space X induces the collapses

- $\text{Rips}_t(X) \searrow T_t$ for all $t \in \mathbb{R}$,
- $\text{Rips}_t(X) \searrow T \searrow \{\ast\}$ for $t \geq \max d(E)$, and
- $\text{Rips}_u(X) \searrow \text{Rips}_t(X)$ whenever no pairwise distance lies in the interval $(t, u]$.

The diameter function of generic trees

Proposition (B, Roll 2022)

Consider a finite weighted tree (V, E) with a generic path length metric (distinct pairwise distances). Then the diameter function $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$ is a generalized discrete Morse function.

- The apparent pairs refine this gradient.

Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space X induces the collapses

- $\text{Rips}_t(X) \searrow T_t$ for all $t \in \mathbb{R}$,
- $\text{Rips}_t(X) \searrow T \searrow \{\ast\}$ for $t \geq \max d(E)$, and
- $\text{Rips}_u(X) \searrow \text{Rips}_t(X)$ whenever no pairwise distance lies in the interval $(t, u]$.

In particular, the persistent homology is trivial in degrees > 0 .

Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree $T = (V, E)$, viewed as a metric space X .

Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree $T = (V, E)$, viewed as a metric space X .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.


Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree $T = (V, E)$, viewed as a metric space X .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for this total order induces the same collapses as in the generic case.




Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree $T = (V, E)$, viewed as a metric space X .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for this total order induces the same collapses as in the generic case.




Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree $T = (V, E)$, viewed as a metric space X .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for this total order induces the same collapses as in the generic case.




Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree $T = (V, E)$, viewed as a metric space X .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for this total order induces the same collapses as in the generic case.




Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like, non-generic metric)?

- Consider a weighted finite tree $T = (V, E)$, viewed as a metric space X .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

Theorem (B, Roll 2022)

The apparent pairs gradient for this total order induces the same collapses as in the generic case.



Thanks for your attention!

Thanks for your attention!



U. Bauer, F. Roll

Gromov hyperbolicity, geodesic defect, and apparent pairs in Vietoris-Rips filtrations

Symposium on Computational Geometry, 2022. doi:10.4230/LIPIcs.SoCG.2022.15



U. Bauer, F. Roll

Wrapping Cycles in Delaunay Complexes: Bridging Persistent Homology and Discrete Morse Theory

Symposium on Computational Geometry, 2024. arXiv:2212.02345



U. Bauer, M. Kerber, F. Roll, and A. Rolle

A Unified View on the Functorial Nerve Theorem and its Variations

Expositiones Mathematicae, 2023. doi:10.1016/j.exmath.2023.04.005



U. Bauer, H. Edelsbrunner

The Morse Theory of Čech and Delaunay Complexes

Transactions of the AMS, 2017. doi:10.1090/tran/6991