



# Induced matchings and the algebraic stability of persistence barcodes


*Ulrich Bauer* (IST Austria), Michael Lesnick (IMA)

IST Austria


Jul 24, 2014
































# What is persistent homology?





# What is persistent homology?





# What is persistent homology?




# What is persistent homology?



# What is persistent homology?




# The pipeline of topological data analysis



# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)


If two functions  $f, g : K \rightarrow \mathbb{R}$  have distance  $\|f - g\|_\infty \leq \delta$   
then there exists a  $\delta$ -matching of their barcodes.



# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions  $f, g : K \rightarrow \mathbb{R}$  have distance  $\|f - g\|_\infty \leq \delta$   
then there exists a  $\delta$ -matching of their barcodes.




- matching  $A \leftrightarrow B$ : bijection of subsets  $A' \subseteq A, B' \subseteq B$

# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions  $f, g : K \rightarrow \mathbb{R}$  have distance  $\|f - g\|_\infty \leq \delta$   
then there exists a  $\delta$ -matching of their barcodes.




- matching  $A \leftrightarrow B$ : bijection of subsets  $A' \subseteq A, B' \subseteq B$
- $\delta$ -matching of barcodes:

# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions  $f, g : K \rightarrow \mathbb{R}$  have distance  $\|f - g\|_\infty \leq \delta$   
then there exists a  $\delta$ -matching of their barcodes.




- matching  $A \leftrightarrow B$ : bijection of subsets  $A' \subseteq A, B' \subseteq B$
- $\delta$ -matching of barcodes:
  - matched intervals have endpoints within distance  $\leq \delta$

# Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions  $f, g : K \rightarrow \mathbb{R}$  have distance  $\|f - g\|_\infty \leq \delta$   
then there exists a  $\delta$ -matching of their barcodes.




- matching  $A \leftrightarrow B$ : bijection of subsets  $A' \subseteq A, B' \subseteq B$
- $\delta$ -matching of barcodes:
  - matched intervals have endpoints within distance  $\leq \delta$
  - unmatched intervals have length  $\leq 2\delta$


# Stability for functions in the big picture




# Stability for functions in the big picture



# Stability for functions in the big picture



# Stability for functions in the big picture



# Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$ ,
- $G_t = g^{-1}(-\infty, t]$ .

# Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$ ,
- $G_t = g^{-1}(-\infty, t]$ .

If  $\|f - g\|_\infty \leq \delta$  then  $F_t \subseteq G_{t+\delta}$  and  $G_t \subseteq F_{t+\delta}$ .


# Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$ ,
- $G_t = g^{-1}(-\infty, t]$ .

If  $\|f - g\|_\infty \leq \delta$  then  $F_t \subseteq G_{t+\delta}$  and  $G_t \subseteq F_{t+\delta}$ .

So the sublevel sets are *interleaved*:




# Interleavings of sublevel sets

Let

- $F_t = f^{-1}(-\infty, t]$ ,
- $G_t = g^{-1}(-\infty, t]$ .

If  $\|f - g\|_\infty \leq \delta$  then  $F_t \subseteq G_{t+\delta}$  and  $G_t \subseteq F_{t+\delta}$ .

So the sublevel sets are *interleaved*:



Therefore their homology groups are interleaved as well.

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

## Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- a vector space  $M_t$  for each  $t \in \mathbb{R}$

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- a vector space  $M_t$  for each  $t \in \mathbb{R}$  (in this talk:  $\dim M_t < \infty$ )

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- a vector space  $M_t$  for each  $t \in \mathbb{R}$  (in this talk:  $\dim M_t < \infty$ )
- a linear map  $M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- a vector space  $M_t$  for each  $t \in \mathbb{R}$  (in this talk:  $\dim M_t < \infty$ )
- a linear map  $M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
- respecting identity:  $\text{id}_{M_t} : M_t \rightarrow M_t$   
and composition:  $(M_r \rightarrow M_s \rightarrow M_t) = (M_r \rightarrow M_t)$

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- a vector space  $M_t$  for each  $t \in \mathbb{R}$  (in this talk:  $\dim M_t < \infty$ )
- a linear map  $M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
- respecting identity:  $\text{id}_{M_t} : M_t \rightarrow M_t$   
and composition:  $(M_r \rightarrow M_s \rightarrow M_t) = (M_r \rightarrow M_t)$

A *morphism*  $f : M \rightarrow N$  is a *natural transformation*:

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- a vector space  $M_t$  for each  $t \in \mathbb{R}$  (in this talk:  $\dim M_t < \infty$ )
- a linear map  $M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
- respecting identity:  $\text{id}_{M_t} : M_t \rightarrow M_t$   
and composition:  $(M_r \rightarrow M_s \rightarrow M_t) = (M_r \rightarrow M_t)$

A *morphism*  $f : M \rightarrow N$  is a *natural transformation*:

- a linear map  $f_t : M_t \rightarrow N_t$  for each  $t \in \mathbb{R}$

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- a vector space  $M_t$  for each  $t \in \mathbb{R}$  (in this talk:  $\dim M_t < \infty$ )
- a linear map  $M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
- respecting identity:  $\text{id}_{M_t} : M_t \rightarrow M_t$   
and composition:  $(M_r \rightarrow M_s \rightarrow M_t) = (M_r \rightarrow M_t)$

A *morphism*  $f : M \rightarrow N$  is a *natural transformation*:

- a linear map  $f_t : M_t \rightarrow N_t$  for each  $t \in \mathbb{R}$
- morphism and transition maps commute:

$$\begin{array}{ccc} M_s & \longrightarrow & M_t \\ f_s \downarrow & & \downarrow f_t \\ N_s & \longrightarrow & N_t \end{array}$$

# Interval Persistence Modules

Let  $\mathcal{I}_{\mathbb{R}}$  denote the set of all intervals in  $\mathbb{R}$ .

For  $I \in \mathcal{I}_{\mathbb{R}}$ , define the *interval persistence module*  $C(I)$  by

$$C(I)_t = \begin{cases} \mathbb{K} & \text{if } t \in I, \\ 0 & \text{otherwise.} \end{cases}$$

$$C(I)_s \rightarrow C(I)_t = \begin{cases} \text{id}_{\mathbb{K}} & \text{if } s, t \in I, \\ 0 & \text{otherwise.} \end{cases}$$

# The structure of persistence modules

Theorem (Crawley-Boevey 2012)

*Let  $M$  be a pointwise finite dimensional persistence module:*

*$M_t$  is finite dimensional for all  $t$ .*

*Then  $M$  is interval-decomposable:*

*there exists a collection of intervals  $B(M)$  such that*

$$M \cong \bigoplus_{I \in B(M)} C(I).$$

Theorem (Azumaya 1950, de Silva et al. 2012)

*If  $M$  is interval-decomposable, then  $B(M)$  is unique.*

$B(M)$  is called the *barcode* of  $M$ .

- Motivates use of homology with field coefficients

# Interleavings of persistence modules

## Definition

Two persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if there are morphisms

$$f : M \rightarrow N(\delta), \quad g : N \rightarrow M(\delta)$$

such that this diagram commutes for all  $t$ :

$$\begin{array}{ccccc} M_t & \xrightarrow{\hspace{2cm}} & M_{t+2\delta} & & \\ f_t \searrow & & \nearrow g_{t+\delta} & & f_{t+2\delta} \searrow \\ & N_{t+\delta} & \xrightarrow{\hspace{2cm}} & N_{t+3\delta} & \end{array}$$

# Interleavings of persistence modules

## Definition

Two persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if there are morphisms

$$f : M \rightarrow N(\delta), \quad g : N \rightarrow M(\delta)$$

such that this diagram commutes for all  $t$ :

$$\begin{array}{ccccc} M_t & \xrightarrow{\hspace{2cm}} & M_{t+2\delta} & & \\ f_t \searrow & & \nearrow g_{t+\delta} & & f_{t+2\delta} \searrow \\ & N_{t+\delta} & \xrightarrow{\hspace{2cm}} & N_{t+3\delta} & \end{array}$$

- define  $M(\delta)$  by  $M(\delta)_t = M_{t+\delta}$

# Interleavings of persistence modules

## Definition

Two persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if there are morphisms

$$f : M \rightarrow N(\delta), \quad g : N \rightarrow M(\delta)$$

such that this diagram commutes for all  $t$ :

$$\begin{array}{ccccc} M_t & \xrightarrow{\hspace{2cm}} & M_{t+2\delta} & & \\ f_t \searrow & & \nearrow g_{t+\delta} & & f_{t+2\delta} \searrow \\ & N_{t+\delta} & \xrightarrow{\hspace{2cm}} & N_{t+3\delta} & \end{array}$$

- define  $M(\delta)$  by  $M(\delta)_t = M_{t+\delta}$   
(shift barcode to the left by  $\delta$ )



# Interleavings of persistence modules

## Definition

Two persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if there are morphisms

$$f : M \rightarrow N(\delta), \quad g : N \rightarrow M(\delta)$$

such that this diagram commutes for all  $t$ :

$$\begin{array}{ccccc} M_t & \xrightarrow{\hspace{2cm}} & M_{t+2\delta} & & \\ f_t \searrow & & \nearrow g_{t+\delta} & & f_{t+2\delta} \searrow \\ & N_{t+\delta} & \xrightarrow{\hspace{2cm}} & N_{t+3\delta} & \end{array}$$


- define  $M(\delta)$  by  $M(\delta)_t = M_{t+\delta}$   
(shift barcode to the left by  $\delta$ )



# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012)


*If two persistence modules are  $\delta$ -interleaved  
then there exists a  $\delta$ -matching of their barcodes.*



# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012)

*If two persistence modules are  $\delta$ -interleaved  
then there exists a  $\delta$ -matching of their barcodes.*



- converse statement also holds (isometry theorem)
- indirect proof, 80 page paper (Chazal et al. 2012)

## Earlier proofs of ASP

Stability for functions (Cohen-Steiner et al. 2005)

- uses interpolation between functions
- subdivides interpolation in sufficiently small pieces

## Earlier proofs of ASP

Stability for functions (Cohen-Steiner et al. 2005)

- uses interpolation between functions
- subdivides interpolation in sufficiently small pieces

Vineyards (Edelsbrunner et al. 2007)

- subdivides at changes of persistence pairings

## Earlier proofs of ASP

Stability for functions (Cohen-Steiner et al. 2005)

- uses interpolation between functions
- subdivides interpolation in sufficiently small pieces

Vineyards (Edelsbrunner et al. 2007)

- subdivides at changes of persistence pairings

Algebraic stability (Chazal et al. 2009)

- uses interpolation between persistence modules
- mimics proof for functions

# Earlier proofs of ASP

Stability for functions (Cohen-Steiner et al. 2005)

- uses interpolation between functions
- subdivides interpolation in sufficiently small pieces

Vineyards (Edelsbrunner et al. 2007)

- subdivides at changes of persistence pairings

Algebraic stability (Chazal et al. 2009)

- uses interpolation between persistence modules
- mimics proof for functions

Algebraic stability (Chazal et al. 2012)

- revision and expansion of previous proof

# Our approach

Our proof takes a different approach:

- direct proof (no interpolation, matching immediately from interleaving)

# Our approach

Our proof takes a different approach:

- direct proof (no interpolation, matching immediately from interleaving)
- shows how morphism induces a matching

# Our approach

Our proof takes a different approach:

- direct proof (no interpolation, matching immediately from interleaving)
- shows how morphism induces a matching
- stability follows from properties of a *single morphism*, not just from a pair of morphisms

# Our approach

Our proof takes a different approach:

- direct proof (no interpolation, matching immediately from interleaving)
- shows how morphism induces a matching
- stability follows from properties of a *single morphism*, not just from a pair of morphisms
- relies on *partial functoriality* of the induced matching

# The matching category

A *matching*  $\sigma : S \leftrightarrow T$  is a bijection  $S' \rightarrow T'$ , where  $S' \subseteq S$ ,  $T' \subseteq T$ .  
Formally:

$$\sigma = \{(s, t) \in S' \times T' \mid \sigma(s) = t\} \subseteq S \times T.$$

Composition of matchings  $\sigma : S \leftrightarrow T$  and  $\tau : T \leftrightarrow U$ :


$$\tau \circ \sigma = \{(s, u) \mid (s, t) \in \sigma, (t, u) \in \tau \text{ for some } t \in T\}.$$

Matchings form a category **Mch**

- objects: sets
- morphisms: matchings

# Barcodes as matching diagrams


We can regard a barcode  $\mathcal{D}$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :



# Barcodes as matching diagrams

We can regard a barcode  $\mathcal{D}$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :


- For each real number  $t$ , let  $\mathcal{D}_t$  be the subset of  $\mathcal{D}$  consisting of intervals which contain  $t$ , and



# Barcodes as matching diagrams

We can regard a barcode  $\mathcal{D}$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :

- For each real number  $t$ , let  $\mathcal{D}_t$  be the subset of  $\mathcal{D}$  consisting of intervals which contain  $t$ , and
- for each  $s \leq t$ , define the matching  $\mathcal{D}_{s,t} : \mathcal{D}_s \leftrightarrow \mathcal{D}_t$  to be the identity on  $\mathcal{D}_s \cap \mathcal{D}_t$ .



# Barcode matchings as interleavings

We can regard matchings of barcodes  $\sigma : \mathcal{C} \leftrightarrow \mathcal{D}$  as morphisms.

- Morphisms  $\mathcal{C} \rightarrow \mathcal{D}$  are natural transformations;  
the matchings  $\sigma_t : \mathcal{C}_t \leftrightarrow \mathcal{D}_t$  are restrictions of  $\sigma$  to  $\mathcal{C}_t \times \mathcal{D}_t$ :

$$\begin{array}{ccc} \mathcal{C}_s & \longleftrightarrow & \mathcal{C}_t \\ \downarrow \sigma_s & & \downarrow \sigma_t \\ \mathcal{D}_s & \longleftrightarrow & \mathcal{D}_t \end{array}$$

# Barcode matchings as interleavings

We can regard matchings of barcodes  $\sigma : \mathcal{C} \leftrightarrow \mathcal{D}$  as morphisms.

- Morphisms  $\mathcal{C} \rightarrow \mathcal{D}$  are natural transformations;  
the matchings  $\sigma_t : \mathcal{C}_t \leftrightarrow \mathcal{D}_t$  are restrictions of  $\sigma$  to  $\mathcal{C}_t \times \mathcal{D}_t$ :

$$\begin{array}{ccc} \mathcal{C}_s & \longleftrightarrow & \mathcal{C}_t \\ \downarrow \sigma_s & & \downarrow \sigma_t \\ \mathcal{D}_s & \longleftrightarrow & \mathcal{D}_t \end{array}$$

This way, we can regard a  $\delta$ -matching of barcodes  $\mathcal{C} \leftrightarrow \mathcal{D}$  as a  $\delta$ -interleaving:

$$\begin{array}{ccccc} & & \mathcal{C}_t & \longleftrightarrow & \mathcal{C}_{t+2\delta} \\ & \nwarrow & & \nearrow & \nwarrow \\ & & \mathcal{D}_{t+\delta} & \longleftrightarrow & \mathcal{D}_{t+3\delta} \end{array}$$

# Non-functoriality of the persistence barcode

## Proposition

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}$  sending each persistence module to its barcode.*

# Non-functoriality of the persistence barcode

## Proposition

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}$  sending each persistence module to its barcode.*

## Lemma

*There exists no functor  $\mathbf{Vect} \rightarrow \mathbf{Mch}$  sending each vector space of dimension  $d$  to a set of cardinality  $d$ .*

# Structure of submodules and quotient modules

## Proposition (B, Lesnick 2013)

For a persistence submodule  $K \subseteq M$ :

- $B(K)$  is obtained from  $B(M)$  by moving left endpoints to the right,



# Structure of submodules and quotient modules

## Proposition (B, Lesnick 2013)

For a persistence submodule  $K \subseteq M$ :

- $B(K)$  is obtained from  $B(M)$  by moving left endpoints to the right,




# Structure of submodules and quotient modules

## Proposition (B, Lesnick 2013)

For a persistence submodule  $K \subseteq M$ :

- $B(K)$  is obtained from  $B(M)$  by moving left endpoints to the right,
- $B(M/K)$  is obtained from  $B(M)$  by moving right endpoints to the left.




# Structure of submodules and quotient modules

## Proposition (B, Lesnick 2013)

For a persistence submodule  $K \subseteq M$ :

- $B(K)$  is obtained from  $B(M)$  by moving left endpoints to the right,
- $B(M/K)$  is obtained from  $B(M)$  by moving right endpoints to the left.




This yields canonical matchings between the barcodes:  
match bars with the same right endpoint (left endpoint)

# Structure of submodules and quotient modules

## Proposition (B, Lesnick 2013)

For a persistence submodule  $K \subseteq M$ :

- $B(K)$  is obtained from  $B(M)$  by moving left endpoints to the right,
- $B(M/K)$  is obtained from  $B(M)$  by moving right endpoints to the left.



This yields canonical matchings between the barcodes:  
match bars with the same right endpoint (left endpoint)

- If multiple bars have same endpoint:  
match in order of decreasing length



## Induced matchings

For any morphism  $f : M \rightarrow N$  between persistence modules:

- decompose into  $M \twoheadrightarrow \text{im } f \hookrightarrow N$

# Induced matchings

For any morphism  $f : M \rightarrow N$  between persistence modules:

- decompose into  $M \twoheadrightarrow \text{im } f \hookrightarrow N$
- $\text{im } f$  is a quotient of  $M$



# Induced matchings

For any morphism  $f : M \rightarrow N$  between persistence modules:

- decompose into  $M \twoheadrightarrow \text{im } f \hookrightarrow N$
- $\text{im } f$  is a quotient of  $M$
- $\text{im } f$  is a submodule of  $N$



# Induced matchings

For any morphism  $f : M \rightarrow N$  between persistence modules:

- decompose into  $M \twoheadrightarrow \text{im } f \hookrightarrow N$
- $\text{im } f$  is a quotient of  $M$
- $\text{im } f$  is a submodule of  $N$
- Composing the canonical matchings yields a matching  $B(f) : B(M) \leftrightarrow B(N)$  induced by  $f$



# Induced matchings

For any morphism  $f : M \rightarrow N$  between persistence modules:

- decompose into  $M \twoheadrightarrow \text{im } f \hookrightarrow N$
- $\text{im } f$  is a quotient of  $M$
- $\text{im } f$  is a submodule of  $N$
- Composing the canonical matchings yields a matching  $B(f) : B(M) \leftrightarrow B(N)$  induced by  $f$



This matching is functorial *for injections*:

$$B(K \hookrightarrow M) = B(L \hookrightarrow M) \circ B(K \hookrightarrow L)$$



# Induced matchings

For any morphism  $f : M \rightarrow N$  between persistence modules:

- decompose into  $M \twoheadrightarrow \text{im } f \hookrightarrow N$
- $\text{im } f$  is a quotient of  $M$
- $\text{im } f$  is a submodule of  $N$
- Composing the canonical matchings yields a matching  $B(f) : B(M) \leftrightarrow B(N)$  induced by  $f$



This matching is functorial *for injections*:

$$B(K \hookrightarrow M) = B(L \hookrightarrow M) \circ B(K \hookrightarrow L)$$



Similarly for surjections.


# The induced matching theorem

Say  $K$  is  $\epsilon$ -trivial if  $K_t \rightarrow K_{t+\epsilon}$  is the zero map for all  $t$ .

Define  $M^\epsilon$  by  $M_t^\epsilon = \text{coim}(M_t \rightarrow M_{t+\epsilon})$ .

## Lemma

Let  $f : M \rightarrow N$  be a morphism such that  $\ker f$  is  $\epsilon$ -trivial.  
Then  $M^\epsilon$  is a quotient module of  $\text{im } f$ .



$$\begin{array}{ccc} M & & \\ \searrow & & \swarrow \\ M^\epsilon & \xleftarrow{\quad} & \text{im } f \end{array}$$

# The induced matching theorem


Say  $K$  is  $\epsilon$ -trivial if  $K_t \rightarrow K_{t+\epsilon}$  is the zero map for all  $t$ .

Define  ${}^\epsilon N$  by  ${}^\epsilon N_t = \text{im}(N_{t-\epsilon} \rightarrow N_t)$ .


## Lemma

Let  $f : M \rightarrow N$  be a morphism such that  $\text{coker } f$  is  $\epsilon$ -trivial.


Then  ${}^\epsilon N$  is a submodule of  $\text{im } f$ .



# The induced matching theorem



# The induced matching theorem



# The induced matching theorem


Theorem (B, Lesnick 2013)

Let  $f : M \rightarrow N$  be a morphism with  $\ker f$  and  $\operatorname{coker} f$   $\epsilon$ -trivial.

Then each interval of length  $\geq \epsilon$  is matched by  $B(f)$ .

If  $B(f)$  matches  $[b, d) \in B(M)$  to  $[b', d') \in B(N)$ , then

$b' \leq b \leq b' + \epsilon$  and  $d - \epsilon \leq d' \leq d$ .




# The induced matching theorem

Let  $f : M \rightarrow N(\delta)$  be an interleaving morphism.

Then  $\ker f$  and  $\text{coker } f$  are  $2\delta$ -trivial.

## Corollary

A  $\delta$ -interleaving between persistence modules induces a  $\delta$ -matching of their persistence barcodes.



# Stability via induced matchings



Thanks for your attention!

# Stability via induced matchings




$B(M)$



$B(N(\delta))$

Thanks for your attention!

# Stability via induced matchings




Thanks for your attention!

# Stability via induced matchings




Thanks for your attention!

# Stability via induced matchings



Thanks for your attention!

# Stability via induced matchings



Thanks for your attention!