


Topological simplification of functions on surfaces

Ulrich Bauer

Georg-August-Universität Göttingen

June 28, 2011







Goal

Given a function f on a surface and $\delta > 0$, find a function f_δ that:

- ▶ minimizes number of critical points
- ▶ stays close to input function: $\|f_\delta - f\|_\infty \leq \delta$

Goal

Given a function f on a surface and $\delta > 0$, find a function f_δ that:

- ▶ minimizes number of critical points
- ▶ stays close to input function: $\|f_\delta - f\|_\infty \leq \delta$

Using:

- ▶ Discrete Morse theory [Forman 1998]
 - ▶ provides notion of critical point in the discrete setting
- ▶ Homological persistence [Edelsbrunner et al. 2002]
 - ▶ quantifies significance of critical points

Regular CW complexes

Think of generalized simplicial complexes


A finite CW complex is constructed inductively:

Regular CW complexes

Think of generalized simplicial complexes

A finite CW complex is constructed inductively:

- ▶ start with a set of points (0-skeleton)




Regular CW complexes

Think of generalized simplicial complexes

A finite CW complex is constructed inductively:

- ▶ start with a set of points (0-skeleton)
- ▶ construct n -skeleton by attaching n -balls along their boundaries to the $(n - 1)$ -skeleton




Regular CW complexes

Think of generalized simplicial complexes

A finite CW complex is constructed inductively:

- ▶ start with a set of points (0-skeleton)
- ▶ construct n -skeleton by attaching n -balls along their boundaries to the $(n - 1)$ -skeleton




Regular CW complexes

Think of generalized simplicial complexes


A finite CW complex is constructed inductively:

- ▶ start with a set of points (0-skeleton)
- ▶ construct n -skeleton by attaching n -balls along their boundaries to the $(n - 1)$ -skeleton
- ▶ *regular* CW complex: all attaching maps are topological embeddings



Discrete Morse theory [Forman, 1998]


Finite regular CW complex \mathcal{K}



Discrete Morse theory [Forman, 1998]

Finite regular CW complex \mathcal{K}



- Discrete vector field:
 - $V \subset \{(\sigma, \tau) : \sigma \text{ is facet of } \tau\}$ s.t.
no cell in more than one pair



Discrete Morse theory [Forman, 1998]

Finite regular CW complex \mathcal{K}



- ▶ Discrete vector field:
 - ▶ $V \subset \{(\sigma, \tau) : \sigma \text{ is facet of } \tau\}$ s.t.
no cell in more than one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths



Discrete Morse theory [Forman, 1998]

Finite regular CW complex \mathcal{K}



- ▶ Discrete vector field:
 - ▶ $V \subset \{(\sigma, \tau) : \sigma \text{ is facet of } \tau\}$ s.t.
no cell in more than one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths
- ▶ Discrete Morse function:



Discrete Morse theory [Forman, 1998]

Finite regular CW complex \mathcal{K}



- ▶ Discrete vector field:
 - ▶ $V \subset \{(\sigma, \tau) : \sigma \text{ is facet of } \tau\}$ s.t.
no cell in more than one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths
- ▶ Discrete Morse function:
 - ▶ *consistent* with gradient field V , i.e.:



Discrete Morse theory [Forman, 1998]

Finite regular CW complex \mathcal{K}



- ▶ Discrete vector field:
 - ▶ $V \subset \{(\sigma, \tau) : \sigma \text{ is facet of } \tau\}$ s.t.
no cell in more than one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths
- ▶ Discrete Morse function:
 - ▶ *consistent* with gradient field V , i.e.:
For all (σ, τ) with σ facet of τ :
 $f(\sigma) < f(\tau)$



Discrete Morse theory [Forman, 1998]

Finite regular CW complex \mathcal{K}

- ▶ Discrete vector field:
 - ▶ $V \subset \{(\sigma, \tau) : \sigma \text{ is facet of } \tau\}$ s.t.
no cell in more than one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths
- ▶ Discrete Morse function:
 - ▶ *consistent* with gradient field V , i.e.:
For all (σ, τ) with σ facet of τ :
 $f(\sigma) < f(\tau)$, if $(\sigma, \tau) \notin V$
 $f(\sigma) \geq f(\tau)$, if $(\sigma, \tau) \in V$



Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in vector field

Discrete Morse theory [Forman, 1998]



- ▶ Critical cell:
 - ▶ not contained in vector field
 - ▶ Examples in 2D:
minimum (0-cell)



Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:


- ▶ not contained in vector field
- ▶ Examples in 2D:
minimum (0-cell), maximum (2-cell)



Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:


- ▶ not contained in vector field
- ▶ Examples in 2D:
 - minimum (0-cell), maximum (2-cell), saddle (1-cell)



Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:

- ▶ not contained in vector field
- ▶ Examples in 2D:
 - minimum (0-cell), maximum (2-cell), saddle (1-cell)





- ▶ No multi-saddle by definition!

Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*


- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*


- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]



- ▶ *Level subcomplex:*

- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]



- ▶ *Level subcomplex:*
 - ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value
- ▶ Homotopy type (and hence homology) changes only at critical cells



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value
- ▶ Homotopy type (and hence homology) changes only at critical cells



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value
- ▶ Homotopy type (and hence homology) changes only at critical cells



Discrete Morse Theory [Forman, 1998]

- ▶ *Level subcomplex:*



- ▶ union of all closed cells below a certain value
- ▶ Homotopy type (and hence homology) changes only at critical cells



Discrete Morse Theory [Forman, 1998]


- ▶ *Level subcomplex:*

- ▶ union of all closed cells below a certain value
- ▶ Homotopy type (and hence homology) changes only at critical cells





Discrete Morse theory [Forman, 1998]

- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: unique path between two critical cells





Discrete Morse theory [Forman, 1998]

- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: unique path between two critical cells
 - ▶ Reversing vector field along path cancels critical cells





Discrete Morse theory [Forman, 1998]

- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: unique path between two critical cells
 - ▶ Reversing vector field along path cancels critical cells
(of gradient vector fields, *not functions*)



Level subcomplexes


Level subcomplex: union of all closed cells below a certain value



Level subcomplexes


Level subcomplex: union of all closed cells below a certain value

• *





Level subcomplexes

Level subcomplex: union of all closed cells below a certain value





Level subcomplexes

Level subcomplex: union of all closed cells below a certain value





Level subcomplexes

Level subcomplex: union of all closed cells below a certain value





Level subcomplexes

Level subcomplex: union of all closed cells below a certain value





Level subcomplexes

Level subcomplex: union of all closed cells below a certain value





Level subcomplexes

Level subcomplex: union of all closed cells below a certain value





Level subcomplexes

Level subcomplex: union of all closed cells below a certain value



Level subcomplexes



Level subcomplex: union of all closed cells below a certain value



Homotopy type (and hence homology) changes only at critical cells

Level subcomplexes



Level subcomplex: union of all closed cells below a certain value



Homotopy type (and hence homology) changes only at critical cells

Level subcomplexes



Level subcomplex: union of all closed cells below a certain value



Homotopy type (and hence homology) changes only at critical cells

Level subcomplexes



Level subcomplex: union of all closed cells below a certain value



Homotopy type (and hence homology) changes only at critical cells


Level subcomplexes

Level subcomplex: union of all closed cells below a certain value




Homotopy type (and hence homology) changes only at critical cells

Cancelling critical points of a gradient vector field



Cancelling critical points of a gradient vector field




Ascending and descending sets

A gradient vector field V imposes inequalities on values of a consistent function

- ▶ *Ascending set* of a cell σ :


$$\{\rho : g(\rho) \geq g(\sigma) \text{ for any } g \text{ consistent with } V\}$$



Ascending and descending sets

A gradient vector field V imposes inequalities on values of a consistent function

- ▶ *Ascending set* of a cell σ :
 $\{\rho : g(\rho) \geq g(\sigma) \text{ for any } g \text{ consistent with } V\}$
- ▶ *Descending set*:
 $\{\phi : g(\phi) \leq g(\sigma) \text{ for any } g \text{ consistent with } V\}$




Ascending and descending sets


A gradient vector field V imposes inequalities on values of a consistent function

- ▶ *Ascending set* of a cell σ :
 $\{\rho : g(\rho) \geq g(\sigma) \text{ for any } g \text{ consistent with } V\}$
- ▶ *Descending set*:
 $\{\phi : g(\phi) \leq g(\sigma) \text{ for any } g \text{ consistent with } V\}$


V induces partial order \prec_V on cells: write $\sigma \prec_V \rho$ and $\phi \prec_V \sigma$




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function



Cancelling critical points of a function




Pseudo-Morse functions



After cancelation, function is no longer Morse


Pseudo-Morse functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones


Pseudo-Morse functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones
 - ▶ For all (σ, τ) s.t. σ is facet of τ :
 $(\sigma, \tau) \notin V \Rightarrow f(\sigma) \leq f(\tau)$
 $(\sigma, \tau) \in V \Rightarrow f(\sigma) \geq f(\tau)$

Pseudo-Morse functions



After cancelation, function is no longer Morse

- ▶ *pseudo-Morse*: replace strict inequalities by weak ones
 - ▶ For all (σ, τ) s.t. σ is facet of τ :
 $(\sigma, \tau) \notin V \Rightarrow f(\sigma) \leq f(\tau)$
 $(\sigma, \tau) \in V \Rightarrow f(\sigma) \geq f(\tau)$
 - ▶ Gradient vector field is no longer unique in general

A symbolic perturbation scheme

Lemma

A function f is a pseudo-Morse function consistent with V if and only if

for every $\epsilon > 0$ there is a discrete Morse function f_ϵ consistent with V and $\|f_\epsilon - f\|_\infty \leq \epsilon$.

A symbolic perturbation scheme

Lemma

A function f is a pseudo-Morse function consistent with V if and only if

for every $\epsilon > 0$ there is a discrete Morse function f_ϵ consistent with V and $\|f_\epsilon - f\|_\infty \leq \epsilon$.

- ▶ treat the pair (f, V) like a Morse function

A symbolic perturbation scheme

Lemma

A function f is a pseudo-Morse function consistent with V if and only if

for every $\epsilon > 0$ there is a discrete Morse function f_ϵ consistent with V and $\|f_\epsilon - f\|_\infty \leq \epsilon$.

- ▶ treat the pair (f, V) like a Morse function
- ▶ can be extended to situations where *unique* critical values are required

Comparing Discrete and PL Morse theory

Why not just use PL functions?

Comparing Discrete and PL Morse theory

Why not just use PL functions?

Discrete Morse theory is often a more convenient and efficient language

Comparing Discrete and PL Morse theory

Why not just use PL functions?

Discrete Morse theory is often a more convenient and efficient language

- ▶ critical points are always non-degenerate (no multiple saddles)

Comparing Discrete and PL Morse theory

Why not just use PL functions?

Discrete Morse theory is often a more convenient and efficient language

- ▶ critical points are always non-degenerate (no multiple saddles)
- ▶ clear and concise notion of vector fields, gradient paths, ascending/descending sets

Comparing Discrete and PL Morse theory

Why not just use PL functions?

Discrete Morse theory is often a more convenient and efficient language

- ▶ critical points are always non-degenerate (no multiple saddles)
- ▶ clear and concise notion of vector fields, gradient paths, ascending/descending sets
- ▶ captures the combinatorial essence of Morse theory (instead of only discretizing)

Comparing Discrete and PL Morse theory

Why not just use PL functions?

Discrete Morse theory is often a more convenient and efficient language

- ▶ critical points are always non-degenerate (no multiple saddles)
- ▶ clear and concise notion of vector fields, gradient paths, ascending/descending sets
- ▶ captures the combinatorial essence of Morse theory (instead of only discretizing)

We can easily convert one type of function into the other

Comparing Discrete and PL Morse theory

Why not just use PL functions?

Discrete Morse theory is often a more convenient and efficient language

- ▶ critical points are always non-degenerate (no multiple saddles)
- ▶ clear and concise notion of vector fields, gradient paths, ascending/descending sets
- ▶ captures the combinatorial essence of Morse theory (instead of only discretizing)

We can easily convert one type of function into the other
(PL → pseudo-Morse → PL: barycentric subdivision)

Complexity of simplexwise linear simplification

Theorem (Gray et al., 2010; B., 2011)

Topological δ -simplification is NP-hard for simplexwise linear functions.

Complexity of simplexwise linear simplification

Theorem (Gray et al., 2010; B., 2011)

Topological δ -simplification is NP-hard for simplexwise linear functions.

- ▶ simplexwise linear: linear on each simplex of the *given* triangulation
- ▶ piecewise linear: simplexwise linear wrt. *some* triangulation


Complexity of simplexwise linear simplification

Theorem (Gray et al., 2010; B., 2011)

Topological δ -simplification is NP-hard for simplexwise linear functions.

- ▶ simplexwise linear: linear on each simplex of the *given* triangulation
- ▶ piecewise linear: simplexwise linear wrt. *some* triangulation

Difficulty comes from multi-saddles



Back to our problem

We have seen how to cancel 2 critical points from a function

Back to our problem

We have seen how to cancel 2 critical points from a function
Strategy: Cancel several pairs of critical points sequentially

Back to our problem

We have seen how to cancel 2 critical points from a function
Strategy: Cancel several pairs of critical points sequentially

Difficulties:

- ▶ Several possible choices at each step (new choices may appear after each step)

Back to our problem

We have seen how to cancel 2 critical points from a function
Strategy: Cancel several pairs of critical points sequentially

Difficulties:

- ▶ Several possible choices at each step (new choices may appear after each step)
- ▶ Keep in mind the tolerance

Back to our problem

We have seen how to cancel 2 critical points from a function
Strategy: Cancel several pairs of critical points sequentially

Difficulties:

- ▶ Several possible choices at each step (new choices may appear after each step)
- ▶ Keep in mind the tolerance
- ▶ How many critical points can be removed by cancelation?


Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets

Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets


Example: connected components in 1D




Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for sublevel sets



Example: connected components in 1D



Persistence diagrams [Cohen-Steiner et al., 2005]



Persistence diagrams [Cohen-Steiner et al., 2005]



Stability of persistence diagrams

Theorem (Cohen-Steiner et al., 2005)

Let $\|f - g\|_\infty \leq \delta$.

The persistence pairs of f that have persistence $> 2\delta$ can be mapped injectively to the persistence pairs of g .


Stability of persistence diagrams

Theorem (Cohen-Steiner et al., 2005)

Let $\|f - g\|_\infty \leq \delta$.

The persistence pairs of f that have persistence $> 2\delta$ can be mapped injectively to the persistence pairs of g .

Corresponding points p_f, p_g in the persistence diagrams have distance $\|p_f - p_g\|_\infty \leq \delta$.



A bound on number of critical points

Corollary

Let f be a discrete Morse function on a surface and let $\delta > 0$.

Then for every function f_δ with $\|f_\delta - f\|_\infty \leq \delta$ we have:

$\# \text{critical points of } f_\delta$

$\geq \# \text{critical points of } f \text{ with persistence} > 2\delta$

A bound on number of critical points

Corollary

Let f be a discrete Morse function on a surface and let $\delta > 0$.

Then for every function f_δ with $\|f_\delta - f\|_\infty \leq \delta$ we have:

$\# \text{critical points of } f_\delta$

$\geq \# \text{critical points of } f \text{ with persistence} > 2\delta$

Strategy: Cancel all persistence pairs of f with persistence $\leq 2\delta$

- ▶ Is cancellation always allowed?

A bound on number of critical points

Corollary

Let f be a discrete Morse function on a surface and let $\delta > 0$.


Then for every function f_δ with $\|f_\delta - f\|_\infty \leq \delta$ we have:

$$\begin{aligned} & \# \text{ critical points of } f_\delta \\ & \geq \# \text{ critical points of } f \text{ with persistence} > 2\delta \end{aligned}$$

Strategy: Cancel all persistence pairs of f with persistence $\leq 2\delta$

- ▶ Is cancelation always allowed?
- ▶ Order of cancelation?

Canceling two nested persistence pairs



Connecting persistence and Morse theory

Theorem (B., Lange, Wardetzky, 2010)

Consider a persistence pair (σ, τ) of a discrete Morse function on a surface. Then (σ, τ) can be canceled after all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with $f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$ have been canceled:

- (i) there exists a gradient path from τ to σ , and*
- (ii) this path is unique.*

Connecting persistence and Morse theory

Theorem (B., Lange, Wardetzky, 2010)

Consider a persistence pair (σ, τ) of a discrete Morse function on a surface. Then (σ, τ) can be canceled after all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with $f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$ have been canceled:

- (i) there exists a gradient path from τ to σ , and*
- (ii) this path is unique.*

Corollary

On a surface, it is possible to cancel just the persistence pairs with persistence $\leq 2\delta$ (without canceling the other pairs).

Connecting persistence and Morse theory

Theorem (B., Lange, Wardetzky, 2010)

Consider a persistence pair (σ, τ) of a discrete Morse function on a surface. Then (σ, τ) can be canceled after all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with $f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$ have been canceled:

- (i) there exists a gradient path from τ to σ , and*
- (ii) this path is unique.*

Corollary

On a surface, it is possible to cancel just the persistence pairs with persistence $\leq 2\delta$ (without canceling the other pairs).

What about the tolerance? Does the error accumulate?

Tightness of the stability bound

Theorem (B., Lange, Wardetzky, 2010)

Let f be a pseudo-Morse function on a surface and let $\delta > 0$.

Then a nested cancelation sequence yields a function f_δ with:

- ▶ $\|f_\delta - f\|_\infty \leq \delta$.
- ▶ *All persistence pairs of f with persistence $\leq 2\delta$ are canceled.*

Tightness of the stability bound

Theorem (B., Lange, Wardetzky, 2010)

Let f be a pseudo-Morse function on a surface and let $\delta > 0$.

Then a nested cancelation sequence yields a function f_δ with:

- ▶ $\|f_\delta - f\|_\infty \leq \delta$.
- ▶ *All persistence pairs of f with persistence $\leq 2\delta$ are canceled.*

This function achieves the minimal number of critical points.

Proof of the tightness theorem

Lemma

Let V_i be a vector field obtained from nested persistence cancellation and let (σ, τ) be a remaining persistence pair. Then for each cell ρ ,

- ▶ $\rho >_{V_i} \sigma$ implies $f(\rho) \geq f(\sigma)$, and
- ▶ $\rho <_{V_i} \tau$ implies $f(\rho) \leq f(\tau)$.

Proof of the tightness theorem

Lemma

Let V_i be a vector field obtained from nested persistence cancellation and let (σ, τ) be a remaining persistence pair. Then for each cell ρ ,

- ▶ $\rho >_{V_i} \sigma$ implies $f(\rho) \geq f(\sigma)$, and
- ▶ $\rho <_{V_i} \tau$ implies $f(\rho) \leq f(\tau)$.

In simple words: the ascending set of σ (descending set of τ) in the *simplified* vector field V_i is consistent with the *original* function f .

Proof of the tightness theorem

Lemma


Let V_i be a vector field obtained from nested persistence cancellation and let (σ, τ) be a remaining persistence pair. Then for each cell ρ ,

- ▶ $\rho >_{V_i} \sigma$ implies $f(\rho) \geq f(\sigma)$, and
- ▶ $\rho <_{V_i} \tau$ implies $f(\rho) \leq f(\tau)$.


In simple words: the ascending set of σ (descending set of τ) in the *simplified* vector field V_i is consistent with the *original* function f .

Need to consider interplay between dimensions!


An example in 2D



An example in 2D



An example in 2D



On surfaces, cancellation may affect other critical values!

An efficient algorithm

A simplified function f_δ can be computed efficiently:

An efficient algorithm

A simplified function f_δ can be computed efficiently:

- ▶ Computing persistence pairs: $O(\text{sort}(n))$
(modified Kruskal's MST algorithm)

An efficient algorithm

A simplified function f_δ can be computed efficiently:

- ▶ Computing persistence pairs: $O(\text{sort}(n))$
(modified Kruskal's MST algorithm)
- ▶ Computing simplified gradient vector field: $O(n)$
(DFS graph traversal)

An efficient algorithm

A simplified function f_δ can be computed efficiently:

- ▶ Computing persistence pairs: $O(\text{sort}(n))$
(modified Kruskal's MST algorithm)
- ▶ Computing simplified gradient vector field: $O(n)$
(DFS graph traversal)
- ▶ Computing simplified function: $O(n)$
(topological sort by DFS graph traversal)

Combination with energy methods

Recall: simplified vector field V_δ imposes inequalities on simplified function

$\|f_\delta - f\|_\infty \leq \delta$: another set of linear inequalities

Combination with energy methods

Recall: simplified vector field V_δ imposes inequalities on simplified function

$\|f_\delta - f\|_\infty \leq \delta$: another set of linear inequalities

- ▶ defines convex set of solutions

Combination with energy methods


Recall: simplified vector field V_δ imposes inequalities on simplified function

$\|f_\delta - f\|_\infty \leq \delta$: another set of linear inequalities


- ▶ defines convex set of solutions
- ▶ find the “best” solution using your favorite energy functional








Removing local extrema from 3D data



Removing local extrema from 3D data



Future work

- ▶ Connection to total variation denoising
- ▶ Extension to higher dimensions (weaker assumptions on critical points)
- ▶ Connection to singularity theory?

Present work

-  U. Bauer.
Persistence in Discrete Morse Theory.
PhD thesis, University of Göttingen, 2011.
-  U. Bauer, C. Lange, and M. Wardetzky.
Optimal topological simplification of discrete functions on surfaces.
arXiv preprint, 2010. To appear in *Discr. Comp. Geometry*.
[arXiv:1001.1269](https://arxiv.org/abs/1001.1269)
-  U. Bauer, C.-B. Schönlieb, and M. Wardetzky.
Total variation meets topological persistence: A first encounter.
Proceedings of ICNAAM 2010, 1022–1026.
[doi:10.1063/1.3497795](https://doi.org/10.1063/1.3497795)

Past work



R. Forman.

A user's guide to discrete Morse theory.

Sém. Loth. de Combinatoire, B48c:1–35, 2002.



H. Edelsbrunner, D. Letscher, and A. Zomorodian.

Topological Persistence and Simplification.

Discret. Comp. Geometry, 28(4):511–533, 2002.



D. Cohen-Steiner, H. Edelsbrunner, and J. Harer.

Stability of Persistence Diagrams.

Discret. Comp. Geometry, 37(1):103–120, 2007.

Thanks for your attention!

