

Topological simplification of functions on surfaces

Ulrich Bauer¹ Carsten Lange² Max Wardetzky¹

¹Georg-August-Universität Göttingen

²Freie Universität Berlin





© 2009 Smithsonian Institution
Image U.S. Geological Survey
Image © 2009 DigitalGlobe
Image State of Oregon

©2009 Google

Goal

Given a function f on a surface and $\delta > 0$, find a function f_δ that:

- ▶ minimizes number of critical points
- ▶ stays close to input function: $\|f - f_\delta\|_\infty \leq \delta$

Goal


Given a function f on a surface and $\delta > 0$, find a function f_δ that:

- ▶ minimizes number of critical points
- ▶ stays close to input function: $\|f - f_\delta\|_\infty \leq \delta$


Using:

- ▶ Discrete Morse theory [Forman 1998]
 - ▶ provides notion of critical point in the discrete setting
- ▶ Homological persistence [Edelsbrunner et al. 2002]
 - ▶ quantifies significance of critical points


Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance




Morse theory at a glance



Morse theory at a glance




Morse theory at a glance



Discrete Morse theory [Forman, 1998]

Finite CW complex \mathcal{K}




Discrete Morse theory [Forman, 1998]

Finite CW complex \mathcal{K}

- Discrete vector field:



- pairs of cells (σ, τ) (arrows),
 σ is a regular facet of τ
- each cell in at most one pair



Discrete Morse theory [Forman, 1998]

Finite CW complex \mathcal{K}



- ▶ Discrete vector field:
 - ▶ pairs of cells (σ, τ) (arrows),
 σ is a regular facet of τ
 - ▶ each cell in at most one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths



Discrete Morse theory [Forman, 1998]

Finite CW complex \mathcal{K}



- ▶ Discrete vector field:
 - ▶ pairs of cells (σ, τ) (arrows),
 σ is a regular facet of τ
 - ▶ each cell in at most one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths
- ▶ Discrete Morse function:



Discrete Morse theory [Forman, 1998]

Finite CW complex \mathcal{K}



- ▶ Discrete vector field:
 - ▶ pairs of cells (σ, τ) (arrows),
 σ is a regular facet of τ
 - ▶ each cell in at most one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths
- ▶ Discrete Morse function:
 - ▶ facets have lower value...



Discrete Morse theory [Forman, 1998]


Finite CW complex \mathcal{K}

- ▶ Discrete vector field:
 - ▶ pairs of cells (σ, τ) (arrows),
 σ is a regular facet of τ
 - ▶ each cell in at most one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths
- ▶ Discrete Morse function:
 - ▶ facets have lower value...
except when there is an arrow





Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair






Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
 - ▶ Examples: minimum





Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
 - ▶ Examples: minimum, maximum




Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
 - ▶ Examples: minimum, maximum, saddle





Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: unique path between two critical cells



Discrete Morse theory [Forman, 1998]


- ▶ Critical cell:
 - ▶ not contained in any pair
- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: unique path between two critical cells
 - ▶ Reversing vector field along path cancels critical cells



Attracting and repelling sets

A gradient vector field V enforces inequalities on cells


- ▶ Attracting set of a (critical) cell σ :
all cells ρ with $g(\rho) \geq g(\sigma)$ for *any* g consistent with V



Attracting and repelling sets

A gradient vector field V enforces inequalities on cells

- ▶ Attracting set of a (critical) cell σ :
all cells ρ with $g(\rho) \geq g(\sigma)$ for *any* g consistent with V
- ▶ Repelling set: analogously for $g(\rho) \leq g(\sigma)$



Comparing Discrete and PL Morse theory

Discrete Morse theory is often a more convenient and efficient language

- ▶ critical points are always non-degenerate (no multiple saddles)
- ▶ clear and concise notion of vector fields, gradient paths, attracting/repelling sets
- ▶ captures the combinatorial essence of Morse theory (instead of only discretizing)

Comparing Discrete and PL Morse theory

Discrete Morse theory is often a more convenient and efficient language

- ▶ critical points are always non-degenerate (no multiple saddles)
- ▶ clear and concise notion of vector fields, gradient paths, attracting/repelling sets
- ▶ captures the combinatorial essence of Morse theory (instead of only discretizing)

We can convert one type of function to the other

Back to our problem

Aim: Cancel critical points from pseudo-Morse function

To do: Cancellation requires three steps:

- ▶ Identify pairs for cancellation (which pairs?)
- ▶ Reverse gradient vector field
- ▶ Make function consistent to new vector field (how?)


Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces

Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces


Example: connected components of sublevel sets



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces


Example: connected components of sublevel sets



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces


Example: connected components of sublevel sets



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces


Example: connected components of sublevel sets



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces


Example: connected components of sublevel sets



Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces


Example: connected components of sublevel sets





Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces

Example: connected components of sublevel sets




Example: level subcomplexes




Example: level subcomplexes

• *





birth (connected component created)



Example: level subcomplexes





Example: level subcomplexes





Example: level subcomplexes





Example: level subcomplexes





Example: level subcomplexes




Example: level subcomplexes




Example: level subcomplexes





Example: level subcomplexes





birth (1-cycle created)





Example: level subcomplexes




Example: level subcomplexes




Example: level subcomplexes




Example: level subcomplexes



death (1-cycle killed)



Example: level subcomplexes



Persistence pair: homology *born* at value 9 *dies* at 12

The lifespan is called *persistence* of the pair

Connecting persistence and Morse theory

How do the two theories fit together?

Connecting persistence and Morse theory

How do the two theories fit together?

Lemma

Consider a discrete Morse function on a surface. Assume that all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with $\sigma < \tilde{\sigma} < \tilde{\tau} < \tau$ have been canceled. Then (σ, τ) can be canceled too.

Connecting persistence and Morse theory

How do the two theories fit together?

Lemma

Consider a discrete Morse function on a surface. Assume that all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with $\sigma < \tilde{\sigma} < \tilde{\tau} < \tau$ have been canceled. Then (σ, τ) can be canceled too.

Corollary

All persistence pairs with persistence $\leq 2\delta$ can be canceled.

Connecting persistence and Morse theory

How do the two theories fit together?

Lemma



Consider a discrete Morse function on a surface. Assume that all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with $\sigma < \tilde{\sigma} < \tilde{\tau} < \tau$ have been canceled. Then (σ, τ) can be canceled too.

Corollary


All persistence pairs with persistence $\leq 2\delta$ can be canceled.

Next: how can we change the function accordingly?


Canceling two nested persistence pairs




Cancelling a persistence pair




Cancelling a persistence pair




Cancelling a persistence pair




Cancelling a persistence pair




Cancelling a persistence pair



Cancelling a persistence pair



Cancelling a persistence pair



Main result

Theorem


Let f be a discrete Morse function on a surface and let $\delta > 0$.

Then there exists a function f_δ with:

- ▶ $\|f - f_\delta\|_\infty \leq \delta$.
- ▶ *All persistence pairs of f with persistence $\leq 2\delta$ are canceled.*

This function achieves the minimal number of critical points.


Comparison with other methods




An efficient algorithm

- ▶ Computing persistence pairs: $O(\text{sort}(n))$
(Kruskal's algorithm)
- ▶ Computing simplified gradient vector field: $O(n)$
(graph traversal)
- ▶ Computing simplified function: $O(n)$
(topological sort)



Example: Simplification of terrain





Example: Simplification of terrain





Example: Medical images



Example: Medical images



Example: Medical images



... thanks for your attention!