

# Persistent Homology and the Stability Theorem

Ulrich Bauer

TUM

February 19, 2018

TAGS – Linking Topology to Algebraic Geometry and Statistics



21



?





2||




?































# What is persistent homology?





# What is persistent homology?





# What is persistent homology?





# What is persistent homology?





# What is persistent homology?



# What is persistent homology?



# What is persistent homology?



# What is persistent homology?

# What is persistent homology?

Persistent homology is the homology of a filtration

# What is persistent homology?

Persistent homology is the homology of a filtration

- ▶ A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - ▶  $\mathbf{R}$  is the poset category of  $(\mathbb{R}, \leq)$
  - ▶ A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - ▶ An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$

# What is persistent homology?

Persistent homology is the homology of a filtration

- ▶ A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - ▶  $\mathbf{R}$  is the poset category of  $(\mathbb{R}, \leq)$
  - ▶ A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - ▶ An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- ▶ Consider homology with coefficients in a field (often  $\mathbb{Z}_2$ )  
 $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$

# What is persistent homology?

Persistent homology is the homology of a filtration

- ▶ A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - ▶  $\mathbf{R}$  is the poset category of  $(\mathbb{R}, \leq)$
  - ▶ A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - ▶ An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- ▶ Consider homology with coefficients in a field (often  $\mathbb{Z}_2$ )  
 $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- ▶ Persistent homology is a diagram  $M : \mathbf{R} \rightarrow \mathbf{Vect}$   
*(persistence module)*

# Homology inference

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $X$  with  $H_*(X) \cong H_*(\Omega)$ .*

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $X$  with  $H_*(X) \cong H_*(\Omega)$ .*

Idea:

- approximate the shape by a thickening  $B_\delta(P)$  covering  $\Omega$

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $X$  with  $H_*(X) \cong H_*(\Omega)$ .*

Idea:

- approximate the shape by a thickening  $B_\delta(P)$  covering  $\Omega$

Requires strong assumptions:


# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $\Omega \subset \mathbb{R}^d$ . Let  $P \subset \Omega$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- the inclusions  $\Omega \hookrightarrow B_\delta(\Omega) \hookrightarrow B_{2\delta}(\Omega)$  preserve homology.

Then  $H_*(\Omega) \cong \text{im } H_*(B_\delta(P) \hookrightarrow B_{2\delta}(P))$ .



# Homological realization

This motivates the *homological realization problem*:

## Problem

*Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that*

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$



# Homological realization

This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$




# Homological realization

This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$



# Homological realization


This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



# Homological realization


This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



# Homological realization


This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



# Homological realization


This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



# Homological realization


This motivates the *homological realization problem*:

## Problem

*Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that*

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



**Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)**

*The homological realization problem is NP-hard, even in  $\mathbb{R}^3$ .*

# Stability


# Persistence and stability: the big picture

Data


point cloud

$P \subset \mathbb{R}^d$


# Persistence and stability: the big picture




# Persistence and stability: the big picture



# Persistence and stability: the big picture



# Persistence and stability: the big picture



# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)


*Let  $\|f - g\|_\infty = \delta$ . Then there exists a matching between the intervals of the persistence barcodes of  $f$  and  $g$  such that*

# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $\|f - g\|_\infty = \delta$ . Then there exists a matching between the intervals of the persistence barcodes of  $f$  and  $g$  such that

- matched intervals have endpoints within distance  $\leq \delta$ , and




# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $\|f - g\|_\infty = \delta$ . Then there exists a matching between the intervals of the persistence barcodes of  $f$  and  $g$  such that

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .




# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  and  $G_t = g^{-1}(-\infty, t]$ .

# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  and  $G_t = g^{-1}(-\infty, t]$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:



# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  and  $G_t = g^{-1}(-\infty, t]$ .


Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:

$$\begin{array}{ccccccc} \dots & \rightarrow & F_t & \hookrightarrow & F_{t+\delta} & \hookrightarrow & F_{t+2\delta} \hookrightarrow F_{t+3\delta} & \dots \rightarrow \\ & & \searrow \times \nearrow & & \searrow \times \nearrow & & \searrow \times \nearrow & \\ \dots & \rightarrow & G_t & \hookrightarrow & G_{t+\delta} & \hookrightarrow & G_{t+2\delta} \hookrightarrow G_{t+3\delta} & \dots \rightarrow \end{array}$$


Applying homology (functor) preserves commutativity

- persistent homology of  $f, g$  yields  
 $\delta$ -interleaved persistence modules  $\mathbf{R} \rightarrow \mathbf{Vect}$


# Geometric interleavings




# Geometric interleavings




# Geometric interleavings




# Geometric interleavings



# Geometric interleavings



# Geometric interleavings



# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- ▶ a vector space  $M_t$  for each  $t \in \mathbb{R}$

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- ▶ a vector space  $M_t$  for each  $t \in \mathbb{R}$
- ▶ a linear map  $m_s^t : M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- ▶ a vector space  $M_t$  for each  $t \in \mathbb{R}$
- ▶ a linear map  $m_s^t : M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
  - ▶ respecting identity:  $m_t^t = \text{id}_{M_t}$
  - ▶ and composition:  $m_s^t \circ m_r^s = m_r^t$

If each  $\dim M_t < \infty$ , we say  $M$  is *pointwise finite dimensional*.

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- ▶ a vector space  $M_t$  for each  $t \in \mathbb{R}$
- ▶ a linear map  $m_s^t : M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
  - ▶ respecting identity:  $m_t^t = \text{id}_{M_t}$
  - ▶ and composition:  $m_s^t \circ m_r^s = m_r^t$

If each  $\dim M_t < \infty$ , we say  $M$  is *pointwise finite dimensional*.

A *morphism*  $f : M \rightarrow N$  is a *natural transformation*:

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- ▶ a vector space  $M_t$  for each  $t \in \mathbb{R}$
- ▶ a linear map  $m_s^t : M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
  - ▶ respecting identity:  $m_t^t = \text{id}_{M_t}$
  - ▶ and composition:  $m_s^t \circ m_r^s = m_r^t$

If each  $\dim M_t < \infty$ , we say  $M$  is *pointwise finite dimensional*.

A *morphism*  $f : M \rightarrow N$  is a *natural transformation*:

- ▶ a linear map  $f_t : M_t \rightarrow N_t$  for each  $t \in \mathbb{R}$

# Persistence modules

A *persistence module*  $M$  is a *diagram* (functor)  $\mathbf{R} \rightarrow \mathbf{Vect}$ :

- ▶ a vector space  $M_t$  for each  $t \in \mathbb{R}$
- ▶ a linear map  $m_s^t : M_s \rightarrow M_t$  for each  $s \leq t$  (*transition maps*)
  - ▶ respecting identity:  $m_t^t = \text{id}_{M_t}$
  - ▶ and composition:  $m_s^t \circ m_r^s = m_r^t$

If each  $\dim M_t < \infty$ , we say  $M$  is *pointwise finite dimensional*.

A *morphism*  $f : M \rightarrow N$  is a *natural transformation*:

- ▶ a linear map  $f_t : M_t \rightarrow N_t$  for each  $t \in \mathbb{R}$
- ▶ morphism and transition maps commute:

$$\begin{array}{ccc} \dots & \rightarrow & M_s \xrightarrow{m_s^t} M_t \rightarrow \dots \\ & & f_s \downarrow \qquad \qquad \qquad \downarrow f_t \\ \dots & \rightarrow & N_s \xrightarrow{n_s^t} N_t \rightarrow \dots \end{array}$$

# Interval Persistence Modules

Let  $\mathbb{K}$  be a field. For an arbitrary interval  $I \subseteq \mathbb{R}$ , define the *interval persistence module*  $\mathbb{K}(I)$  by

$$\mathbb{K}(I)_t = \begin{cases} \mathbb{K} & \text{if } t \in I, \\ 0 & \text{otherwise,} \end{cases}$$

with transition maps of maximal rank.

Schematic example:

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{K} \rightarrow \mathbb{K} \cdots \rightarrow \mathbb{K} \rightarrow 0 \rightarrow 0 \rightarrow \cdots$$

# Barcodes: the structure of persistence modules

Theorem (Krull–Schmidt; Crawley-Boevey 2015)

*Let  $M$  be a pointwise finite-dimensional persistence module.*

# Barcodes: the structure of persistence modules

Theorem (Krull–Schmidt; Crawley-Boevey 2015)

*Let  $M$  be a pointwise finite-dimensional persistence module.*

*Then  $M$  is interval-decomposable:*

*there exists a unique collection of intervals  $B(M)$  such that*

$$M \cong \bigoplus_{I \in B(M)} \mathbb{K}(I).$$

$B(M)$  is called the *barcode* of  $M$ .



# Barcodes: the structure of persistence modules

Theorem (Krull–Schmidt; Crawley-Boevey 2015)

Let  $M$  be a pointwise finite-dimensional persistence module.

Then  $M$  is interval-decomposable:

there exists a unique collection of intervals  $B(M)$  such that

$$M \cong \bigoplus_{I \in B(M)} \mathbb{K}(I).$$

$B(M)$  is called the *barcode* of  $M$ .





- ▶ The decomposition itself is not unique.
- ▶ This is why we use homology with coefficients in a field.

# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:


- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .



# Barcodes as diagrams


# The matching category

A *matching*  $\sigma : S \nrightarrow T$  is a bijection  $S' \rightarrow T'$ , where  $S' \subseteq S$ ,  $T' \subseteq T$ .




# The matching category

A *matching*  $\sigma : S \nrightarrow T$  is a bijection  $S' \rightarrow T'$ , where  $S' \subseteq S$ ,  $T' \subseteq T$ .



Composition of matchings  $\sigma : S \nrightarrow T$  and  $\tau : T \nrightarrow U$ :




# The matching category

A *matching*  $\sigma : S \not\rightarrow T$  is a bijection  $S' \rightarrow T'$ , where  $S' \subseteq S$ ,  $T' \subseteq T$ .



Composition of matchings  $\sigma : S \not\rightarrow T$  and  $\tau : T \not\rightarrow U$ :




Matchings form a category **Mch**

- objects: sets
- morphisms: matchings

# Barcodes as matching diagrams


We can regard a barcode  $B$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :



# Barcodes as matching diagrams

We can regard a barcode  $B$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :


- For each real number  $t$ , let  $B_t$  be the set of intervals of  $B$  that contain  $t$ , and



# Barcodes as matching diagrams

We can regard a barcode  $B$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :

- For each real number  $t$ , let  $B_t$  be the set of intervals of  $B$  that contain  $t$ , and
- for each  $s \leq t$ , define the matching  $B_s \rightarrow B_t$  to be the identity on  $B_s \cap B_t$ .



# Stability via functoriality?

$$\begin{array}{ccc} F_t & \xleftarrow{\quad} & F_{t+2\delta} \\ \curvearrowright & & \curvearrowright \\ G_{t+\delta} & \xleftarrow{\quad} & G_{t+3\delta} \end{array}$$

# Stability via functoriality?

$$\begin{array}{ccc} H_*(F_t) & \longrightarrow & H_*(F_{t+2\delta}) \\ \searrow & & \nearrow \\ H_*(G_{t+\delta}) & \longrightarrow & H_*(G_{t+3\delta}) \end{array}$$

# Stability via functoriality?

$$\begin{array}{ccc} B(H_*(F_t)) & \rightarrow & B(H_*(F_{t+2\delta})) \\ \searrow & & \nearrow \\ & B(H_*(G_{t+\delta})) & \rightarrow B(H_*(G_{t+3\delta})) \end{array}$$

# Stability via functoriality?

$$\begin{array}{ccc} B(H_*(F_t)) & \rightarrow & B(H_*(F_{t+2\delta})) \\ \searrow & & \nearrow \\ & B(H_*(G_{t+\delta})) & \rightarrow B(H_*(G_{t+3\delta})) \end{array}$$



# Non-functoriality of the persistence barcode

## Theorem (B, Lesnick 2015)

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}^R$  sending each persistence module to its barcode.*

# Non-functoriality of the persistence barcode

## Theorem (B, Lesnick 2015)

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}^R$  sending each persistence module to its barcode.*

## Proposition

*There exists no functor  $\mathbf{Vect} \rightarrow \mathbf{Mch}$  sending each vector space of dimension  $d$  to a set of cardinality  $d$ .*

# Induced barcode matchings


# Structure of persistence submodules / quotients

## Proposition

Let  $f : M \rightarrow N$  be a monomorphism of persistence modules:  
each  $f_t : M_t \rightarrow N_t$  is injective.

Then  $f$  induces an injective map  $B(M) \rightarrow B(N)$   
mapping each  $I \in B(M)$  to some  $J \in B(N)$   
with larger or same left and same right endpoint.

$$\begin{array}{ccccccc} \dots & \rightarrow & M_s & \longrightarrow & M_t & \dots & \rightarrow \\ & & \downarrow & & \downarrow & & \\ \dots & \rightarrow & N_s & \longrightarrow & N_t & \dots & \rightarrow \end{array}$$



Dually for epimorphisms (left and right exchanged).

# Induced matchings

For a general morphism  $f : M \rightarrow N$  of persistence modules:  
consider *epi-mono factorization*


$$M \twoheadrightarrow \text{im } f \hookrightarrow N.$$

# Induced matchings

For a general morphism  $f : M \rightarrow N$  of persistence modules:  
consider *epi-mono factorization*


$$M \twoheadrightarrow \text{im } f \hookrightarrow N.$$

- $\text{im } f \hookrightarrow N$  induces injection  $B(\text{im } f) \hookrightarrow B(N)$
- $M \twoheadrightarrow \text{im } f$  induces injection  $B(\text{im } f) \hookrightarrow B(M)$
- compose to a matching  $B(M) \leftrightarrow B(N)$ :




# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



# Stability from interleavings


Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



- ▶  $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- ▶  $M_t \twoheadrightarrow \text{im } f_t$

# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




- ▶  $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- ▶  $M_t \twoheadrightarrow \text{im } f_t$

---

B(N)

# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




- ▶  $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- ▶  $M_t \twoheadrightarrow \text{im } f_t$



# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




- ▶  $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- ▶  $M_t \twoheadrightarrow \text{im } f_t$




# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




- $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- $M_t \twoheadrightarrow \text{im } f_t$



# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




- $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- $M_t \twoheadrightarrow \text{im } f_t$




# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




- $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- $M_t \twoheadrightarrow \text{im } f_t$




# Stability from interleavings

Consider interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



- $\text{im } n_{t-\delta,t+\delta} \hookrightarrow \text{im } f_t \hookrightarrow N_{t+\delta}$
- $M_t \twoheadrightarrow \text{im } f_t$



Sending persistence into  
Hilbert space

# Extending the TDA pipeline

Mapping barcodes into a Hilbert space?

- ▶ desirable for (kernel-based) machine learning methods and statistics
- ▶ stability (Lipschitz continuity): important for reliable predictions
- ▶ inverse stability (bi-Lipschitz): avoid loss of information

# Extending the TDA pipeline

Mapping barcodes into a Hilbert space?

- ▶ desirable for (kernel-based) machine learning methods and statistics
- ▶ stability (Lipschitz continuity): important for reliable predictions
- ▶ inverse stability (bi-Lipschitz): avoid loss of information

Existing kernels for persistence diagrams:

- ▶ stability bounds only for 1–Wasserstein distance
- ▶ Lipschitz constants depend on bound on number and range of bars
- ▶ no bi-Lipschitz bounds known

# Extending the TDA pipeline

Mapping barcodes into a Hilbert space?

- ▶ desirable for (kernel-based) machine learning methods and statistics
- ▶ stability (Lipschitz continuity): important for reliable predictions
- ▶ inverse stability (bi-Lipschitz): avoid loss of information

Existing kernels for persistence diagrams:

- ▶ stability bounds only for 1–Wasserstein distance
- ▶ Lipschitz constants depend on bound on number and range of bars
- ▶ no bi-Lipschitz bounds known

Can we hope for something better?

# No bi-Lipschitz feature maps for persistence

## Theorem (B, Carrière 2018)

*There is no bi-Lipschitz map from the persistence diagrams  
(with the interleaving or any  $p$ -Wasserstein distance)  
into any finite-dimensional Hilbert space,  
even when restricting to bounded range or number of bars.*

# No bi-Lipschitz feature maps for persistence

## Theorem (B, Carrière 2018)

*There is no bi-Lipschitz map from the persistence diagrams  
(with the interleaving or any  $p$ -Wasserstein distance)  
into any finite-dimensional Hilbert space,  
even when restricting to bounded range or number of bars.*

## Theorem (B, Carrière 2018)

*If there was such a bi-Lipschitz map into some Hilbert space,  
the ratio of the Lipschitz constants would have to go to  $\infty$   
together with the bounds on number or range of bars.*

# History

# When was persistent homology invented?

- ▶ [Edelsbrunner/Letscher/Zomorodian 2000]

# When was persistent homology invented?

- ▶ [Edelsbrunner/Letscher/Zomorodian 2000]
- ▶ [Robbins 1999]

# When was persistent homology invented?

- ▶ [Edelsbrunner/Letscher/Zomorodian 2000]
- ▶ [Robbins 1999]
- ▶ [Frosini 1990]

# When was persistent homology invented?

- ▶ [Edelsbrunner/Letscher/Zomorodian 2000]
- ▶ [Robbins 1999]
- ▶ [Frosini 1990]
- ▶ [Leray 1946]?

# When was persistent homology invented first?

# When was persistent homology invented first?

ANNALS OF MATHEMATICS  
Vol. 41, No. 2, April, 1940

## RANK AND SPAN IN FUNCTIONAL TOPOLOGY

BY MARSTON MORSE

(Received August 9, 1939)

### 1. Introduction.

The analysis of functions  $F$  on metric spaces  $M$  of the type which appear in variational theories is made difficult by the fact that the critical limits, such as absolute minima, relative minima, minimax values etc., are in general infinite in number. These limits are associated with relative  $k$ -cycles of various dimensions and are classified as 0-limits, 1-limits etc. The number of  $k$ -limits suitably counted is called the  $k^{\text{th}}$  type number  $m_k$  of  $F$ . The theory seeks to establish relations between the numbers  $m_k$  and the connectivities  $p_k$  of  $M$ . The numbers  $p_k$  are finite in the most important applications. It is otherwise with the numbers  $m_k$ .

The theory has been able to proceed provided one of the following hypotheses is satisfied. The critical limits cluster at most at  $+\infty$ ; the critical points are isolated;<sup>1</sup> the problem is defined by analytic functions; the critical limits taken in their natural order are well-ordered. These conditions are not generally fulfilled. The generality of the theory rested upon the fact that the cases treated approximate in a certain sense the most general problems which it is

# When was persistent homology invented first?

Web Images More... Sign in

Google

Scholar 9 results (0.02 sec) My Citations ▾

All citations Rank and span in functional topology

Articles  Search within citing articles

Case law

My library

Exact homomorphism sequences in homology theory ed.ac.uk [PDF]

JL Kelley, E Pitcher - Annals of Mathematics, 1947 - JSTOR

The developments of this paper stem from the attempts of one of the authors to deduce relations between homology groups of a complex and homology groups of a complex which is its image under a simplicial map. Certain relations were deduced (see [EP 1] and [EP 2] ...)

Any time Cited by 46 Related articles All 3 versions Cite Save More

Since 2016

Since 2015

Since 2012

Custom range...

Sort by relevance Cited by 24 Related articles All 4 versions Cite Save More

Sort by date

include citations

Create alert

Unstable minimal surfaces of higher topological structure

M Morse, CB Tompkins - Duke Math. J., 1941 - projecteuclid.org

1. Introduction. We are concerned with extending the calculus of variations in the large to multiple integrals. The problem of the existence of minimal surfaces of unstable type contains many of the typical difficulties, especially those of a topological nature. Having studied this ...

Cited by 19 Related articles All 2 versions Cite Save

[PDF] Persistence in discrete Morse theory psu.edu [PDF]

U Bauer - 2011 - Citeseer

The goal of this thesis is to bring together two different theories about critical points of a scalar function and their relation to topology: Discrete Morse theory and Persistent Homology. The thesis is divided into three main parts: the first part is an introduction to the different fields, the second part is a detailed description of the two theories, and the third part is an application of the two theories to a specific problem.

27 / 31

# When was persistent homology invented first?

BULLETIN (New Series) OF THE  
AMERICAN MATHEMATICAL SOCIETY  
Volume 3, Number 3, November 1980

## MARSTON MORSE AND HIS MATHEMATICAL WORKS

BY RAOUL BOTT<sup>1</sup>

**1. Introduction.** Marston Morse was born in 1892, so that he was 33 years old when in 1925 his paper *Relations between the critical points of a real-valued function of n independent variables* appeared in the Transactions of the American Mathematical Society. Thus Morse grew to maturity just at the time when the subject of Analysis Situs was being shaped by such masters<sup>2</sup> as Poincaré, Veblen, L. E. J. Brouwer, G. D. Birkhoff, Lefschetz and Alexander, and it was Morse's genius and destiny to discover one of the most beautiful and far-reaching relations between this fledgling and Analysis; a relation which is now known as *Morse Theory*.

In retrospect all great ideas take on a certain simplicity and inevitability, partly because they shape the whole subsequent development of the subject. And so to us, today, Morse Theory seems natural and inevitable. However one only has to glance at these early papers to see what a tour de force it was in the 1920's to go from the mini-max principle of Birkhoff to the Morse inequalities, let alone extend these inequalities to function spaces, so that by the early 30's Morse could establish the theorem that for any Riemann

# When was persistent homology invented first?

inequalities between the dimensions of the  $H_i$  and those of  $H(A_i)$ . Thus the Morse inequalities already reflect a certain part of the “Spectral Sequence magic”, and a modern and tremendously general account of Morse’s work on rank and span in the framework of Leray’s theory was developed by Deheuvels [D] in the 50’s.

Unfortunately both Morse’s and Deheuvel’s papers are not easy reading. On the other hand there is no question in my mind that the papers [36] and [44] constitute another tour de force by Morse. Let me therefore illustrate rather than explain some of the ideas of the rank and span theory in a very simple and tame example.

In the figure which follows I have drawn a homeomorph of  $M = S^1$  in the plane, and I will be studying the height function  $F = y$  on  $M$ .





FIGURE 8

# Morse's functional topology


Key aspects:

- ▶ early precursor of persistence and spectral sequences
- ▶ uses Vietoris homology with field coefficients
- ▶ applies to a broad class of functions on metric spaces  
(not necessarily continuous)
- ▶ inclusions of sublevel sets have finite rank homology  
( $q$ -tame persistent homology)
- ▶ focus on controlled behavior in pathological cases


# Motivation and application: minimal surfaces



(a)




(b)



(c)



(from Dierkes et al.: Minimal Surfaces, Springer 2010)

# Motivation and application: minimal surfaces



(from Dierkes et al.: Minimal Surfaces, Springer 2010)

# Motivation and application: minimal surfaces



(from Dierkes et al.: Minimal Surfaces, Springer 2010)

# Existence of unstable minimal surfaces


Using persistent homology:

- ▶ Number of  $\epsilon$ -persistent critical points (minimal surfaces) is finite for any  $\epsilon > 0$
- ▶ Morse inequalities for  $\epsilon$ -persistent critical points

Theorem (Morse, Tompkins 1939)

*There is a  $C_1$  curve bounding an unstable minimal surface (an index 1 critical point of the area functional).*

# Thanks for your attention!



# Thanks for your attention!

