

# Homological reconstruction and simplification

Ulrich Bauer

IST Austria

April 7, 2013


EMS/DMF Joint Mathematical Weekend


Joint work with:


Dominique Attali, Olivier Devillers, Marc Glisse, André Lieutier




















- ▶ Homology transforms topological spaces to groups



- ▶ Homology transforms topological spaces to groups  
... and continuous maps into group homomorphisms




- ▶ Homology transforms topological spaces to groups  
... and continuous maps into group homomorphisms
- ▶ *Persistent homology group:*  $\text{im } H_*(\text{yellow circle} \hookrightarrow \text{blue circle})$



- ▶ Homology transforms topological spaces to groups  
... and continuous maps into group homomorphisms
- ▶ *Persistent homology group:*  $\text{im } H_*(\text{yellow} \hookrightarrow \text{blue})$
- ▶ Under mild sampling conditions on :

$$\text{im } H_*(\text{yellow} \hookrightarrow \text{blue}) \sim H_*(\text{donut})$$

[Edelsbrunner et al. 2005, Chazal et al. 2005]



- ▶  $A \subset B \subset C$ : thickenings of
- ▶  $X \subset Y$ : thickenings of

$$\begin{array}{ccccc} H_*(A) & & H_*(B) & & H_*(C) \\ \searrow & & \nearrow & & \searrow \\ & H_*(X) & & H_*(Y) & \end{array}$$

- ▶  $A \subset B \subset C$ : thickenings of 
- ▶  $X \subset Y$ : thickenings of 

$$\begin{array}{ccccc}
 H_*(A) & \xleftarrow{\quad} & H_*(B) & \xrightarrow{\quad} & H_*(C) \\
 \searrow & & \nearrow & & \searrow \\
 & H_*(X) & & H_*(Y) &
 \end{array}$$

- ▶  $A \subset B \subset C$ : thickenings of  (with same homology)
- ▶  $X \subset Y$ : thickenings of 

$$\begin{array}{ccccc}
 H_*(A) & \xleftarrow{\quad} & H_*(B) & \xrightarrow{\quad} & H_*(C) \\
 \searrow & & \nearrow & & \curvearrowleft \qquad \curvearrowright \\
 & H_*(X) & & H_*(Y) &
 \end{array}$$

- ▶  $A \subset B \subset C$ : thickenings of  (with same homology)
- ▶  $X \subset Y$ : thickenings of 

$$\begin{array}{ccccc}
 H_*(A) & \xleftarrow{\quad} & H_*(B) & \xrightarrow{\quad} & H_*(C) \\
 \searrow & \nearrow & \curvearrowleft & \searrow & \nearrow \\
 & H_*(X) & \xrightarrow{\quad} & H_*(Y) &
 \end{array}$$

- ▶  $A \subset B \subset C$ : thickenings of  (with same homology)
- ▶  $X \subset Y$ : thickenings of 

$$\begin{array}{ccccc}
 H_*(A) & \xleftarrow{\quad} & H_*(B) & \xrightarrow{\quad} & H_*(C) \\
 \searrow & \nearrow & \uparrow \downarrow \sim & \curvearrowleft & \nearrow \\
 H_*(X) & \twoheadrightarrow & \text{im } H_*(X \hookrightarrow Y) & \hookrightarrow & H_*(Y)
 \end{array}$$


- ▶  $A \subset B \subset C$ : thickenings of  (with same homology)
- ▶  $X \subset Y$ : thickenings of 
- ▶ Then


$$\text{im } H_*(X \hookrightarrow Y) \sim H_*(B)$$


$$\begin{array}{ccccc}
 H_*(A) & \xleftarrow{\quad} & H_*(B) & \xrightarrow{\quad} & H_*(C) \\
 \searrow & \nearrow & \uparrow \downarrow \sim & \curvearrowleft & \nearrow \\
 H_*(X) & \twoheadrightarrow & \text{im } H_*(X \hookrightarrow Y) & \hookrightarrow & H_*(Y)
 \end{array}$$


- ▶  $A \subset B \subset C$ : thickenings of  (with same homology)
- ▶  $X \subset Y$ : thickenings of 
- ▶ Then


$$\text{im } H_*(X \hookrightarrow Y) \sim H_*(B) \sim H_*(\text{doughnut})$$











# Homological reconstruction

(Throughout this talk: homology with field coefficients)

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological reconstruction* of  $(K, L)$ .

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological reconstruction* of  $(K, L)$ .

- ▶ If  $L \subset X \subset K$  then  $\beta_*(X) \geq \text{rank } H_*(L \hookrightarrow K)$

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)


*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological reconstruction* of  $(K, L)$ .

- ▶ If  $L \subset X \subset K$  then  $\beta_*(X) \geq \text{rank } H_*(L \hookrightarrow K)$
- ▶ Equality iff  $X$  is a homological reconstruction


# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist




# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist



- ▶ A homological reconstruction may exist, but not as a subcomplex of  $X$



# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

This means: NP-complete problems, e.g. 3-SAT, can be transformed into homological reconstruction problems (*reduction*)

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

Theorem (Attali, B., Devillers, Glisse, Lieutier)

Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.

This means: NP-complete problems, e.g. 3-SAT, can be transformed into homological reconstruction problems (*reduction*)

3-SAT: satisfiability of Boolean formulas in 3-CNF  
(3-conjunctive normal form)

- $\Phi = (\neg t \wedge u \wedge v) \vee (t \wedge \neg v \wedge \neg w) \vee (\neg u \wedge \neg v \wedge w)$

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

Theorem (Attali, B., Devillers, Glisse, Lieutier)

Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.

This means: NP-complete problems, e.g. 3-SAT, can be transformed into homological reconstruction problems (*reduction*)

3-SAT: satisfiability of Boolean formulas in 3-CNF  
(3-conjunctive normal form)

- ▶  $\Phi = (\neg t \wedge u \wedge v) \vee (t \wedge \neg v \wedge \neg w) \vee (\neg u \wedge \neg v \wedge w)$
- ▶  $t, u, v, \dots$ : variables

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

Theorem (Attali, B., Devillers, Glisse, Lieutier)

Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.

This means: NP-complete problems, e.g. 3-SAT, can be transformed into homological reconstruction problems (*reduction*)

3-SAT: satisfiability of Boolean formulas in 3-CNF  
(3-conjunctive normal form)

- ▶  $\Phi = (\neg t \wedge u \wedge v) \vee (t \wedge \neg v \wedge \neg w) \vee (\neg u \wedge \neg v \wedge w)$
- ▶  $t, u, v, \dots$ : variables
- ▶  $\neg t, u, v, \neg v, \dots$ : literals (variables or negated variables)

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

Theorem (Attali, B., Devillers, Glisse, Lieutier)

Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.

This means: NP-complete problems, e.g. 3-SAT, can be transformed into homological reconstruction problems (*reduction*)

3-SAT: satisfiability of Boolean formulas in 3-CNF  
(3-conjunctive normal form)

- ▶  $\Phi = (\neg t \wedge u \wedge v) \vee (t \wedge \neg v \wedge \neg w) \vee (\neg u \wedge \neg v \wedge w)$
- ▶  $t, u, v, \dots$ : variables
- ▶  $\neg t, u, v, \neg v, \dots$ : literals (variables or negated variables)
- ▶  $(\neg t \wedge u \wedge v), \dots$ : clauses (disjunction of 3 literals)

# Reduction from 3-SAT

# Reduction from 3-SAT

Outline of proof: reduction from 3-SAT

- ▶ Given a Boolean expression  $\Phi$ , construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$

# Reduction from 3-SAT

Outline of proof: reduction from 3-SAT

- ▶ Given a Boolean expression  $\Phi$ , construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$

# Reduction from 3-SAT

Outline of proof: reduction from 3-SAT

- ▶ Given a Boolean expression  $\Phi$ , construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:

# Reduction from 3-SAT

Outline of proof: reduction from 3-SAT

- ▶ Given a Boolean expression  $\Phi$ , construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:
  - ▶  $\Phi$  has a satisfying assignment

# Reduction from 3-SAT

Outline of proof: reduction from 3-SAT


- ▶ Given a Boolean expression  $\Phi$ , construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:
  - ▶  $\Phi$  has a satisfying assignment
  - ▶  $(K, L)$  has a reconstruction

# Reduction from 3-SAT

Outline of proof: reduction from 3-SAT


- ▶ Given a Boolean expression  $\Phi$ , construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:
  - ▶  $\Phi$  has a satisfying assignment
  - ▶  $(K, L)$  has a reconstruction
  - ▶  $(K, L)$  has a reconstruction as a subcomplex of  $K$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  can contain at most one of the edges  $\text{True}_i, \text{False}_i$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  can contain at most one of the edges  $\text{True}_i, \text{False}_i$
- ▶ edges  $\text{True}_i, \text{False}_i$  correspond to possible values of variable  $x_i$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget





- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$
- ▶ edges  $a, b, c$  correspond to literals in a clause

# Reduction from 3-SAT: an example




$(t \vee \neg v \vee \neg w)$




- ▶ For every clause: identify edges *a*, *b*, *c* of clause gadget with edges of variable gadgets corresponding to the literals

# Reduction from 3-SAT: an example




- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals

# Reduction from 3-SAT: an example




- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals

# Reduction from 3-SAT: an example





- For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals
- We can transform a satisfying assignment into a homological reconstruction of  $(K, L)$  and vice versa

# Reduction from 3-SAT: an example



- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals
- ▶ We can transform a satisfying assignment into a homological reconstruction of  $(K, L)$  and vice versa





# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .
- ▶ For any  $X$  with  $L \subset X \subset K$ , there is  $g$  with  $G_{\leq t} = X$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .
- ▶ For any  $X$  with  $L \subset X \subset K$ , there is  $g$  with  $G_{\leq t} = X$ .

Thus we are looking for  $X$  with  $L \subset X \subset K$  minimizing  $\beta(X)$ .

# Simplification of sublevel sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

# Simplification of sublevel sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Sketch of proof:

- ▶ Assume we can minimize  $\beta(G_{\leq t})$  efficiently.

# Simplification of sublevel sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Sketch of proof:

- ▶ Assume we can minimize  $\beta(G_{\leq t})$  efficiently.
- ▶ Given  $(K, L)$ , construct a function  $f$  with  $K = F_{\leq t+\delta}$  and  $L = F_{\leq t-\delta}$ .

# Simplification of sublevel sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Sketch of proof:

- ▶ Assume we can minimize  $\beta(G_{\leq t})$  efficiently.
- ▶ Given  $(K, L)$ , construct a function  $f$  with  $K = F_{\leq t+\delta}$  and  $L = F_{\leq t-\delta}$ .
- ▶ Then  $(K, L)$  has a homological reconstruction  $\iff g$  is a *sublevel set reconstruction*, i.e.,  $\beta(G_{\leq t}) = \text{rank } H_*(L \hookrightarrow K)$ .

# Level set simplification

Let  $F_t = f^{-1}(t)$  denote the  $t$ -level set of  $f$ .

## Problem

*Given a PL function  $f : \mathbb{S}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_t)$ .*

## Simplification of level sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

# Simplification of level sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Idea of proof:

# Simplification of level sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Idea of proof:

- ▶ Construct  $f$  from  $(K, L)$  as before; let  $g$  minimize  $\beta_*(G_t)$

# Simplification of level sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Idea of proof:

- ▶ Construct  $f$  from  $(K, L)$  as before; let  $g$  minimize  $\beta_*(G_t)$
- ▶  $\beta_*(G_t) = \beta_*(G_{<t}) + \beta_*(G_{\leq t})$   
(Mayer-Vietoris, Alexander duality)

# Simplification of level sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Idea of proof:

- ▶ Construct  $f$  from  $(K, L)$  as before; let  $g$  minimize  $\beta_*(G_t)$
- ▶  $\beta_*(G_t) = \beta_*(G_{<t}) + \beta_*(G_{\leq t})$   
(Mayer-Vietoris, Alexander duality)
- ▶  $g$  can be shown to have regular value  $t$ , so  
$$\beta_*(G_t) = 2\beta_*(G_{\leq t})$$

# Simplification of level sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Idea of proof:

- ▶ Construct  $f$  from  $(K, L)$  as before; let  $g$  minimize  $\beta_*(G_t)$
- ▶  $\beta_*(G_t) = \beta_*(G_{<t}) + \beta_*(G_{\leq t})$   
(Mayer-Vietoris, Alexander duality)
- ▶  $g$  can be shown to have regular value  $t$ , so  
$$\beta_*(G_t) = 2\beta_*(G_{\leq t})$$
- ▶ Hence  $g$  also minimizes  $\beta_*(G_{\leq t})$

# Simplification of level sets

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Idea of proof:

- ▶ Construct  $f$  from  $(K, L)$  as before; let  $g$  minimize  $\beta_*(G_t)$
- ▶  $\beta_*(G_t) = \beta_*(G_{<t}) + \beta_*(G_{\leq t})$   
(Mayer-Vietoris, Alexander duality)
- ▶  $g$  can be shown to have regular value  $t$ , so  
$$\beta_*(G_t) = 2\beta_*(G_{\leq t})$$
- ▶ Hence  $g$  also minimizes  $\beta_*(G_{\leq t})$
- ▶ Conversely, a minimizer of  $\beta_*(G_{\leq t})$  also minimizes  $\beta_*(G_t)$

## Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a,b]$ .

**Definition (Edelsbrunner, Morozov, Patel 2011)**

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

## Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a,b]$ .

**Definition (Edelsbrunner, Morozov, Patel 2011)**

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

- ▶ Intuition: a  $\delta$ -robust version of level set homology  $H_*(F_t)$

## Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a,b]$ .

**Definition (Edelsbrunner, Morozov, Patel 2011)**

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

- ▶ Intuition: a  $\delta$ -robust version of level set homology  $H_*(F_t)$
- ▶  $\beta(G_t) \geq \text{rank } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]}) \geq \text{rank } W_*(f, t, \delta)$

# Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a,b]$ .

## Definition (Edelsbrunner, Morozov, Patel 2011)

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

- ▶ Intuition: a  $\delta$ -robust version of level set homology  $H_*(F_t)$
- ▶  $\beta(G_t) \geq \text{rank } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]}) \geq \text{rank } W_*(f, t, \delta)$
- ▶ *Realization* of well group: a function  $g$  with  $\|g - f\|_\infty \leq \delta$  and

$$\beta(G_t) = \text{rank } W_*(f, t, \delta).$$

# Well group realization is level set reconstruction

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Let  $f$  be a PL function on  $\mathbb{S}^n$  with  $t \pm \delta \in (\text{im } f)$ .*

*A PL function  $g$  realizes the well group  $W_*(f, t, \delta)$  if and only if it is a  $(t, \delta)$ -level set reconstruction of  $f$ .*

# Well group realization is level set reconstruction

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Let  $f$  be a PL function on  $\mathbb{S}^n$  with  $t \pm \delta \in (\text{im } f)$ .*

*A PL function  $g$  realizes the well group  $W_*(f, t, \delta)$  if and only if it is a  $(t, \delta)$ -level set reconstruction off.*

*Level set reconstruction:* a simplification achieving the lower bound on  $\beta_*(G_t)$

# Well group realization is level set reconstruction

Theorem (Attali, B., Devillers, Glisse, Lieutier)

*Let  $f$  be a PL function on  $\mathbb{S}^n$  with  $t \pm \delta \in (\text{im } f)$ .*

*A PL function  $g$  realizes the well group  $W_*(f, t, \delta)$  if and only if it is a  $(t, \delta)$ -level set reconstruction off.*

*Level set reconstruction:* a simplification achieving the lower bound on  $\beta_*(G_t)$

## Corollary

*Realization of a well group on  $\mathbb{S}^3$  is NP-hard.*

# Simplification and reconstruction in 2D

Things look brighter in 2D (on surfaces) ...

# Simplification and reconstruction in 2D

Things look brighter in 2D (on surfaces) . . .

**Problem (Function simplification)**

*Given a PL function  $f$  and  $\delta \geq 0$ , find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing the number of critical points.*

# Simplification and reconstruction in 2D

Things look brighter in 2D (on surfaces) . . .

**Problem (Function simplification)**

*Given a PL function  $f$  and  $\delta \geq 0$ , find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing the number of critical points.*

**Theorem (B., Lange, Wardetzky 2012)**

*On surfaces, the function simplification problem can be solved in time  $O(n \log n)$ .*

# Simplification and reconstruction in 2D

Things look brighter in 2D (on surfaces) . . .

## Problem (Function simplification)

*Given a PL function  $f$  and  $\delta \geq 0$ , find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing the number of critical points.*

## Theorem (B., Lange, Wardetzky 2012)

*On surfaces, the function simplification problem can be solved in time  $O(n \log n)$ .*

- ▶ As a consequence, homological reconstruction and (sub)level set simplification can be solved in  $O(n \log n)$

# Simplification and reconstruction in 2D

Things look brighter in 2D (on surfaces) ...

## Problem (Function simplification)

*Given a PL function  $f$  and  $\delta \geq 0$ , find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing the number of critical points.*

## Theorem (B., Lange, Wardetzky 2012)


*On surfaces, the function simplification problem can be solved in time  $O(n \log n)$ .*

- ▶ As a consequence, homological reconstruction and (sub)level set simplification can be solved in  $O(n \log n)$
- ▶ On a surface homological reconstructions always exist






Thanks for your attention!



Thanks for your attention!



... and have a safe trip home!