


Persistence simplification of discrete Morse functions on surfaces

Ulrich Bauer¹ Carsten Lange² Max Wardetzky¹

¹Georg-August-Universität Göttingen

²Freie Universität Berlin

June 11, 2009



Goal

Topological denoising of functions on surfaces

- ▶ minimize number of critical points
- ▶ stay close to input function

Goal


Topological denoising of functions on surfaces

- ▶ minimize number of critical points
- ▶ stay close to input function


Using:

- ▶ Discrete Morse theory [Forman 1998]
 - ▶ provides notion of critical point in the discrete setting
- ▶ Homological persistence [Edelsbrunner et al. 2002]
 - ▶ quantifies significance of critical points


Morse theory at a glance




Morse theory at a glance




Morse theory at a glance



Morse theory at a glance




Morse theory at a glance



Discrete Morse theory [Forman, 1998]


Consider finite CW complex \mathcal{K} .



Discrete Morse theory [Forman, 1998]

Consider finite CW complex \mathcal{K} .



- ▶ Discrete vector field:
 - ▶ a set of *pairs* of cells (σ, τ) ,
where σ is a regular facet of τ
(arrow from σ to τ)
 - ▶ each cell in at most one pair



Discrete Morse theory [Forman, 1998]


Consider finite CW complex \mathcal{K} .

- ▶ Discrete vector field:
 - ▶ a set of *pairs* of cells (σ, τ) , where σ is a regular facet of τ (arrow from σ to τ)
 - ▶ each cell in at most one pair
- ▶ Discrete *gradient* vector field:
 - ▶ no closed paths




Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair





Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: path between two critical cells





Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: path between two critical cells
 - ▶ Reversing vector field along path cancels critical cells




Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: path between two critical cells
 - ▶ Reversing vector field along path cancels critical cells





Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: path between two critical cells
 - ▶ Reversing vector field along path cancels critical cells
 - ▶ If path is unique: preserves gradient vector field property



Discrete Morse theory [Forman, 1998]

- ▶ Critical cell:
 - ▶ not contained in any pair
- ▶ Cancellation of critical cells:
 - ▶ Prerequisite: path between two critical cells
 - ▶ Reversing vector field along path cancels critical cells
 - ▶ If path is unique: preserves gradient vector field property





Discrete Morse function [Forman, 1998]

A function $f : \{\text{cells of } \mathcal{K}\} \rightarrow \mathbb{R}$ and a gradient vector field V_f such that:

- ▶ For all σ facet of τ :
 - ▶ If there is an arrow $\sigma \rightarrow \tau$: $f(\sigma) \geq f(\tau)$
 - ▶ Otherwise: $f(\sigma) < f(\tau)$

f is *consistent* with V_f .





Discrete Morse function (Pseudo-Morse function)

A function $f : \{\text{cells of } \mathcal{K}\} \rightarrow \mathbb{R}$ and a gradient vector field V_f such that:

- ▶ For all σ facet of τ :
 - ▶ If there is an arrow $\sigma \rightarrow \tau$: $f(\sigma) \geq f(\tau)$
 - ▶ Otherwise: $f(\sigma) < f(\tau)$ ($f(\sigma) \leq f(\tau)$)


f is *consistent* with V_f .



Attracting and repelling sets

A gradient vector field V enforces inequalities on cells

- ▶ Attracting set of a (critical) cell σ :
all cells ρ with $g(\rho) \geq g(\sigma)$ for *any* g consistent with V
- ▶ Repelling set: analogously for $g(\rho) \leq g(\sigma)$



Back to our problem

Aim: Cancel critical points from pseudo-Morse function (g, V)

To do: Cancelation requires two steps:

- ▶ Reverse gradient vector field (which pairs?)
- ▶ Make function consistent to new vector field (how?)

Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces

Given:

- ▶ CW complex \mathcal{K}
- ▶ A (Pseudo-)Morse function (f, V)
- ▶ Critical cells $\{\rho_1, \dots, \rho_N\}$ of V such that $f(\rho_i) < f(\rho_{i+1})$

Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces

Given:

- ▶ CW complex \mathcal{K}
- ▶ A (Pseudo-)Morse function (f, V)
- ▶ Critical cells $\{\rho_1, \dots, \rho_N\}$ of V such that $f(\rho_i) < f(\rho_{i+1})$

Morse theory: homology depends on critical cells

Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces

Given:

- ▶ CW complex \mathcal{K}
- ▶ A (Pseudo-)Morse function (f, V)
- ▶ Critical cells $\{\rho_1, \dots, \rho_N\}$ of V such that $f(\rho_i) < f(\rho_{i+1})$

Morse theory: homology depends on critical cells

- ▶ Level Subcomplex $\mathcal{K}(\rho)$:
all cells ϕ with $f(\phi) \leq f(\rho)$ and their faces

Persistent homology [Edelsbrunner et al., 2002]

Investigate change of homology for growing spaces



Given:

- ▶ CW complex \mathcal{K}
- ▶ A (Pseudo-)Morse function (f, V)
- ▶ Critical cells $\{\rho_1, \dots, \rho_N\}$ of V such that $f(\rho_i) < f(\rho_{i+1})$

Morse theory: homology depends on critical cells

- ▶ Level Subcomplex $\mathcal{K}(\rho)$:
all cells ϕ with $f(\phi) \leq f(\rho)$ and their faces
- ▶ Investigate change of homology of $\mathcal{K}(\rho_i)$ as i increases

Example: level subcomplexes



Example: Persistent homology

•


$$\mathcal{K}(\rho_1)$$

$$\beta_0 = 1$$

$$\beta_1 = 0$$

ρ_1 positive cell

Example: Persistent homology




$\mathcal{K}(\rho_1)$

$\beta_0 = 1$

$\beta_1 = 0$

ρ_1 positive cell

Example: Persistent homology



$\mathcal{K}(\rho_1)$

$\beta_0 = 1$

$\beta_1 = 0$

ρ_1 positive cell


$\mathcal{K}(\rho_2)$

$\beta_0 = 1$

$\beta_1 = 1$

ρ_2 positive cell

Example: Persistent homology



$\mathcal{K}(\rho_1)$

$\beta_0 = 1$

$\beta_1 = 0$

ρ_1 positive cell


$\mathcal{K}(\rho_2)$

$\beta_0 = 1$


$\beta_1 = 1$

ρ_2 positive cell

Example: Persistent homology






Example: Persistent homology




- ▶ (ρ_2, ρ_3) is a *persistence pair*:
 ρ_3 kills homology created at ρ_2

Example: Persistent homology


			
$\mathcal{K}(\rho_1)$		$\mathcal{K}(\rho_2)$	$\mathcal{K}(\rho_3)$
$\beta_0 = 1$		$\beta_0 = 1$	$\beta_0 = 1$
$\beta_1 = 0$		$\beta_1 = 1$	$\beta_1 = 0$
ρ_1 positive cell		ρ_2 positive cell	ρ_3 negative cell

- ▶ (ρ_2, ρ_3) is a *persistence pair*:
 ρ_3 kills homology created at ρ_2
- ▶ $f(\rho_3) - f(\rho_2)$ is the *persistence* of (ρ_2, ρ_3)


Cancelling a persistence pair




Cancelling a persistence pair





Cancelling a persistence pair



Cancelling a persistence pair



Canceling two nested persistence pairs



The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value
- ▶ For each low-persistence pair (σ, τ) in this order do:

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value
- ▶ For each low-persistence pair (σ, τ) in this order do:
 - ▶ Let $v = \frac{f(\sigma) + f(\tau)}{2}$

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value
- ▶ For each low-persistence pair (σ, τ) in this order do:
 - ▶ Let $v = \frac{f(\sigma) + f(\tau)}{2}$
 - ▶ For each cell ρ in the attracting set of σ :
$$f_{i+1}(\rho) := \min(f_i(\rho), v)$$

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value
- ▶ For each low-persistence pair (σ, τ) in this order do:
 - ▶ Let $v = \frac{f(\sigma) + f(\tau)}{2}$
 - ▶ For each cell ρ in the attracting set of σ :
$$f_{i+1}(\rho) := \min(f_i(\rho), v)$$
 - ▶ For each cell ρ in the repelling set of τ :
$$f_{i+1}(\rho) := \max(f_i(\rho), v)$$

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value
- ▶ For each low-persistence pair (σ, τ) in this order do:
 - ▶ Let $v = \frac{f(\sigma) + f(\tau)}{2}$
 - ▶ For each cell ρ in the attracting set of σ :
$$f_{i+1}(\rho) := \min(f_i(\rho), v)$$
 - ▶ For each cell ρ in the repelling set of τ :
$$f_{i+1}(\rho) := \max(f_i(\rho), v)$$
 - ▶ For all other cells ρ : $f_{i+1}(\rho) := f_i(\rho)$

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value
- ▶ For each low-persistence pair (σ, τ) in this order do:
 - ▶ Let $v = \frac{f(\sigma) + f(\tau)}{2}$
 - ▶ For each cell ρ in the attracting set of σ :
$$f_{i+1}(\rho) := \min(f_i(\rho), v)$$
 - ▶ For each cell ρ in the repelling set of τ :
$$f_{i+1}(\rho) := \max(f_i(\rho), v)$$
 - ▶ For all other cells ρ : $f_{i+1}(\rho) := f_i(\rho)$
 - ▶ Obtain V_{i+1} by reversing V_i along the path from τ to σ

The algorithm

Given a function $f = f_0$ with gradient vector field V_0 :

- ▶ Sort low-persistence pairs ($\leq 2\delta$) w.r.t. maximum value
- ▶ For each low-persistence pair (σ, τ) in this order do:
 - ▶ Let $v = \frac{f(\sigma) + f(\tau)}{2}$
 - ▶ For each cell ρ in the attracting set of σ :
$$f_{i+1}(\rho) := \min(f_i(\rho), v)$$
 - ▶ For each cell ρ in the repelling set of τ :
$$f_{i+1}(\rho) := \max(f_i(\rho), v)$$
 - ▶ For all other cells ρ : $f_{i+1}(\rho) := f_i(\rho)$
 - ▶ Obtain V_{i+1} by reversing V_i along the path from τ to σ
- ▶ The function f_n is the desired function

Natural questions

- ▶ Can we cancel all low-persistence pairs?
 - ▶ Are the assumptions for canceling satisfied?
- ▶ If yes, how close can we stay to the original function?

Main result


Let (g, V) be a discrete pseudo-Morse function on a combinatorial surface and $\delta > 0$.

Then there exists a pseudo-Morse function (g_δ, V_δ) with:


- ▶ $\|g - g_\delta\|_\infty < \delta$
- ▶ All persistence pairs of (g, V) with persistence $< 2\delta$ are canceled

This function achieves the minimal number of critical points.


Comparison with other methods



Original function





after [Edelsbrunner et al. 2006,
Attali et al. 2009]





our method



Example: Simplification of terrain



Example: Medical images



Example: Medical images



... thanks for your attention!