

# Topological Data Analysis


Part III: The Morse theory of  
Čech and Delaunay complexes

Ulrich Bauer

TUM

February 7, 2015

# Connect the dots: topology from geometry



# Connect the dots: topology from geometry




# Connect the dots: topology from geometry




$$\text{Cech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

# Connect the dots: topology from geometry



$$\text{Cech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$


# Connect the dots: topology from geometry



$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

$$\text{Vor}(p, X) = \left\{ a \in \mathbb{R}^n \mid d(a, p) = d(a, X) \right\}$$


# Connect the dots: topology from geometry



$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

$$\text{Vor}(p, X) = \left\{ a \in \mathbb{R}^n \mid d(a, p) = d(a, X) \right\}$$

# Connect the dots: topology from geometry



$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

$$\text{Vor}(p, X) = \left\{ a \in \mathbb{R}^n \mid d(a, p) = d(a, X) \right\}$$

$$\text{Delaunay}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} (B_r(p) \cap \text{Vor}(p, X)) \neq \emptyset \right\}$$

# Čech and Delaunay functions


$X \subset \mathbb{R}^d$ : finite point set (in general position)



# Čech and Delaunay functions

$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$



# Čech and Delaunay functions

$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$



# $\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$

Two functions on simplices  $Q \subseteq X$ :

*$\check{\text{C}}\text{ech function}$*   $f_C(Q)$ :

radius of smallest enclosing sphere of  $Q$



# $\check{\text{C}}\text{ech}$ and Delaunay functions

$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$


Two functions on simplices  $Q \subseteq X$ :

*$\check{\text{C}}\text{ech function}$*   $f_C(Q)$ :

radius of smallest enclosing sphere of  $Q$

*Delaunay function*  $f_D(Q)$ :

radius of smallest empty circumsphere



# Čech and Delaunay functions


$X \subset \mathbb{R}^d$ : finite point set (in general position)

*Simplices*  $\Delta(X)$ : nonempty subsets of  $X$

Two functions on simplices  $Q \subseteq X$ :

*Čech function*  $f_C(Q)$ :

radius of smallest enclosing sphere of  $Q$



*Delaunay function*  $f_D(Q)$ :

radius of smallest empty circumsphere

- defined only if  $Q$  has an empty circumsphere:  $Q \in \text{Del}(X)$

# Čech and Delaunay complexes from functions

Define for any radius  $r$ :

# Čech and Delaunay complexes from functions

Define for any radius  $r$ :

- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$

# Čech and Delaunay complexes from functions


Define for any radius  $r$ :

- Čech complex  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$

# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- *Čech complex*  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- *Delaunay–Čech complex*  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- *Čech complex*  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- *Delaunay–Čech complex*  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- *Čech complex*  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- *Delaunay–Čech complex*  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices
- *Delaunay complex ( $\alpha$ -shape, for  $\alpha = r$ )*  $\text{Del}_r = f_D^{-1}(-\infty, r]$



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :


- *Čech complex*  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- *Delaunay–Čech complex*  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices
- *Delaunay complex ( $\alpha$ -shape, for  $\alpha = r$ )*  $\text{Del}_r = f_D^{-1}(-\infty, r]$ 
  - all simplices having an empty circumsphere of radius  $\leq r$



# Čech and Delaunay complexes from functions

Define for any radius  $r$ :

- *Čech complex*  $\text{Cech}_r = f_C^{-1}(-\infty, r]$ 
  - all simplices having an enclosing sphere of radius  $\leq r$
- *Delaunay–Čech complex*  $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$ 
  - restriction of Čech complex to Delaunay simplices
- *Delaunay complex ( $\alpha$ -shape, for  $\alpha = r$ )*  $\text{Del}_r = f_D^{-1}(-\infty, r]$ 
  - all simplices having an empty circumsphere of radius  $\leq r$



# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?

# Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$


- Are all three complexes homotopy equivalent?
- Are they related by a sequence of simplicial collapses?

# Discrete Morse theory

# Simplicial collapses

Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.




# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.



# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.
- We can *collapse*  $K$  to a subcomplex  $L$  by removing a free face  $F$ , along with its proper coface.



# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.
- We can *collapse*  $K$  to a subcomplex  $L$  by removing a free face  $F$ , along with its proper coface.
- $L$  is homotopy equivalent to  $K$ .  
In particular, they have isomorphic homology.



# Simplicial collapses

## Definition (Whitehead 1938)

Let  $K$  be a simplicial complex.

- A *free face* of  $K$  is a simplex that has a unique proper coface.
- We can *collapse*  $K$  to a subcomplex  $L$  by removing a free face  $F$ , along with its proper coface.
- $L$  is homotopy equivalent to  $K$ .  
In particular, they have isomorphic homology.




If there is a sequence of such elementary collapses from  $K$  to  $M$ , we say that  $K$  *collapses* to  $M$  (written as  $K \searrow M$ ).

# Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014)

$\check{\text{C}}\text{ech}$ , Delaunay– $\check{\text{C}}\text{ech}$ , Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




# Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014)

$\check{\text{C}}\text{ech}$ , Delaunay– $\check{\text{C}}\text{ech}$ , Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*

# Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014)

$\check{\text{C}}\text{ech}$ , Delaunay– $\check{\text{C}}\text{ech}$ , Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*
- Also works for weighted point sets

# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a simplicial complex  
is a partition of the simplices  
into singletons and pairs  $\{L, U\}$ ,  
where  $L$  is a face of  $U$  with codimension 1.




# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs  $\{L, U\}$ , where  $L$  is a face of  $U$  with codimension 1.

- indicated by an arrow from  $L$  to  $U$




# Discrete Morse theory

## Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs  $\{L, U\}$ , where  $L$  is a face of  $U$  with codimension 1.

- indicated by an arrow from  $L$  to  $U$




The singletons are called *critical simplices*.

# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes



# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex  
is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$   
are subcomplexes

•  
1

# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes
- the level sets  $f^{-1}(t)$  form a discrete vector field (the *discrete gradient* of  $f$ )




# Discrete Morse theory

## Definition (Forman 1998)

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *discrete Morse function* if for all  $t \in \mathbb{R}$ :


- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes
- the level sets  $f^{-1}(t)$  form a discrete vector field (the *discrete gradient* of  $f$ )

If  $f^{-1}(t) = \{Q\}$  then  $t$  is a *critical value*.



# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**

*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**

*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

Theorem (Forman 1998)

If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**


*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*



Let  $V$  be a discrete gradient field on a simplicial complex  $K$ ,  
and let  $L$  be a subcomplex of  $K$ .

**Corollary**

*If  $K \setminus L$  is the union of some pairs of  $V$ ,  
then  $K \searrow L$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**


*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*



Let  $V$  be a discrete gradient field on a simplicial complex  $K$ ,  
and let  $L$  be a subcomplex of  $K$ .

**Corollary**

*If  $K \setminus L$  is the union of some pairs of  $V$ ,  
then  $K \searrow L$ .*




# Collapses from Morse functions and gradients

Let  $f$  be a discrete Morse function on a simplicial complex  $K$ .

**Theorem (Forman 1998)**


*If  $(s, t]$  contains no critical value of  $f$ ,  
then  $K_t \searrow K_s$ .*



Let  $V$  be a discrete gradient field on a simplicial complex  $K$ ,  
and let  $L$  be a subcomplex of  $K$ .

**Corollary**

*If  $K \setminus L$  is the union of some pairs of  $V$ ,  
then  $K \searrow L$ .*



We say that  $V$  induces the collapse  $K \searrow L$ .


## Unfortunately...

Neither the Čech nor the Delaunay functions  
are discrete Morse functions!


# Unfortunately...

Neither the Čech nor the Delaunay functions are discrete Morse functions!

- Example: two simplices  $Q, Q'$  with  $f_C(Q) = f_C(Q')$  such that neither is a face of the other:



$Q$




$Q'$

# Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




# Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




- indicated by an arrow from  $L$  to  $U$

# Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$



- indicated by an arrow from  $L$  to  $U$


The singletons are called *critical simplices*.

# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes



# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex  
is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$   
are subcomplexes

•  
1

# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :

- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes




# Generalized discrete Morse theory

## Definition

A function  $f : K \rightarrow \mathbb{R}$  on a simplicial complex is a *generalized discrete Morse function* if for  $t \in \mathbb{R}$ :


- the sublevel sets  $K_t = f^{-1}(-\infty, t]$  are subcomplexes
- the level sets  $f^{-1}(t)$  form a generalized vector field (the *discrete gradient* of  $f$ )



# Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :




# Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :

- choose an arbitrary vertex  $x \in U \setminus L$




# Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :

- choose an arbitrary vertex  $x \in U \setminus L$
- partition  $[L, U]$  into facet pairs  $(Q \setminus \{x\}, Q \cup \{x\})$  for all  $Q \in [L, U]$ .




# Refining generalized vector fields

A generalized vector field  $V$  can be refined to a vector field.

For each non-critical interval  $[L, U] \in V$ :

- choose an arbitrary vertex  $x \in U \setminus L$
- partition  $[L, U]$  into facet pairs  $(Q \setminus \{x\}, Q \cup \{x\})$  for all  $Q \in [L, U]$ .



Therefore the collapsing theorems also hold for generalized discrete Morse functions.

# Čech and Delaunay functions

# Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

# Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*

- $f_D(Q) = f_C(Q)$

# Morse theory of Čech and Delaunay complexes

## Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*

- $f_D(Q) = f_C(Q)$
- $Q$  is a critical simplex of  $f_C$

# Morse theory of Čech and Delaunay complexes

## Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*

- $f_D(Q) = f_C(Q)$
- $Q$  is a critical simplex of  $f_C$
- $Q$  is a critical simplex of  $f_D$


# Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function  
are generalized discrete Morse functions.*

*The following are equivalent for each simplex  $Q \subseteq X$ :*


- $f_D(Q) = f_C(Q)$
- $Q$  is a critical simplex of  $f_C$
- $Q$  is a critical simplex of  $f_D$
- $Q$  is a centered Delaunay simplex  
(containing the circumcenter in the interior)



# Čech intervals

## Lemma

*Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff*




$Q$

# Čech intervals


## Lemma

Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff

On  $S \subseteq Q'$



$Q$




On  $S$

# Čech intervals


## Lemma

Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff


$$\text{On } S \subseteq Q' \subseteq \text{Encl } S.$$



$Q$



$\text{On } S$




$\text{Encl } S$

# Čech intervals


## Lemma

Let  $Q \subseteq X$  be a simplex with smallest enclosing sphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest enclosing sphere  $S$  iff


$$Q' \in [\text{On } S, \text{Encl } S].$$



$Q$



$\text{On } S$



$\text{Encl } S$

## The front and back faces of a simplex

Let  $\text{On } S$  denote the points of  $X$  on the sphere  $S$ .

Write the center  $z$  of  $S$  as an affine combination

$$z = \sum_{x \in \text{On } S} \mu_x x, \quad 1 = \sum_{x \in \text{On } S} \mu_x.$$

## The front and back faces of a simplex

Let  $\text{On } S$  denote the points of  $X$  on the sphere  $S$ .

Write the center  $z$  of  $S$  as an affine combination

$$z = \sum_{x \in \text{On } S} \mu_x x, \quad 1 = \sum_{x \in \text{On } S} \mu_x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$

# The front and back faces of a simplex

Let  $\text{On } S$  denote the points of  $X$  on the sphere  $S$ .


Write the center  $z$  of  $S$  as an affine combination

$$z = \sum_{x \in \text{On } S} \mu_x x, \quad 1 = \sum_{x \in \text{On } S} \mu_x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



# The front and back faces of a simplex

Let  $\text{On } S$  denote the points of  $X$  on the sphere  $S$ .



Write the center  $z$  of  $S$  as an affine combination

$$z = \sum_{x \in \text{On } S} \mu_x x, \quad 1 = \sum_{x \in \text{On } S} \mu_x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$






# Delaunay intervals

## Lemma

Let  $Q \subseteq X$  be a simplex with smallest empty circumsphere  $S$ .  
Then  $Q' \subseteq X$  has the same smallest empty circumsphere  $S$  iff

$$Q' \in [\text{Front } S, \text{On } S].$$



## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

minimize  
 $r, z$

$r$

subject to

$$\|z - q\| \leq r, \quad q \in Q,$$

$$\|z - e\| \geq r, \quad e \in E.$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

minimize  
 $r, z$

$r$

subject to

$$\begin{aligned}\|z - q\| \leq r, & \quad q \in Q, \\ \|z - e\| \geq r, & \quad e \in E.\end{aligned}$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

- Čech function: choose  $E = \emptyset$
- Delaunay function: choose  $E = X$

## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

$$\begin{array}{ll} \text{minimize}_{r,z} & r^2 \\ \text{subject to} & \|z - q\|^2 \leq r^2, \quad q \in Q, \\ & \|z - e\|^2 \geq r^2, \quad e \in E. \end{array}$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

- Čech function: choose  $E = \emptyset$
- Delaunay function: choose  $E = X$

## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

$$\begin{array}{lll} \underset{a,z}{\text{minimize}} & \|z\|^2 - a & (r^2 = \|z\|^2 - a) \\ \text{subject to} & \|z - q\|^2 \leq \|z\|^2 - a, & q \in Q, \\ & \|z - e\|^2 \geq \|z\|^2 - a, & e \in E. \end{array}$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

- Čech function: choose  $E = \emptyset$
- Delaunay function: choose  $E = X$

## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

$$\underset{a,z}{\text{minimize}} \quad \|z\|^2 - a$$

$$\begin{aligned} \text{subject to} \quad & \|z\|^2 - 2\langle z, q \rangle + \|q\|^2 \leq \|z\|^2 - a, \quad q \in Q, \\ & \|z\|^2 - 2\langle z, e \rangle + \|e\|^2 \geq \|z\|^2 - a, \quad e \in E. \end{aligned}$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

- Čech function: choose  $E = \emptyset$
- Delaunay function: choose  $E = X$

## Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere  $S$  satisfying certain constraints:

$$\begin{array}{ll} \text{minimize}_{a,z} & \|z\|^2 - a \\ \text{subject to} & 2\langle z, q \rangle \geq \|q\|^2 + a, \quad q \in Q, \\ & 2\langle z, e \rangle \leq \|e\|^2 + a, \quad e \in E. \end{array}$$

Here  $r$  is the radius of the sphere  $S$ , and  $z$  is the center of  $S$ .

- Čech function: choose  $E = \emptyset$
- Delaunay function: choose  $E = X$

# The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem of the form

minimize<sub>x</sub>

$$f(x)$$

subject to

$$g_i(x) \leq 0, \quad i \in I,$$

$$h_j(x) = 0, \quad j \in J,$$

where the function  $f$  is convex and  $g_i, h_j$  are affine.

# The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem of the form

$$\begin{array}{ll}\text{minimize}_x & f(x) \\ \text{subject to} & g_i(x) \leq 0, \quad i \in I, \\ & h_j(x) = 0, \quad j \in J,\end{array}$$

where the function  $f$  is convex and  $g_i, h_j$  are affine.

**Theorem (Karush 1939, Kuhn–Tucker 1951)**

A feasible point  $x$  to the above problem is an optimal solution iff there exist Lagrange multipliers  $(\nu_i)_{i \in I}$  and  $(\lambda_j)_{j \in J}$  such that

$$\nabla f(x) + \sum_{i \in I} \nu_i \nabla g_i(x) + \sum_{j \in J} \lambda_j \nabla h_j(x) = 0, \quad (\text{stationarity})$$

$$\nu_i g_i(x) = 0, \quad i \in I, \quad (\text{complementary slackness})$$

$$\lambda_j \geq 0, \quad j \in J. \quad (\text{dual feasibility})$$

# KKT conditions for the smallest sphere problem

The KKT conditions for our sphere optimization problem are:

## Proposition

*A sphere  $S$  enclosing  $Q$  and excluding  $E$  is minimal iff its center  $z$  can be written as an affine combination*

$$z = \sum_{x \in Q \cup E} \mu_x x, \quad 1 = \sum_{x \in Q \cup E} \mu_x$$

*such that*

- $\mu_x = 0$  for  $x \notin Q \cup E$ ,
- $\mu_x \geq 0$  for  $x \notin E$ , and
- $\mu_x \leq 0$  for  $x \notin Q$ .

# Čech and Delaunay intervals from KKT

## Proposition

*A sphere  $S$  enclosing  $Q$  and excluding  $E$  is minimal iff*

$$z \in \text{Aff}(\text{On } S),$$

$$Q \in [\text{Front } S, \text{Encl } S], \text{ and}$$

$$E \in [\text{Back } S, \text{Excl } S].$$

# $\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

## Proposition

*A sphere  $S$  enclosing  $Q$  and excluding  $E$  is minimal iff*

$$z \in \text{Aff}(\text{On } S),$$

$$Q \in [\text{Front } S, \text{Encl } S], \text{ and}$$

$$E \in [\text{Back } S, \text{Excl } S].$$

## Corollary

*The  $\check{\text{C}}\text{ech}$  intervals are of the form  $[\text{On } S, \text{Encl } S]$ .*

# $\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

## Proposition

*A sphere  $S$  enclosing  $Q$  and excluding  $E$  is minimal iff*

$$z \in \text{Aff}(\text{On } S),$$

$$Q \in [\text{Front } S, \text{Encl } S], \text{ and}$$

$$E \in [\text{Back } S, \text{Excl } S].$$

## Corollary

*The  $\check{\text{C}}\text{ech}$  intervals are of the form  $[\text{On } S, \text{Encl } S]$ .*

*The Delaunay intervals are of the form  $[\text{Front } S, \text{On } S]$ .*

## Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$

## Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$

# Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex*  $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

# Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex*  $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

## Theorem (B, Edelsbrunner 2014)

Let  $E \subseteq E'$ . Then

$$\text{Del}_r(X, E) \downarrow \text{Del}_r(X, E) \cap \text{Del}(X, E') \downarrow \text{Del}_r(X, E').$$

# Selective Delaunay complexes

Define for finite point sets  $X, E \subset \mathbb{R}^d$ :

- *E-Delaunay function*  $f_E(Q)$ :  
radius of smallest sphere enclosing  $Q$  and excluding  $E$ 
  - defined only if  $Q$  has an  $E$ -empty enclosing sphere:  
 $Q \in \text{Del}(X, E)$
- *E-Delaunay complex*  $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

## Theorem (B, Edelsbrunner 2014)

Let  $E \subseteq E'$ . Then

$$\text{Del}_r(X, E) \downarrow \text{Del}_r(X, E) \cap \text{Del}(X, E') \downarrow \text{Del}_r(X, E').$$

Note: choosing  $E = \emptyset$  and  $E' = X$  yields


$$\text{Cech}_r(X) \downarrow \text{DelCech}_r(X) \downarrow \text{Del}_r(X).$$

# Collapsing from Čech to Delaunay

# Collapsing the Delaunay–Čech complex

To construct the collapse  $\text{DelCech}_r \searrow \text{Del}_r$ :


- Consider a non-critical Delaunay simplex  $Q$



# Collapsing the Delaunay–Čech complex

To construct the collapse  $\text{DelCech}_r \searrow \text{Del}_r$ :


- Consider a non-critical Delaunay simplex  $Q$
- There is a point  $p$  inside the Čech sphere and on the Delaunay sphere



# Collapsing the Delaunay–Čech complex

To construct the collapse  $\text{DelCech}_r \searrow \text{Del}_r$ :


- Consider a non-critical Delaunay simplex  $Q$
- There is a point  $p$  inside the Čech sphere and on the Delaunay sphere
- Now  $Q' = Q \setminus \{p\}$  and  $Q'' = Q \cup \{p\}$  have the same Čech and Delaunay sphere



# Collapsing the Delaunay–Čech complex

To construct the collapse  $\text{DelCech}_r \searrow \text{Del}_r$ :

- Consider a non-critical Delaunay simplex  $Q$
- There is a point  $p$  inside the Čech sphere and on the Delaunay sphere
- Now  $Q' = Q \setminus \{p\}$  and  $Q'' = Q \cup \{p\}$  have the same Čech and Delaunay sphere




## Lemma

*The pairs  $(Q', Q'')$  yield a discrete gradient.*

# Collapsing the Delaunay–Čech complex

To construct the collapse  $\text{DelCech}_r \searrow \text{Del}_r$ :

- Consider a non-critical Delaunay simplex  $Q$
- There is a point  $p$  inside the Čech sphere and on the Delaunay sphere
- Now  $Q' = Q \setminus \{p\}$  and  $Q'' = Q \cup \{p\}$  have the same Čech and Delaunay sphere



## Lemma


The pairs  $(Q', Q'')$  yield a discrete gradient.

This gradient induces a collapse  $\text{DelCech}_r \searrow \text{Del}_r$ , for any radius  $r$ .

# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :


- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :


- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :


- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :


- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :

- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :

- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :


- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude more and more points until no feasible sphere exists
- Let  $p$  be the last excluded point



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :


- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude more and more points until no feasible sphere exists
- Let  $p$  be the last excluded point
- Now  $Q' = Q \setminus \{p\}$  and  $Q'' = Q \cup \{p\}$  have the same Čech sphere and no Delaunay sphere



# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :

- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude more and more points until no feasible sphere exists
- Let  $p$  be the last excluded point
- Now  $Q' = Q \setminus \{p\}$  and  $Q'' = Q \cup \{p\}$  have the same Čech sphere and no Delaunay sphere




## Lemma

The pairs  $(Q', Q'')$  yield a discrete gradient.

# Collapsing non-Delaunay simplices

To construct the collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ :

- Consider a non-Delaunay simplex  $Q$  together with its Čech sphere  $S$
- $S$  is also the smallest sphere enclosing  $Q$  and excluding  $\text{Excl } S$
- Exclude more and more points until no feasible sphere exists
- Let  $p$  be the last excluded point
- Now  $Q' = Q \setminus \{p\}$  and  $Q'' = Q \cup \{p\}$  have the same Čech sphere and no Delaunay sphere



## Lemma


The pairs  $(Q', Q'')$  yield a discrete gradient.

This gradient induces a collapse  $\text{Cech}_r \searrow \text{DelCech}_r$ , for any radius  $r$ .

# Weighted point sets

All constructions can be generalized to weighted point sets


- Intuition: spheres with given squared radius
- Instead of circumspheres, consider *orthogonal spheres*



# Wrap complexes


# Wrap complexes

Generalizes and greatly simplifies the surface reconstruction algorithm *Wrap* (Edelsbrunner 1995, Geomagic)



# Wrap complexes


Consider the Delaunay function  $f_D$  of  $X$ .



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .


- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$



# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$




# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$




# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$




# Wrap complexes

Consider the Delaunay function  $f_D$  of  $X$ .

- The face relation induces a partial order on the Delaunay intervals  $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$
- Let  $C \in \text{Crit}_r$  be a critical simplex with value  $f_D(C) \leq r$
- Let  $\downarrow C$  denote the descending set of  $C$

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$



The Delaunay intervals induce a collapse  $\text{Del}_r \searrow \text{Wrap}_r$ .

# Wrapping up

- Čech and Delaunay complexes from Morse functions
- Explicit homotopy equivalence by simplicial collapses
- Simple definition and generalization of *Wrap* complexes