

Sending persistence into space

On the metric distortion of embedding persistence
diagrams into reproducing kernel Hilbert spaces

Ulrich Bauer

TUM

June 28, 2018

Curves & Surfaces, Arcachon

Joint work with Mathieu Carrière (INRIA/Columbia)



\approx



?





\approx




?























































Persistence barcodes and persistence diagrams




The pipeline of topological data analysis



Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)


If two functions $f, g : K \rightarrow \mathbb{R}$ have
distance $\|f - g\|_\infty \leq \delta$, then there exists
a δ -matching of their barcodes.



Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have
distance $\|f - g\|_\infty \leq \delta$, then there exists
a δ -matching of their barcodes.




- ▶ matching of sets X, Y : bijection of subsets $X' \subseteq X, Y' \subseteq Y$

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have
distance $\|f - g\|_\infty \leq \delta$, then there exists
a δ -matching of their barcodes.




- ▶ matching of sets X, Y : bijection of subsets $X' \subseteq X, Y' \subseteq Y$
- ▶ δ -matching of barcodes:

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$, then there exists a δ -matching of their barcodes.




- ▶ matching of sets X, Y : bijection of subsets $X' \subseteq X, Y' \subseteq Y$
- ▶ δ -matching of barcodes:
 - ▶ matched intervals have endpoints within distance $\leq \delta$

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$, then there exists a δ -matching of their barcodes.




- ▶ matching of sets X, Y : bijection of subsets $X' \subseteq X, Y' \subseteq Y$
- ▶ δ -matching of barcodes:
 - ▶ matched intervals have endpoints within distance $\leq \delta$
 - ▶ unmatched intervals have length $\leq 2\delta$

Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

If two functions $f, g : K \rightarrow \mathbb{R}$ have distance $\|f - g\|_\infty \leq \delta$, then there exists a δ -matching of their barcodes.



- ▶ matching of sets X, Y : bijection of subsets $X' \subseteq X, Y' \subseteq Y$
- ▶ δ -matching of barcodes:
 - ▶ matched intervals have endpoints within distance $\leq \delta$
 - ▶ unmatched intervals have length $\leq 2\delta$
- ▶ Bottleneck distance d_∞ : infimum δ admitting a δ -matching


Topological machine learning




Task: shape retrieval



Task: object recognition




Task: texture recognition



⋮




Topological machine learning




Task: shape retrieval



Task: object recognition



Task: texture recognition




- SVM
- PCA
- k-Means

↔ Topological data analysis ↔

↔ Machine learning ↔


Topological machine learning



Linear machine learning methods

Many machine learning methods solve linear geometric problems on vector spaces



- ▶ Principal component analysis (PCA): find orthogonal directions of largest variance



Linear machine learning methods

Many machine learning methods solve linear geometric problems on vector spaces

- ▶ Principal component analysis (PCA): find orthogonal directions of largest variance
- ▶ Support vector machines (SVM): find best-separating hyperplane



Kernels and kernel methods

Let X be a set, and \mathcal{H} a vector space with inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$.

- ▶ Consider a map $\Phi : X \rightarrow \mathcal{H}$
- ▶ Then

$$k(\cdot, \cdot) = \langle \Phi(\cdot), \Phi(\cdot) \rangle_{\mathcal{H}} : X \times X \rightarrow \mathbb{R}$$

is a *kernel* (and Φ is the associated *feature map*)

Kernels and kernel methods

Let X be a set, and \mathcal{H} a vector space with inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$.

- ▶ Consider a map $\Phi : X \rightarrow \mathcal{H}$
- ▶ Then

$$k(\cdot, \cdot) = \langle \Phi(\cdot), \Phi(\cdot) \rangle_{\mathcal{H}} : X \times X \rightarrow \mathbb{R}$$


is a *kernel* (and Φ is the associated *feature map*)

For many linear methods:


- ▶ Feature map Φ not required explicitly
- ▶ No explicit basis for \mathcal{H} required
- ▶ Computations can be performed by evaluating $k(\cdot, \cdot)$

This is called the *kernel trick*.


Persistence scale space kernel




Persistence scale space kernel




Persistence scale space kernel




Persistence scale space kernel




Persistence scale space kernel




Persistence scale space kernel



Persistence scale space kernel



Persistence scale space kernel



Extending the TDA pipeline

Mapping barcodes into a Hilbert space:

- ▶ desirable for (kernel-based) machine learning methods and statistics
- ▶ stability (Lipschitz continuity): important for reliable predictions
- ▶ inverse stability (bi-Lipschitz): avoid loss of information

Metric distortion of the PSS kernel

Let $d_{\text{PSS}}(D_f, D_g)$ be the L^2 distance between smoothings of the persistence diagrams D_f, D_g (i.e., a *kernel distance*).

Theorem (Reininghaus, Huber, B, Kwitt 2015)

For two persistence diagrams D_f and D_g and $\sigma > 0$ we have

$$d_{\text{PSS}}(D_f, D_g) \leq \frac{C}{\sigma} d_1(D_f, D_g).$$

Metric distortion of the PSS kernel

Let $d_{\text{PSS}}(D_f, D_g)$ be the L^2 distance between smoothings of the persistence diagrams D_f, D_g (i.e., a *kernel distance*).

Theorem (Reininghaus, Huber, B, Kwitt 2015)

For two persistence diagrams D_f and D_g and $\sigma > 0$ we have

$$d_{\text{PSS}}(D_f, D_g) \leq \frac{C}{\sigma} d_1(D_f, D_g).$$

- Here: stability with respect to *Wasserstein distance*:

$$d_p(D_f, D_g) = \left(\inf_{\mu: D_f \leftrightarrow D_g} \sum_{x \in D_f} \|x - \mu(x)\|_\infty^p \right)^{\frac{1}{p}}$$

- Note: bottleneck distance $d_B(D_f, D_g) = \lim_{p \rightarrow \infty} d_p(D_f, D_g)$

Sliced Wasserstein distance and kernel

Another construction: the *sliced Wasserstein distance* d_{SW} between persistence diagrams.

Theorem (Carrière, Cuturi, Oudot 2017)

For persistence diagrams D_f and D_g with at most N bars, we have

$$\frac{1}{8N^2} d_1(D_f, D_g) \leq d_{\text{SW}}(D_f, D_g) \leq \sqrt{8} d_1(D_f, D_g).$$

Sliced Wasserstein distance and kernel

Another construction: the *sliced Wasserstein distance* d_{SW} between persistence diagrams.

Theorem (Carrière, Cuturi, Oudot 2017)

For persistence diagrams D_f and D_g with at most N bars, we have

$$\frac{1}{8N^2} d_1(D_f, D_g) \leq d_{\text{SW}}(D_f, D_g) \leq \sqrt{8} d_1(D_f, D_g).$$

- ▶ SW is *conditionally negative definite*. This implies that $k_{\text{SW}}(\cdot, \cdot) = \exp(-\text{SW}(\cdot, \cdot)^2)$ is a kernel.
- ▶ Let d_{kSW} denote the corresponding RKHS distance.
- ▶ Then d_{kSW} is not the same as d_{SW} .
- ▶ But on bounded subspaces, the two are strongly equivalent (bi-Lipschitz).

Conditional bounds

Existing kernels for persistence diagrams:

- ▶ stability bounds only for 1–Wasserstein distance
- ▶ Lipschitz constants depend on bound on number and range of bars
- ▶ no bi-Lipschitz bounds known

Conditional bounds

Existing kernels for persistence diagrams:

- stability bounds only for 1–Wasserstein distance
- Lipschitz constants depend on bound on number and range of bars
- no bi-Lipschitz bounds known

Can we hope for something better?

No bi-Lipschitz feature maps for persistence

Theorem (B, Carrière 2018)

*There is no bi-Lipschitz map from the persistence diagrams
(with the bottleneck or any p -Wasserstein distance)
into any Euclidean space \mathbb{R}^d ,
even when restricting to bounded range or number of bars.*

No bi-Lipschitz feature maps for persistence

Theorem (B, Carrière 2018)

*There is no bi-Lipschitz map from the persistence diagrams
(with the bottleneck or any p -Wasserstein distance)
into any Euclidean space \mathbb{R}^d ,
even when restricting to bounded range or number of bars.*

Theorem (B, Carrière 2018)

*If there was such a bi-Lipschitz map into any Hilbert space,
the ratio of the Lipschitz constants would have to go to ∞
if either bound on the number or range of bars go to ∞ .*

Properties of the space of persistence diagrams

Known properties:

- ▶ The space of persistence diagrams (with p -distance, $p < \infty$) is complete and separable [Mileyko et al. 2011]
- ▶ There are no upper bounds on curvature [Turner et al. 2014]

These properties provide no immediate obstructions to bi-Lipschitz embeddings. But:

Persistence spaces and doubling spaces

Definition

A metric space is *doubling* if every ball of radius r can be covered by a constant number of balls of radius $\frac{r}{2}$.

Persistence spaces and doubling spaces

Definition

A metric space is *doubling* if every ball of radius r can be covered by a constant number of balls of radius $\frac{r}{2}$.

Proposition

Any Euclidean space is doubling.

*The image of a doubling space under
a bi-Lipschitz map is again doubling.*

Persistence spaces and doubling spaces

Definition

A metric space is *doubling* if every ball of radius r can be covered by a constant number of balls of radius $\frac{r}{2}$.


Proposition

Any Euclidean space is doubling.

The image of a doubling space under a bi-Lipschitz map is again doubling.

Theorem (B, Carrière 2018)

The space of persistence diagrams (with p -distance) is not doubling.



Thus, there is no bi-Lipschitz map to Euclidean space.

Mapping from non-compact to compact?

What about embedding into Hilbert space?

- ▶ Consider the set $X = \{Dg_n\}_n$ of persistence diagrams, each Dg_n having one interval of length n .
- ▶ Then X is non-compact and metrically complete.

Lemma

Assume that there was a bi-Lipschitz map from persistence diagrams to some Hilbert space.



Such a map would send X to a compact set.

But this would contradict the fact that X is not compact.

Corollary

If there is a bi-Lipschitz map from a bounded subset of persistence diagrams, the Lipschitz constants have to go to ∞ together with the bounds.


Examples



Empirical metric distortion bounds for persistence diagrams of different cardinalities (uniformly sampled points in upper half-square).

Kernels used: Sliced Wasserstein (red), Persistence Scale Space (blue), Persistence Images (purple); Persistence Landscapes (green), Persistence Vectors (orange)

Examples



Empirical distribution of metric distortion ratios, for image (*spider*), shape (*camel*), and texture (*outex*) data sets.

Kernels used: Sliced Wasserstein (red), Persistence Scale Space (blue), Persistence Images (purple); Persistence Landscapes (green), Persistence Vectors (orange)