

Wrapping clouds

Ulrich Bauer

TUM

Mathematisches Kolloquium, U Osnabrück

January 23, 2018

Joint work with Herbert Edelsbrunner (IST Austria)

Wrapping clouds

The Morse theory of Čech and Delaunay complexes


Ulrich Bauer


TUM


Mathematisches Kolloquium, U Osnabrück


January 23, 2018


Joint work with Herbert Edelsbrunner (IST Austria)












































Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?

Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have


$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?
- Are they related by a sequence of *simplicial collapses*?

Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.




Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex that has a unique proper coface.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex that has a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face and its proper coface.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex that has a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face and its proper coface.
- L is homotopy equivalent to K .
In particular, they have isomorphic homology.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex that has a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face and its proper coface.
- L is homotopy equivalent to K .
In particular, they have isomorphic homology.




If there is a sequence of such elementary collapses from K to M , we say that K *collapses* to M (written as $K \searrow M$).

Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2017)

Čech, Delaunay–Čech, Delaunay, and Wrap complexes are homotopy equivalent through a sequence of collapses

$$\text{Cech}_r X \searrow \text{Cech}_r X \cap \text{Del } X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$




Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2017)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent through a sequence of collapses

$$\text{Cech}_r X \searrow \text{Cech}_r X \cap \text{Del } X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$




- Extends to weighted Delaunay triangulations
- All collapses are induced by a single *discrete gradient field*
- This yields explicit chain maps, inducing isomorphisms in persistent homology

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U




Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:


- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- level sets $f^{-1}(t)$ form a discrete vector field (*gradient of f*)




Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U



A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:


- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- level sets $f^{-1}(t)$ form a discrete vector field (*gradient of f*)



If $f^{-1}(t) = \{Q\}$ then t is a *critical value*.

Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.



Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.



Corollary

K is homotopy-equivalent to a CW complex with one d -cell for each critical d -simplex.

Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.



Corollary

K is homotopy-equivalent to a CW complex with one d -cell for each critical d -simplex.

Let V be a discrete gradient field on a simplicial complex K , and let L be a subcomplex of K .

Corollary

If $K \setminus L$ is a union of pairs of V , then $K \searrow L$.



Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.




Corollary

K is homotopy-equivalent to a CW complex with one d -cell for each critical d -simplex.

Let V be a discrete gradient field on a simplicial complex K , and let L be a subcomplex of K .

Corollary

If $K \setminus L$ is a union of pairs of V , then $K \searrow L$.



Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.



Corollary


K is homotopy-equivalent to a CW complex with one d -cell for each critical d -simplex.

Let V be a discrete gradient field on a simplicial complex K , and let L be a subcomplex of K .

Corollary

If $K \setminus L$ is a union of pairs of V , then $K \searrow L$.

We say that V induces the collapse $K \searrow L$.



Čech and Delaunay functions

Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)




Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)

Two radius functions:

- Čech radius function $f_{\text{Čech}}(Q)$:
radius of *smallest enclosing sphere* of Q




Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)

Two radius functions:

- Čech radius function $f_{\text{Cech}}(Q)$:
radius of *smallest enclosing sphere* of Q
- Delaunay radius function $f_{\text{Del}}(Q)$:
radius of *smallest empty circumsphere*
 - defined only on $\text{Del}(X)$



$\check{\text{C}}$ ech and Delaunay complexes from functions

For any radius r :


- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$



Čech and Delaunay complexes from functions

For any radius r :


- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$
- $\text{DelCech}_r(X) = \text{Del}(X) \cap \text{Cech}_r(X)$
 - restriction of Čech complex to Delaunay simplices



Čech and Delaunay complexes from functions

For any radius r :


- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$
- $\text{DelCech}_r(X) = \text{Del}(X) \cap \text{Cech}_r(X)$
 - restriction of Čech complex to Delaunay simplices
- $\text{Del}_r(X) = f_{\text{Del}}^{-1}(-\infty, r]$
 - simplices with smallest empty circumradius $\leq r$



Čech and Delaunay complexes from functions

For any radius r :

- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$
- $\text{DelCech}_r(X) = \text{Del}(X) \cap \text{Cech}_r(X)$
 - restriction of Čech complex to Delaunay simplices
- $\text{Del}_r(X) = f_{\text{Del}}^{-1}(-\infty, r]$
 - simplices with smallest empty circumradius $\leq r$




Unfortunately...

Neither the Čech nor the Delaunay functions
are discrete Morse functions!


Unfortunately...

Neither the Čech nor the Delaunay functions are discrete Morse functions!

- Example: two simplices Q, Q' with $f_C(Q) = f_C(Q')$ where neither is a face of the other:



Q



Q'


Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex K is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$

- indicated by an arrow from L to U




Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex K is a partition of the simplices into intervals of the face poset:


$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$

- indicated by an arrow from L to U



A *generalized discrete Morse function* $f : K \rightarrow \mathbb{R}$ satisfies:


- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes (for all $t \in \mathbb{R}$)
- the level sets $f^{-1}(t)$ form a generalized vector field (the *discrete gradient* of f)



Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$



Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.



Therefore the collapsing theorems also hold for generalized discrete Morse functions.

Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*


Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

The following are equivalent:


- *Q is a critical simplex of $f_{\text{Čech}}$*
- *Q is a critical simplex of f_{Del}*
- $f_{\text{Čech}}(Q) = f_{\text{Del}}(Q)$
- *Q is a centered Delaunay simplex
(containing the circumcenter in the interior)*



Čech intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff




Q

Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff

On $S \subseteq Q'$



Q




On S

Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$\text{On } S \subseteq Q' \subseteq \text{Encl } S.$$



Q



$\text{On } S$




$\text{Encl } S$

Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$Q' \in [\text{On } S, \text{Encl } S].$$



Q



$\text{On } S$



$\text{Encl } S$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex On $S \subseteq X$.

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.
Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

The front and back faces of a simplex


Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.
Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.
Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$






Delaunay intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff

$$Q' \in [\text{Front } S, \text{On } S].$$



Selective Delaunay complexes

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

minimize
 r, z

r

subject to

$$\|z - q\| \leq r, \quad q \in Q,$$

$$\|z - e\| \geq r, \quad e \in E.$$

Here r is the radius of the sphere S , and z is the center of S .

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

minimize
 r, z

r

subject to

$$\|z - q\| \leq r, \quad q \in Q,$$

$$\|z - e\| \geq r, \quad e \in E.$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{ll} \text{minimize}_{r,z} & r^2 \\ \text{subject to} & \|z - q\|^2 \leq r^2, \quad q \in Q, \\ & \|z - e\|^2 \geq r^2, \quad e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{lll} \text{minimize}_{a,z} & \|z\|^2 - a & (r^2 = \|z\|^2 - a) \\ \text{subject to} & \|z - q\|^2 \leq \|z\|^2 - a, & q \in Q, \\ & \|z - e\|^2 \geq \|z\|^2 - a, & e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\underset{a,z}{\text{minimize}} \quad \|z\|^2 - a$$

$$\begin{aligned} \text{subject to} \quad & \|z\|^2 - 2\langle z, q \rangle + \|q\|^2 \leq \|z\|^2 - a, \quad q \in Q, \\ & \|z\|^2 - 2\langle z, e \rangle + \|e\|^2 \geq \|z\|^2 - a, \quad e \in E. \end{aligned}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{ll} \underset{a,z}{\text{minimize}} & \|z\|^2 - a \\ \text{subject to} & 2\langle z, q \rangle \geq \|q\|^2 + a, \quad \forall q \in Q, \\ & 2\langle z, e \rangle \leq \|e\|^2 + a, \quad \forall e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem of the form

minimize_x

$$f(x)$$

subject to

$$g_j(x) \geq 0, \quad \forall j \in J,$$

$$g_k(x) = 0, \quad \forall k \in K,$$

$$g_l(x) \leq 0, \quad \forall l \in L.$$

where the function f is convex and g_i are affine ($i \in I = J \cup K \cup L$).

The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem of the form

$$\begin{array}{ll} \underset{x}{\text{minimize}} & f(x) \\ \text{subject to} & g_j(x) \geq 0, \quad \forall j \in J, \\ & g_k(x) = 0, \quad \forall k \in K, \\ & g_l(x) \leq 0, \quad \forall l \in L. \end{array}$$

where the function f is convex and g_i are affine ($i \in I = J \cup K \cup L$).

Theorem (Karush 1939, Kuhn–Tucker 1951)

A feasible x is optimal iff there exist $(\lambda_i)_{i \in I}$ such that

$$\nabla f(x) + \sum_{i \in I} \lambda_i \nabla g_i(x) = 0, \quad (\text{stationarity})$$

$$\lambda_i g_i(x) = 0, \quad \forall i \in I, \quad (\text{complementary slackness})$$

$$\begin{aligned} \lambda_j &\leq 0, & \forall j \in J, \\ \lambda_l &\geq 0, & \forall l \in L. \end{aligned} \quad (\text{dual feasibility})$$

KKT conditions for the smallest sphere problem

The KKT conditions for our sphere optimization problem are:

Proposition

A sphere S enclosing Q and excluding E is minimal iff its center z can be written as an affine combination

$$z = \sum_{x \in Q \cup E} \lambda_x x, \quad 1 = \sum_{x \in Q \cup E} \lambda_x$$

such that

- $\lambda_x = 0$ whenever x does not lie on S ,
- $\lambda_x \leq 0$ whenever $x \in E \setminus Q$, and
- $\lambda_x \geq 0$ whenever $x \in Q \setminus E$.

Čech and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

Proposition (Geometric KKT conditions)

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $Q \in [\text{Front } S, \text{Encl } S]$, and
- $E \in [\text{Back } S, \text{Excl } S]$.

$\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The Karush–Kuhn–Tucker optimality conditions yield:

Proposition (Geometric KKT conditions)

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $Q \in [\text{Front } S, \text{Encl } S]$, and
- $E \in [\text{Back } S, \text{Excl } S]$.

Corollary

The $\check{\text{C}}\text{ech}$ intervals are of the form $[\text{On } S, \text{Encl } S]$.

$\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The Karush–Kuhn–Tucker optimality conditions yield:

Proposition (Geometric KKT conditions)

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $Q \in [\text{Front } S, \text{Encl } S]$, and
- $E \in [\text{Back } S, \text{Excl } S]$.

Corollary

The $\check{\text{C}}\text{ech}$ intervals are of the form $[\text{On } S, \text{Encl } S]$.

The Delaunay intervals are of the form $[\text{Front } S, \text{On } S]$.

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if such a sphere exists: $Q \in \text{Del}(X, E)$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if such a sphere exists: $Q \in \text{Del}(X, E)$
- *E-Delaunay complex* $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if such a sphere exists: $Q \in \text{Del}(X, E)$
- *E-Delaunay complex* $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2017)

Let $E \subseteq F \subseteq X$. Then

$$\text{Del}_r(X, E) \downarrow \text{Del}_r(X, E) \cap \text{Del}(X, F) \downarrow \text{Del}_r(X, F).$$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *E-Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if such a sphere exists: $Q \in \text{Del}(X, E)$
- *E-Delaunay complex* $\text{Del}_r(X, E) = f_E^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2017)

Let $E \subseteq F \subseteq X$. Then


$$\text{Del}_r(X, E) \subsetneq \text{Del}_r(X, E) \cap \text{Del}(X, F) \subsetneq \text{Del}_r(X, F).$$

Note: choosing $E = \emptyset$ and $F = X$ yields


$$\text{Cech}_r(X) \subsetneq \text{DelCech}_r(X) \subsetneq \text{Del}_r(X).$$

Connecting different Delaunay complexes

X




Y



Connecting different Delaunay complexes

$B_r(X)$





$B_r(Y)$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



$\text{Del}_r(X)$





$\text{Del}_r(X \cup Y)$



$\text{Del}_r(Y)$


Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



$$\begin{array}{ccc} \text{Del}_r(X) & & \text{Del}_r(X \cup Y) & & \text{Del}_r(Y) \\ \searrow & & \swarrow \simeq & & \swarrow \\ & \text{Del}_r(X \cup Y, X) & & \text{Del}_r(X \cup Y, Y) & \end{array}$$


Collapsing from Čech to Delaunay

Collapsing the Delaunay–Čech complex

Construct gradient pairs inducing the collapse

$\text{DelCech}_r \searrow \text{Del}_r$:

- Consider a non-critical Delaunay simplex Q




Collapsing the Delaunay–Čech complex

Construct gradient pairs inducing the collapse

$\text{DelCech}_r \searrow \text{Del}_r$:

- Consider a non-critical Delaunay simplex Q
- There must be a point p *inside* the Čech and *on* the Delaunay sphere




Collapsing the Delaunay–Čech complex

Construct gradient pairs inducing the collapse

$\text{DelCech}_r \searrow \text{Del}_r$:

- Consider a non-critical Delaunay simplex Q
- There must be a point p *inside* the Čech and *on* the Delaunay sphere
- Now by the KKT conditions:
 $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$
have the same Čech and Delaunay sphere




Collapsing the Delaunay–Čech complex

Construct gradient pairs inducing the collapse

$\text{DelCech}_r \searrow \text{Del}_r$:

- Consider a non-critical Delaunay simplex Q
- There must be a point p *inside* the Čech and *on* the Delaunay sphere
- Now by the KKT conditions:
 $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$
have the same Čech and Delaunay sphere




Lemma

The pairs (Q', Q'') yield a discrete gradient, inducing a collapse $\text{DelCech}_r \searrow \text{Del}_r$ (for any radius r).

Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:


- Consider a non-Delaunay simplex Q together with its Čech sphere S



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:


- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:


- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:


- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:


- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$
- Exclude **more and more points** until no feasible sphere exists



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:


- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$
- Exclude more and more points until no feasible sphere exists
- Let p be the last excluded point



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:


- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$
- Exclude more and more points until no feasible sphere exists
- Let p be the last excluded point
- Now by KKT: $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$ both have the same Čech sphere and no Delaunay sphere



Collapsing non-Delaunay simplices

Construct gradient pairs for the collapse $\text{Cech}_r \searrow \text{DelCech}_r$:

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- By the KKT conditions, S is the smallest sphere enclosing Q and excluding $\text{Excl } S$
- Exclude more and more points until no feasible sphere exists
- Let p be the last excluded point
- Now by KKT: $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$ both have the same Čech sphere and no Delaunay sphere



Lemma

The pairs (Q', Q'') yield a discrete gradient, inducing a collapse $\text{Cech}_r \searrow \text{DelCech}_r$ (for any radius r).

Wrap complexes

Wrap complexes

Generalize and simplify the surface reconstruction method

Wrap (Edelsbrunner 1995, Geomagic)

Wrap complexes

Generalize and simplify the surface reconstruction method


Wrap (Edelsbrunner 1995, Geomagic)



Wrap complexes

Generalize and simplify the surface reconstruction method


Wrap (Edelsbrunner 1995, Geomagic)



Wrap complexes

Generalize and simplify the surface reconstruction method


Wrap (Edelsbrunner 1995, Geomagic)



Wrap complexes

Generalize and simplify the surface reconstruction method

Wrap (Edelsbrunner 1995, Geomagic)




"It would be interesting to elucidate the connection between the two approaches to a discrete Morse theory."

Herbert Edelsbrunner, 2002


Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).



Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).




Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$ (intervals [Front S , On S] of simplices with same Delaunay sphere S).



Define $\text{Wrap}_r(X)$ as the smallest complex

- consisting of intervals of V_D and
- containing all critical simplices with circumradius $\leq r$.

Corollary

The Delaunay intervals induce a collapse $\text{Del}_r \searrow \text{Wrap}_r$.

Wrapping up

What we learned in this talk

- Čech and Delaunay complexes from Morse functions
- Explicit homotopy equivalence by simplicial collapses
- Simple definition and generalization of *Wrap* complexes