


The Morse theory of Čech and Delaunay filtrations

Ulrich Bauer Herbert Edelsbrunner

IST Austria

SoCG 2014

Connect the dots: topology from geometry



Connect the dots: topology from geometry




Connect the dots: topology from geometry




$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

Connect the dots: topology from geometry




$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

Connect the dots: topology from geometry




$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

Connect the dots: topology from geometry



$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

Connect the dots: topology from geometry



$$\text{Čech}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} B_r(p) \neq \emptyset \right\}$$

$$\text{Delaunay}_r(X) = \left\{ Q \subseteq X \mid \bigcap_{p \in Q} (B_r(p) \cap \text{Vor}_X(p)) \neq \emptyset \right\}$$

Čech and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)


Simplices $\Delta(X)$: nonempty subsets of X



$\check{\text{C}}$ ech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X



$\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

Two functions on simplices $Q \subseteq X$:

$\check{\text{C}}\text{ech function}$ $f_C(Q)$:

radius of smallest enclosing sphere of Q



$\check{\text{C}}\text{ech}$ and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X


Two functions on simplices $Q \subseteq X$:

$\check{\text{C}}\text{ech function}$ $f_C(Q)$:

radius of smallest enclosing sphere of Q

Delaunay function $f_D(Q)$:

radius of smallest empty circumsphere



$\check{\text{C}}\text{ech}$ and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X

Two functions on simplices $Q \subseteq X$:

$\check{\text{C}}\text{ech function}$ $f_C(Q)$:

radius of smallest enclosing sphere of Q



Delaunay function $f_D(Q)$:

radius of smallest empty circumsphere

- defined only if Q has an empty circumsphere: $Q \in \text{Del}(X)$

Čech and Delaunay complexes from functions

Define for any radius r :

Čech and Delaunay complexes from functions

Define for any radius r :

- Čech complex $\text{Cech}_r = f_C^{-1}(-\infty, r]$

Čech and Delaunay complexes from functions


Define for any radius r :

- Čech complex $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$

Čech and Delaunay complexes from functions

Define for any radius r :


- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Cech}_r \cap \text{Del}$



Čech and Delaunay complexes from functions

Define for any radius r :


- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Cech}_r \cap \text{Del}$
 - restriction of Čech complex to Delaunay simplices



Čech and Delaunay complexes from functions

Define for any radius r :


- Čech complex $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- Delaunay–Čech complex $\text{DelCech}_r = \text{Cech}_r \cap \text{Del}$
 - restriction of Čech complex to Delaunay simplices
- Delaunay complex (α -complex, for $\alpha = r$) $\text{Del}_r = f_D^{-1}(-\infty, r]$



Čech and Delaunay complexes from functions

Define for any radius r :


- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Cech}_r \cap \text{Del}$
 - restriction of Čech complex to Delaunay simplices
- *Delaunay complex (α -complex, for $\alpha = r$)* $\text{Del}_r = f_D^{-1}(-\infty, r]$
 - all simplices having an empty circumsphere of radius $\leq r$



Čech and Delaunay complexes from functions

Define for any radius r :

- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Cech}_r \cap \text{Del}$
 - restriction of Čech complex to Delaunay simplices
- *Delaunay complex (α -complex, for $\alpha = r$)* $\text{Del}_r = f_D^{-1}(-\infty, r]$
 - all simplices having an empty circumsphere of radius $\leq r$



Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?

Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have


$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?
- Are they related by a sequence of simplicial collapses?

Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.




Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex which is contained in a unique proper coface.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex which is contained in a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face F , along with its proper coface.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex which is contained in a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face F , along with its proper coface.
- L is homotopy equivalent to K .
In particular, they have isomorphic homology.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex which is contained in a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face F , along with its proper coface.
- L is homotopy equivalent to K .
In particular, they have isomorphic homology.




If there is a sequence of these elementary collapses from K to L , we say that K *collapses* to L (written as $K \searrow L$).

Main result: a sequence of collapses

Theorem (B, Edelsbrunner)

Čech, Delaunay–Čech, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




Main result: a sequence of collapses

Theorem (B, Edelsbrunner)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*

Main result: a sequence of collapses

Theorem (B, Edelsbrunner)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*
- The filtrations have isomorphic persistent homology

Main result: a sequence of collapses

Theorem (B, Edelsbrunner)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent. In particular,

$$\text{Cech}_r \searrow \text{DelCech}_r \searrow \text{Del}_r \searrow \text{Wrap}_r.$$




- All collapses are induced by a single *discrete gradient field*
- The filtrations have isomorphic persistent homology
- Also works for weighted point sets

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and *facet pairs* (Q, R) (Q is a face of R with codimension 1).




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and *facet pairs* (Q, R) (Q is a face of R with codimension 1).

The singletons are called *critical simplices*.




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes

Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ form a discrete vector field (the *discrete gradient* of f)




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:


- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ form a discrete vector field (the *discrete gradient* of f)



If $f^{-1}(t) = \{Q\}$ then t is a *critical value*.

Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s (written as $K_t \searrow K_s$).*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s (written as $K_t \searrow K_s$).*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s (written as $K_t \searrow K_s$).*



Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)


*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s (written as $K_t \searrow K_s$).*



Let V be a discrete gradient field on a simplicial complex K ,
and let L be a subcomplex of K .

Corollary

*If $K \setminus L$ is the union of facet pairs of V ,
then $K \searrow L$.*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)


*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s (written as $K_t \searrow K_s$).*



Let V be a discrete gradient field on a simplicial complex K ,
and let L be a subcomplex of K .

Corollary

*If $K \setminus L$ is the union of facet pairs of V ,
then $K \searrow L$.*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)


*If $(s, t]$ contains no critical value of f ,
then K_t collapses to K_s (written as $K_t \searrow K_s$).*



Let V be a discrete gradient field on a simplicial complex K ,
and let L be a subcomplex of K .

Corollary

*If $K \setminus L$ is the union of facet pairs of V ,
then $K \searrow L$.*



We say that V induces the collapse $K \searrow L$.



Unfortunately...

Neither the Čech nor the Delaunay functions
are discrete Morse functions!

Unfortunately...

Neither the Čech nor the Delaunay functions are discrete Morse functions!

- Example: two simplices Q, Q' with $f_C(Q) = f_C(Q')$ that do not form a facet pair:




Generalized discrete Morse theory

Definition (Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}.$$




Generalized discrete Morse theory

Definition (Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}.$$




The singletons are called *critical simplices*.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes

•
1

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes




Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:


- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ form a generalized vector field (the *discrete gradient* of f)



Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into facet pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.



Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into facet pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.



Therefore the collapsing theorems also hold for generalized discrete Morse functions.

Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$

Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$
- Q is a critical simplex of f_C

Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$
- Q is a critical simplex of f_C
- Q is a critical simplex of f_D


Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

The following are equivalent for each simplex $Q \subseteq X$:

- $f_D(Q) = f_C(Q)$
- Q is a critical simplex of f_C
- Q is a critical simplex of f_D
- Q is a centered Delaunay simplex
(containing the circumcenter in the interior)



Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$Q' \in [L, U],$$

where


- L : points of X on S ,
- U : points of X on or inside of S .



Q



L



U

The front face of a simplex

Let $T \subseteq X$ be a simplex with smallest circumsphere S .

Write the center z of S as an affine combination



$$z = \sum_{x \in T} \mu_x x, \quad 1 = \sum_{x \in T} \mu_x.$$

We call

$$\text{front } T = \{x \in T \mid \mu_x > 0\},$$

$$\text{back } T = \{x \in T \mid \mu_x < 0\}$$

the *front face* and the *back face* of T , respectively.



Delaunay intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff


$$Q' \in [F, T],$$

where


- T : points of X on S ,
- F : front face of T .



Q



F



T

Sphere optimization problems

Both Čech and Delaunay function are defined using the smallest sphere S satisfying certain constraints:

$$\underset{r,z}{\text{minimize}} \quad r$$

$$\begin{aligned} \text{subject to} \quad & \|z - q\| \leq r, \quad q \in Q, \\ & \|z - e\| \geq r, \quad e \in E. \end{aligned}$$

Here r is the radius and z is the center of the sphere S .

Sphere optimization problems

Both Čech and Delaunay function are defined using the smallest sphere S satisfying certain constraints:

$$\begin{aligned} & \underset{r,z}{\text{minimize}} && r \\ & \text{subject to} && \|z - q\| \leq r, \quad q \in Q, \\ & && \|z - e\| \geq r, \quad e \in E. \end{aligned}$$


Here r is the radius and z is the center of the sphere S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Weighted point sets

All constructions can be generalized to weighted point sets


- Intuition: spheres with given squared radius
- Instead of circumspheres, consider *orthogonal spheres*



Weighted point sets

All constructions can be generalized to weighted point sets

- Intuition: spheres with given squared radius
- Instead of circumspheres, consider *orthogonal spheres*



The optimization problem is:

$$\underset{r,z}{\text{minimize}} \quad r^2$$


$$\text{subject to} \quad \|z - q\|^2 \leq r^2 + w_q, \quad q \in Q,$$

$$\|z - e\|^2 \geq r^2 + w_e, \quad e \in E.$$

Weighted point sets

All constructions can be generalized to weighted point sets

- Intuition: spheres with given squared radius
- Instead of circumspheres, consider *orthogonal spheres*



The optimization problem (substituting $a = \|z\|^2 - r^2$) is:

$$\underset{a,z}{\text{minimize}} \quad \|z\|^2 - a$$


$$\text{subject to} \quad \|z - q\|^2 \leq \|z\|^2 - a + w_q, \quad q \in Q,$$

$$\|z - e\|^2 \geq \|z\|^2 - a + w_e, \quad e \in E.$$

Weighted point sets

All constructions can be generalized to weighted point sets

- Intuition: spheres with given squared radius
- Instead of circumspheres, consider *orthogonal spheres*



The optimization problem (substituting $a = \|z\|^2 - r^2$) is:

$$\underset{a,z}{\text{minimize}} \quad \|z\|^2 - a$$

$$\text{subject to} \quad 2\langle z, q \rangle - a \geq \|q\|^2 - w_q, \quad q \in Q,$$

$$2\langle z, e \rangle - a \leq \|e\|^2 - w_e, \quad e \in E.$$

The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem

$$\begin{aligned} & \underset{x}{\text{minimize}} && f(x) \\ & \text{subject to} && g_i(x) \leq 0, \quad i \in I, \\ & && h_j(x) = 0, \quad j \in J. \end{aligned}$$

The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem

$$\underset{x}{\text{minimize}} \quad f(x)$$

$$\text{subject to} \quad g_i(x) \leq 0, \quad i \in I,$$

$$h_j(x) = 0, \quad j \in J.$$

Assume that the functions g_i, h_j are affine and f is convex.
Then a feasible point x is an optimal solution iff there exist
Lagrange multipliers $(v_i)_{i \in I}$ and $(\lambda_j)_{j \in J}$ such that

$$\nabla f(x) + \sum_{i \in I} v_i \nabla g_i(x) + \sum_{j \in J} \lambda_j \nabla h_j(x) = 0,$$
$$v_i \geq 0, \quad i \in I,$$

$$v_i g_i(x) = 0, \quad i \in I.$$

KKT conditions for smallest spheres

The KKT conditions for our sphere optimization problem are:

Proposition

A sphere S enclosing Q and empty of E is minimal iff its center z can be written as an affine combination

$$z = \sum_{x \in Q \cup E} \mu_x x, \quad 1 = \sum_{x \in Q \cup E} \mu_x$$

such that

- $\mu_x \geq 0$ for $x \notin E$,
- $\mu_x \leq 0$ for $x \notin Q$, and
- $\mu_x = 0$ for $x \notin S$.

Čech and Delaunay intervals from KKT

Corollary

A sphere S enclosing Q and empty of E is minimal iff

$$z \in \text{aff } T,$$

$$Q \in [F, U], \text{ and}$$

$$E \in [B, W],$$

where


- z : center of S ,
- T, U, W : points of X on S , on or inside S , on or outside of S ,
- F, B : front, back face of T .

Intersections of Čech and Delaunay intervals

Lemma (Excluded Singularity)

The intersection of non-singular Čech and Delaunay intervals is a non-singular interval.

- Consider a non-critical Delaunay simplex Q




Intersections of Čech and Delaunay intervals

Lemma (Excluded Singularity)

The intersection of non-singular Čech and Delaunay intervals is a non-singular interval.

- Consider a non-critical Delaunay simplex Q
- There is a point p inside the Čech sphere and on the Delaunay sphere




Intersections of Čech and Delaunay intervals

Lemma (Excluded Singularity)

The intersection of non-singular Čech and Delaunay intervals is a non-singular interval.

- Consider a non-critical Delaunay simplex Q
- There is a point p inside the Čech sphere and on the Delaunay sphere
- $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$ have the same Čech and Delaunay sphere




Intersections of Čech and Delaunay intervals

Lemma (Excluded Singularity)

The intersection of non-singular Čech and Delaunay intervals is a non-singular interval.

- Consider a non-critical Delaunay simplex Q
- There is a point p inside the Čech sphere and on the Delaunay sphere
- $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$ have the same Čech and Delaunay sphere




Corollary

The pairs (Q', Q'') yield a vector field that induces a collapse $\text{DelCech}_r \searrow \text{Del}_r$.


Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S




Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set
 $E = X \setminus \text{int } S$




Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set $E = X \setminus \text{int } S$
- Exclude more and more points until no feasible sphere exists




Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set $E = X \setminus \text{int } S$
- Exclude more and more points until no feasible sphere exists



Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set $E = X \setminus \text{int } S$
- Exclude more and more points until no feasible sphere exists



Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set $E = X \setminus \text{int } S$
- Exclude more and more points until no feasible sphere exists



Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set $E = X \setminus \text{int } S$
- Exclude more and more points until no feasible sphere exists
- Let p be the last excluded point



Collapsing non-Delaunay simplices

- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set $E = X \setminus \text{int } S$
- Exclude more and more points until no feasible sphere exists
- Let p be the last excluded point
- $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$ have the same Čech sphere and no Delaunay sphere



Collapsing non-Delaunay simplices


- Consider a non-Delaunay simplex Q together with its Čech sphere S
- Start with the excluded set $E = X \setminus \text{int } S$
- Exclude more and more points until no feasible sphere exists
- Let p be the last excluded point
- $Q' = Q \setminus \{p\}$ and $Q'' = Q \cup \{p\}$ have the same Čech sphere and no Delaunay sphere

Corollary

The pairs (Q', Q'') yield a vector field that induces a collapse $\text{Cech}_r \searrow \text{DelCech}_r$.


Wrap complexes

Generalizes and greatly simplifies the surface reconstruction algorithm *Wrap* (Edelsbrunner 1995, Geomagic)



Wrap complexes


Consider the Delaunay function f_D of X .



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)



Wrap complexes

Consider the Delaunay function f_D of X .

- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)




Wrap complexes

Consider the Delaunay function f_D of X .

- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$




Wrap complexes

Consider the Delaunay function f_D of X .

- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$




Wrap complexes

Consider the Delaunay function f_D of X .

- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of Delaunay intervals (with the partial order induced by the face relation)

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$



By construction: $\text{Del}_r \searrow \text{Wrap}_r$.

Wrapping up

- Čech and Delaunay complexes from Morse functions
- Explicit construction of simplicial collapses
- Simple definition and generalization of *Wrap* complexes