

Topological simplification problems

Ulrich Bauer

TUM

October 8, 2018


Mathematical Signal Processing and Data Analysis – University of Bremen

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a cell complex
is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .



A function $f : K \rightarrow \mathbb{R}$ on a cell complex is a *discrete Morse function* if


- sublevel sets are subcomplexes, and

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .



A function $f : K \rightarrow \mathbb{R}$ on a cell complex is a *discrete Morse function* if

- sublevel sets are subcomplexes, and



Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .



A function $f : K \rightarrow \mathbb{R}$ on a cell complex is a *discrete Morse function* if

- sublevel sets are subcomplexes, and




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .



A function $f : K \rightarrow \mathbb{R}$ on a cell complex is a *discrete Morse function* if

- sublevel sets are subcomplexes, and




Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .



A function $f : K \rightarrow \mathbb{R}$ on a cell complex is a *discrete Morse function* if

- sublevel sets are subcomplexes, and



Discrete Morse theory



Definition (Forman 1998)

A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .

A function $f : K \rightarrow \mathbb{R}$ on a cell complex is a *discrete Morse function* if

- sublevel sets are subcomplexes, and



Discrete Morse theory

Definition (Forman 1998)


A *discrete vector field* on a cell complex is a partition of the set of simplices into

- singleton sets $\{\phi\}$ (*critical cells*), and
- pairs $\{\sigma, \tau\}$, where σ is a facet of τ .



A function $f : K \rightarrow \mathbb{R}$ on a cell complex is a *discrete Morse function* if

- sublevel sets are subcomplexes, and
- level sets form a discrete vector field.



Fundamental theorem of discrete Morse theory


Let f be a discrete Morse function on a cell complex K .

Fundamental theorem of discrete Morse theory

Let f be a discrete Morse function on a cell complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f ,
then the sublevel set K_t collapses to K_s
(written as $K_t \searrow K_s$).




Fundamental theorem of discrete Morse theory

Let f be a discrete Morse function on a cell complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f ,
then the sublevel set K_t collapses to K_s
(written as $K_t \searrow K_s$).




Fundamental theorem of discrete Morse theory

Let f be a discrete Morse function on a cell complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f ,
then the sublevel set K_t collapses to K_s
(written as $K_t \searrow K_s$).




Fundamental theorem of discrete Morse theory

Let f be a discrete Morse function on a cell complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f ,
then the sublevel set K_t collapses to K_s
(written as $K_t \searrow K_s$).



Corollary

K is homotopy-equivalent to a cell complex M built from the critical cells of f .

Fundamental theorem of discrete Morse theory

Let f be a discrete Morse function on a cell complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f ,
then the sublevel set K_t collapses to K_s
(written as $K_t \searrow K_s$).



Corollary

K is homotopy-equivalent to a cell complex M built from the critical cells of f .

This homotopy equivalence is compatible with the filtration.

Corollary

K and M have isomorphic persistent homology (with regard to f).

Homology inference

Homology inference

Given: finite sample $P \subset \Omega$ of unknown shape $\Omega \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(\Omega)$.*

Homology inference

Given: finite sample $P \subset \Omega$ of unknown shape $\Omega \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(\Omega)$.*

Problem (Homological reconstruction)

Construct a shape X with $H_(X) \cong H_*(\Omega)$.*

Homology inference

Given: finite sample $P \subset \Omega$ of unknown shape $\Omega \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(\Omega)$.*

Problem (Homological reconstruction)

Construct a shape X with $H_(X) \cong H_*(\Omega)$.*

Idea:

- approximate the shape by a thickening $B_\delta(P)$ covering Ω

Homology inference

Given: finite sample $P \subset \Omega$ of unknown shape $\Omega \subset \mathbb{R}^d$

Problem (Homology inference)

Determine the homology $H_(\Omega)$.*

Problem (Homological reconstruction)

Construct a shape X with $H_(X) \cong H_*(\Omega)$.*

Idea:

- approximate the shape by a thickening $B_\delta(P)$ covering Ω

Requires strong assumptions:


Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.




Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let M be a submanifold of \mathbb{R}^d . Let $P \subset M$, $\delta > 0$ be such that

- $B_\delta(P)$ covers M , and
- $\delta < \sqrt{3/20} \text{reach}(M)$.

Then $H_*(M) \cong H_*(B_{2\delta}(P))$.





Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\Omega \subset \mathbb{R}^d$. Let $P \subset \Omega$, $\delta > 0$ be such that

- $B_\delta(P)$ covers Ω , and
- the inclusions $\Omega \hookrightarrow B_\delta(\Omega) \hookrightarrow B_{2\delta}(\Omega)$ preserve homology.

Then $H_*(\Omega) \cong \text{im } H_*(B_\delta(P) \hookrightarrow B_{2\delta}(P))$.



Homological realization

This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$



Homological realization

This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$




Homological realization

This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$



Homological realization

This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



$$L \hookrightarrow X? \hookrightarrow K$$

Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



Homological realization


This motivates the *homological realization problem*:

Problem


Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$

This is not always possible:



$$L \hookrightarrow X? \hookrightarrow K$$



$$L \hookrightarrow X \hookrightarrow K$$

Homological realization


This motivates the *homological realization problem*:

Problem

Given a simplicial pair $L \subseteq K$, find X with $L \subseteq X \subseteq K$ such that

$$H_*(X) = \text{im } H_*(L \hookrightarrow K).$$


This is not always possible:




Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)

The homological realization problem is NP-hard, even in \mathbb{R}^3 .

Simplification





Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

*Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$ minimizing $\dim H_*(G_{\leq t})$.*

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$ minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$ minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$ minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.
- For any X with $L \subseteq X \subseteq K$, there is g with $G_{\leq t} = X$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$ minimizing $\dim H_*(G_{\leq t})$.

Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.
- For any X with $L \subseteq X \subseteq K$, there is g with $G_{\leq t} = X$.

Thus we seek X with $L \subseteq X \subseteq K$ minimizing $\dim H_*(X)$.

Sublevel set simplification

Let $F_{\leq t} = f^{-1}(-\infty, t]$ denote the t -sublevel set of f .

Problem (Sublevel set simplification)

Given a function $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ and $t \in \mathbb{R}, \delta > 0$,
find a function g with $\|g - f\|_\infty \leq \delta$ minimizing $\dim H_*(G_{\leq t})$.


Let $K = F_{\leq t+\delta}$, $L = F_{\leq t-\delta}$.

- For any g , we have $L \subseteq G_{\leq t} \subseteq K$.
- For any X with $L \subseteq X \subseteq K$, there is g with $G_{\leq t} = X$.

Thus we seek X with $L \subseteq X \subseteq K$ minimizing $\dim H_*(X)$.

Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)

Sublevel set simplification in \mathbb{R}^3 is NP-hard.







Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

Given a function f and a real number $\delta \geq 0$, find a function f_δ with the minimal number of critical points subject to $\|f_\delta - f\|_\infty \leq \delta$.

Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

Given a function f and a real number $\delta \geq 0$, find a function f_δ with the minimal number of critical points subject to $\|f_\delta - f\|_\infty \leq \delta$.

Persistent homology:

- Relates homology of different sublevel set
- Identifies pairs of critical points (birth and death of homological feature) and quantifies their *persistence*

Topological simplification of functions

Consider the following problem:

Problem (Topological simplification)

Given a function f and a real number $\delta \geq 0$, find a function f_δ with the minimal number of critical points subject to $\|f_\delta - f\|_\infty \leq \delta$.

Persistent homology:

- Relates homology of different sublevel set
- Identifies pairs of critical points (birth and death of homological feature) and quantifies their *persistence*

(Discrete) Morse theory:

- Relates critical points to homology of sublevel sets
- Provides method for canceling pairs of critical points

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)


Let $\|f - g\|_\infty = \delta$. Then there exists a matching between the intervals of the persistence barcodes of f and g such that

Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\|f - g\|_\infty = \delta$. Then there exists a matching between the intervals of the persistence barcodes of f and g such that

- matched intervals have endpoints within distance $\leq \delta$, and




Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\|f - g\|_\infty = \delta$. Then there exists a matching between the intervals of the persistence barcodes of f and g such that

- matched intervals have endpoints within distance $\leq \delta$, and
- unmatched intervals have length $\leq 2\delta$.




Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let $\|f - g\|_\infty = \delta$. Then there exists a matching between the intervals of the persistence barcodes of f and g such that

- matched intervals have endpoints within distance $\leq \delta$, and
- unmatched intervals have length $\leq 2\delta$.




Corollary

Let f be a discrete Morse function and let $\delta > 0$.

Then for every function f_δ with $\|f_\delta - f\|_\infty \leq \delta$ we have:


$$\begin{aligned} &\# \text{ critical points of } f_\delta \\ &\geq \# \text{ critical points of } f \text{ with persistence} > 2\delta. \end{aligned}$$

Canceling critical points of a gradient field



from *Lectures on the h-cobordism theorem* (Milnor, 1965)

Canceling critical points of a gradient field



from *Lectures on the h-cobordism theorem* (Milnor, 1965)

Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006,
Attali et al. 2009]:

- remove all persistence pairs of f with persistence $\leq 2\delta$

Side-effects of elimination


Idea for simplifying critical points [Edelsbrunner et al. 2006,
Attali et al. 2009]:

- remove all persistence pairs of f with persistence $\leq 2\delta$
- leave all other persistence pairs unmodified

Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:


- remove all persistence pairs of f with persistence $\leq 2\delta$
- leave all other persistence pairs unmodified



Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:


- remove all persistence pairs of f with persistence $\leq 2\delta$
- leave all other persistence pairs unmodified



Side-effects of elimination

Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:


- remove all persistence pairs of f with persistence $\leq 2\delta$
- leave all other persistence pairs unmodified



Side-effects of elimination


Idea for simplifying critical points [Edelsbrunner et al. 2006, Attali et al. 2009]:

- remove all persistence pairs of f with persistence $\leq 2\delta$
- leave all other persistence pairs unmodified




For an optimal solution, we must allow the critical values to change!


Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function




Cancelling critical points of a function



Cancelling critical points of a function



Canceling critical points of a function



Combining persistence and Morse theory

Theorem (B., Lange, Wardetzky, 2011)

Let f be a discrete Morse function on a surface (with distinct critical values) with gradient field V . Any persistence pair (σ, τ) can be canceled in V after all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with

$$f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$$

have been canceled.

Combining persistence and Morse theory

Theorem (B., Lange, Wardetzky, 2011)

Let f be a discrete Morse function on a surface (with distinct critical values) with gradient field V . Any persistence pair (σ, τ) can be canceled in V after all persistence pairs $(\tilde{\sigma}, \tilde{\tau})$ with

$$f(\sigma) < f(\tilde{\sigma}) < f(\tilde{\tau}) < f(\tau)$$

have been canceled.

Corollary

The function f_δ constructed from f by canceling all persistence pairs with persistence $\leq 2\delta$ achieves the lower bound on the number of critical points subject to $\|f_\delta - f\|_\infty \leq \delta$.

Combination with energy methods

Recall: simplified vector field V_δ imposes inequalities on simplified function consistent with V_δ

Combination with energy methods

Recall: simplified vector field V_δ imposes inequalities on simplified function consistent with V_δ

$\|f_\delta - f\|_\infty \leq \delta$: another set of linear inequalities

Combination with energy methods

Recall: simplified vector field V_δ imposes inequalities on simplified function consistent with V_δ

$\|f_\delta - f\|_\infty \leq \delta$: another set of linear inequalities

- defines convex polytope of solutions:
any function g consistent with V_δ and $\|g - f\|_\infty \leq \delta$ is a solution

Combination with energy methods


Recall: simplified vector field V_δ imposes inequalities on simplified function consistent with V_δ

$\|f_\delta - f\|_\infty \leq \delta$: another set of linear inequalities

- defines convex polytope of solutions:
any function g consistent with V_δ and $\|g - f\|_\infty \leq \delta$ is a solution
- find the “best” solution using your favorite energy functional


Cancelation conflicts

The result does not generalize beyond surfaces!



Cancelation conflicts

The result does not generalize beyond surfaces!



Room for discussions...