

# Connect the dots

## From data through complexes to persistent homology

Ulrich Bauer

Technical University of Munich (TUM)

September 28, 2023

MathSEE Symposium *Mathematics in Sciences, Engineering and Economics*

Funded by



Deutsche  
Forschungsgemeinschaft  
German Research Foundation



SFB  
TRR  
109

Discretization  
In Geometry  
and Dynamics

Technical  
University  
of Munich



MCSL  
Munich Center for Machine Learning



$\equiv$



?




$\equiv$




?

# Geometry and topology of biomolecules




Gramicidin (an antibiotic functioning as an ion channel)


# Geometry and topology of biomolecules




Gramicidin (an antibiotic functioning as an ion channel)





























# What is homology?




# What is homology?




# What is homology?




# What is homology?




# What is homology?



## What is homology?




# What is homology?




# Geometric complexes


## Čech complexes




## Čech complexes




## Čech complexes




## Čech complexes




## Čech complexes




## Čech complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes



## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes




## Delaunay complexes



# Discrete Morse theory




# Discrete Morse theory



# Discrete Morse theory

## Theorem (Forman 1998)


A simplicial complex with a discrete Morse function  $f$  is homotopy equivalent to a CW complex built from the critical simplices of  $f$ .



# Discrete Morse theory

## Theorem (Forman 1998)

A simplicial complex with a discrete Morse function  $f$  is homotopy equivalent to a CW complex built from the critical simplices of  $f$ .




Discrete Morse functions – and their gradients – encode collapses of sublevel sets:




# Discrete Morse theory

## Theorem (Forman 1998)

A simplicial complex with a discrete Morse function  $f$  is homotopy equivalent to a CW complex built from the critical simplices of  $f$ .




Discrete Morse functions – and their gradients – encode collapses of sublevel sets:



# Generalizing discrete Morse theory

Generalized gradients are a partition into intervals in the face poset (instead of just facet pairs):



# Morse theory for Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech complexes and the Delaunay complexes (alpha shapes) are sublevel sets of (generalized) discrete Morse functions.*

# Morse theory for Čech and Delaunay complexes


## Proposition (B, Edelsbrunner 2014)

*The Čech complexes and the Delaunay complexes (alpha shapes) are sublevel sets of (generalized) discrete Morse functions.*

## Theorem (B., Edelsbrunner 2017)


*Čech, Delaunay, and Wrap complexes (at any scale  $r$ ) are related by collapses encoded by a single discrete gradient field:*

$$\text{Cech}_r X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$




## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .



## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .




**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**

$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).

## Wrap complexes

Consider the gradient  $V$  of the Delaunay radius function  $\text{Del}(X) \rightarrow \mathbb{R}$ .



**Definition (Edelsbrunner 1995; B, Edelsbrunner 2017)**


$\text{Wrap}_r(X)$  is the *descending complex* of  $V$  on  $\text{Del}_r X$ : the smallest subcomplex of  $\text{Del}_r X$  that

- contains all critical simplices
- is compatible with  $V$  (a union of intervals of  $V$ ).


## Delaunay and Wrap complexes




## Delaunay and Wrap complexes




## Wrap complexes and lexicographically minimal cycles




## Wrap complexes and lexicographically minimal cycles




## Wrap complexes and lexicographically minimal cycles




## Wrap complexes and lexicographically minimal cycles




## Wrap complexes and lexicographically minimal cycles




## Wrap complexes and lexicographically minimal cycles






## Wrap complexes and lexicographically minimal cycles



## Wrap complexes and lexicographically minimal cycles





## Point cloud reconstruction with minimal cycles



# Minimal cycles within Wrap complexes

## Theorem (B, Roll 2022)

Let  $X \subset \mathbb{R}$  be a finite subset in general position and let  $r \in \mathbb{R}$ . Then the lexicographically minimal cycles of  $\text{Del}_r(X)$  are supported on  $\text{Wrap}_r(X)$ .



# Homology inference

## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .

## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



# Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

## Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \operatorname{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \operatorname{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



# Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

## Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .


- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .



## Homology reconstruction by thickening

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .

- Approach: approximate the shape by a thickening of  $P$  (union of balls) that covers  $X$ .

### Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .

## Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

## Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the *common features*.

## Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .


- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_*: H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).



## Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_*: H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).

**Proof.**

$$\begin{array}{ccccccc} X & & X_\delta & & X_{2\delta} \\ & \searrow & \nearrow & \searrow & \nearrow \\ & P_\delta & & P_{2\delta} & \end{array}$$

## Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_*: H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).

**Proof.**

$$\begin{array}{ccccccc} X & & & X_\delta & & & X_{2\delta} \\ & \searrow & & \nearrow & & \searrow & \nearrow \\ & & P_\delta & & & P_{2\delta} & \end{array}$$

## Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_* : H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).

**Proof.**

$$\begin{array}{ccc} H_*(X) & & H_*(X_\delta) & & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & H_*(P_\delta) & & H_*(P_{2\delta}) & \end{array}$$

# Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_* : H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).

**Proof.**

$$\begin{array}{ccccc} H_*(X) & \xrightarrow{\quad} & H_*(X_\delta) & \xleftarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_* : H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).

**Proof.**

$$\begin{array}{ccccc} H_*(X) & \xrightarrow{\quad} & H_*(X_\delta) & \xleftarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_* : H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).

**Proof.**

$$\begin{array}{ccccc} H_*(X) & \xrightarrow{\quad} & H_*(X_\delta) & \xleftarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & H_*(P_\delta) & & H_*(P_{2\delta}) & \end{array}$$

# Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_*: H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).

**Proof.**

$$\begin{array}{ccccc} H_*(X) & \xrightarrow{\quad} & H_*(X_\delta) & \xrightarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \downarrow \cong & \swarrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \\ & & \searrow & \downarrow & \swarrow \\ & & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & \end{array}$$

# Homology inference using persistence

Given a finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$ , determine the homology  $H_*(X)$ .

- Approach: consider of  $P$  at two different scales, keeping only the common features.

**Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)**

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the inclusions  $X \hookrightarrow X_\delta \hookrightarrow X_{2\delta}$  of thickenings induce isomorphisms in homology.

Then  $H_*(X) \cong \text{im}(i_*: H_*(P_\delta) \rightarrow H_*(P_{2\delta}))$  (the map induced by inclusion  $P_\delta \hookrightarrow P_{2\delta}$ ).


**Proof.**

$$\begin{array}{ccccc} H_*(X) & \xhookrightarrow{\quad \cong \quad} & H_*(X_\delta) & \longrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \downarrow \cong & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \\ & & \searrow & \downarrow & \nearrow \\ & & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & \end{array}$$


□

# Persistent homology


# What is persistent homology?




# What is persistent homology?




# What is persistent homology?




# What is persistent homology?




# What is persistent homology?



# What is persistent homology?



# What is persistent homology?



# What is persistent homology?

## What is persistent homology?

Persistent homology is the homology of a filtration.

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$  of topological spaces, indexed over the poset of real numbers  $\mathbf{R} := (\mathbb{R}, \leq)$

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space  $K_t$  for each  $t \in \mathbb{R}$
- an inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$  of topological spaces, indexed over the poset of real numbers  $\mathbf{R} := (\mathbb{R}, \leq)$

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space  $K_t$  for each  $t \in \mathbb{R}$
- an inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$  of topological spaces, indexed over the poset of real numbers  $\mathbf{R} := (\mathbb{R}, \leq)$

$$\dots \rightarrow K_s \hookrightarrow K_t \rightarrow \dots$$

- a topological space  $K_t$  for each  $t \in \mathbb{R}$
- an inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- Persistent homology is a diagram  $M = H_* \circ K : \mathbf{R} \rightarrow \mathbf{Vect}$  (*persistence module*):

$$\dots \rightarrow M_s \longrightarrow M_t \rightarrow \dots$$





# Barcodes: the structure of persistence modules

## Theorem (Crawley-Boevey 2015)

*Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules*

$$\rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow$$

*(in an essentially unique way).*

# Barcodes: the structure of persistence modules


## Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$\rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow$$

(in an essentially unique way).

- The supporting intervals form the *persistence barcode*.



# Stability


# Persistence and stability: the big picture

Data


point cloud

$$P \subset \mathbb{R}^d$$


# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture



# Persistence and stability: the big picture



# Persistence and stability: the big picture




## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .


Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:




# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:




Applying homology, the persistence modules  $H_*(F), H_*(G) : \mathbf{R} \rightarrow \mathbf{Vect}$  are  $\delta$ -interleaved:



# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2016; B, Lesnick 2015)

*If two persistence modules are  $\delta$ -interleaved,  
then their barcodes admit a  $\delta$ -matching:*




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2016; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then their barcodes admit a  $\delta$ -matching:

- matched intervals have endpoints within distance  $\leq \delta$ , and




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2016; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then their barcodes admit a  $\delta$ -matching:

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .



## Structure of persistence sub-/quotient modules

Proposition (B, Lesnick 2015)

Let  $M \twoheadrightarrow N$  be an epimorphism of persistence modules.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that

if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial (compatible with composition).



# Structure of persistence sub-/quotient modules

Proposition (B, Lesnick 2015)

Let  $M \twoheadrightarrow N$  be an epimorphism of persistence modules.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that

if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial (compatible with composition).

Dually, for a monomorphism  $N \hookrightarrow O$  there is an injection  $B(N) \hookrightarrow B(O)$ .



## Induced matchings

Any morphism of pfd persistence modules  $f : M \rightarrow N$  factors through its image as an epimorphism followed by a monomorphism:

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

This gives an *induced matching* between their barcodes:

## Induced matchings

Any morphism of pfd persistence modules  $f : M \rightarrow N$  factors through its image as an epimorphism followed by a monomorphism:

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

This gives an *induced matching* between their barcodes:

- compose the injections  $B(M) \hookleftarrow B(\text{im } f) \hookrightarrow B(N)$  from before to a matching  $B(M) \leftrightarrow B(N)$



## Induced matchings

Any morphism of pfd persistence modules  $f : M \rightarrow N$  factors through its image as an epimorphism followed by a monomorphism:

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

This gives an *induced matching* between their barcodes:

- compose the injections  $B(M) \hookleftarrow B(\text{im } f) \hookrightarrow B(N)$  from before to a matching  $B(M) \leftrightarrow B(N)$




If  $f$  is a  $\delta$ -interleaving morphism, then this induced matching is a  $\delta$ -matching.

# Persistence of Vietoris–Rips complexes

## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite, } \text{diam } S \leq t\}.$$



## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



## Vietoris–Rips complexes

For a metric space  $X$ , the *Vietoris–Rips complex* at  $t > 0$  is the simplicial complex



$$\text{Rips}_t(X) = \{S \subseteq X \mid S \neq \emptyset \text{ finite}, \text{diam } S \leq t\}.$$



# Ripser: software for computing Vietoris–Rips persistence barcodes

Open source software ([ripser.org](http://ripser.org))

- significantly faster / more memory efficient than previous codes
- de facto standard for topological data analysis applications
- most popular GitHub project on persistent homology



Ripser users from 935 different locations

Computational improvements based on

- *implicit matrix representations*
- *apparent pairs*, connecting persistence to discrete Morse theory

## Apparent pairs

Ripser uses the following pairing of simplices (for the filtration, we break ties lexicographically):

### Definition (B 2016, 2021)

In a simplexwise filtration ( $K_i = \{\sigma_1, \dots, \sigma_i\}$ )<sub>i</sub>, two simplices  $(\sigma_i, \sigma_j)$  form an *apparent pair* if

- $\sigma_i$  is the latest proper face of  $\sigma_j$ , and
- $\sigma_j$  is the earliest proper coface of  $\sigma_i$ .

## Apparent pairs

Ripser uses the following pairing of simplices (for the filtration, we break ties lexicographically):

### Definition (B 2016, 2021)

In a simplexwise filtration ( $K_i = \{\sigma_1, \dots, \sigma_i\}$ )<sub>i</sub>, two simplices  $(\sigma_i, \sigma_j)$  form an *apparent pair* if

- $\sigma_i$  is the latest proper face of  $\sigma_j$ , and
- $\sigma_j$  is the earliest proper coface of  $\sigma_i$ .

Special cases and equivalent definitions have been (re)discovered independently multiple times

- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

## Apparent pairs

Ripser uses the following pairing of simplices (for the filtration, we break ties lexicographically):

### Definition (B 2016, 2021)

In a simplexwise filtration ( $K_i = \{\sigma_1, \dots, \sigma_i\}$ )<sub>i</sub>, two simplices  $(\sigma_i, \sigma_j)$  form an *apparent pair* if

- $\sigma_i$  is the latest proper face of  $\sigma_j$ , and
- $\sigma_j$  is the earliest proper coface of  $\sigma_i$ .

Special cases and equivalent definitions have been (re)discovered independently multiple times

- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

### Proposition (B 2021)

The apparent pairs are both

- gradient pairs (in the sense of discrete Morse theory) and
- persistence pairs (creating/destroying a feature in homology).

## Apparent pairs

Ripser uses the following pairing of simplices (for the filtration, we break ties lexicographically):

### Definition (B 2016, 2021)

In a simplexwise filtration ( $K_i = \{\sigma_1, \dots, \sigma_i\}$ )<sub>i</sub>, two simplices  $(\sigma_i, \sigma_j)$  form an *apparent pair* if

- $\sigma_i$  is the latest proper face of  $\sigma_j$ , and
- $\sigma_j$  is the earliest proper coface of  $\sigma_i$ .

Special cases and equivalent definitions have been (re)discovered independently multiple times

- Kahle 2011, Robbins 2015, Henselmann-Petrusek 2017, Lampret 2020, Feichtner-Kozlov 2020, ...

### Proposition (B 2021)


The apparent pairs are both

- gradient pairs (in the sense of discrete Morse theory) and
- persistence pairs (creating/destroying a feature in homology).

### Proposition (B, Roll 2022)


Any generalized discrete Morse function is refined by apparent pairs.

# Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadan, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

# Topology of viral evolution



Joint work with: A. Ott, M. Bleher, L. Hahn (Heidelberg), R. Rabadian, J. Patiño-Galindo (Columbia), M. Carrière (INRIA)

Observation: Ripser runs unusually fast on genetic distance data


- SARS-CoV2 RNA sequences (spike protein)
- 25556 data points ( $2.8 \times 10^{12}$  simplices in 2-skeleton)
- 120 s computation time (with data points ordered appropriately)

# Gromov-hyperbolicity

## Definition (Gromov 1988)

A metric space  $X$  is  $\delta$ -hyperbolic (for  $\delta \geq 0$ ) if for all  $w, x, y, z \in X$  we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$




# Gromov-hyperbolicity

## Definition (Gromov 1988)

A metric space  $X$  is  $\delta$ -hyperbolic (for  $\delta \geq 0$ ) if for all  $w, x, y, z \in X$  we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$



- The hyperbolic plane is  $(\ln 2)$ -hyperbolic.




# Gromov-hyperbolicity

## Definition (Gromov 1988)

A metric space  $X$  is  $\delta$ -hyperbolic (for  $\delta \geq 0$ ) if for all  $w, x, y, z \in X$  we have

$$d(w, x) + d(y, z) \leq \max\{d(w, y) + d(x, z), d(w, z) + d(x, y)\} + 2\delta.$$



- The hyperbolic plane is  $(\ln 2)$ -hyperbolic.



- The 0-hyperbolic spaces are precisely the metric trees and their subspaces.



## Rips Contractibility

Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

# Rips Contractibility

## Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

- What about non-geodesic spaces? In particular, finite metric spaces?
- What about collapsibility?
- What about the filtration?
- Connection to computation of persistent homology?

# Rips Contractibility

## Theorem (Rips; Gromov 1988)

*Let  $X$  be a  $\delta$ -hyperbolic geodesic metric space. Then  $\text{Rips}_t(X)$  is contractible for all  $t \geq 4\delta$ .*

- What about non-geodesic spaces? In particular, finite metric spaces?
- What about collapsibility?
- What about the filtration?
- Connection to computation of persistent homology?


## Theorem (B, Roll 2022)

*Let  $X$  be a finite  $\delta$ -hyperbolic space. Then there is a single discrete gradient encoding the collapses*


$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \searrow \{\ast\}$$

*for all  $u > t \geq 4\delta + 2\nu$ , where  $\nu$  is the geodesic defect of  $X$ .*


## Geodesic defect




## Geodesic defect



## Geodesic defect



## Geodesic defect



Definition (Bonk, Schramm 2000)

A metric space  $X$  is  $v$ -geodesic if for all points  $x, y \in X$  and all  $r, s \geq 0$  with  $r + s = d(x, y)$  we have

$$B_{r+v}(x) \cap B_{s+v}(y) \neq \emptyset.$$

The infimum of all such  $v$  is the *geodesic defect* of  $X$ .

## The diameter function of generic trees

Proposition (B, Roll 2022)

*Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.*

# The diameter function of generic trees

## Proposition (B, Roll 2022)

Consider a finite weighted tree  $(V, E)$  with a generic path length metric (distinct pairwise distances). Then the diameter function  $\text{diam}: \Delta(V) \rightarrow \mathbb{R}$  is a generalized discrete Morse function.

- The apparent pairs refine this gradient.

## Theorem (B, Roll 2022)

The apparent pairs of the diameter function for a generic tree metric space  $X$  induces the collapses

$$\text{Rips}_t(X) \searrow T_t \quad \text{for all } t \in \mathbb{R}, \text{ with}$$

$$\text{Rips}_t(X) \searrow T \searrow \{\ast\} \quad \text{for } t \geq \max l(E), \text{ and}$$

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \quad \text{whenever } l(e) \notin (t, u] \text{ for all } e \in E.$$

In particular, the persistent homology is trivial in degrees  $> 0$ .

## Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.

# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.


## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case:*

$$\text{Rips}_t(X) \searrow T_t \quad \text{for all } t \in \mathbb{R}, \text{ with}$$

$$\text{Rips}_t(X) \searrow T \searrow \{\ast\} \quad \text{for } t \geq \max l(E), \text{ and}$$

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \quad \text{whenever } l(e) \notin (t, u] \text{ for all } e \in E.$$



# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.


## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case:*

$$\text{Rips}_t(X) \searrow T_t \quad \text{for all } t \in \mathbb{R}, \text{ with}$$

$$\text{Rips}_t(X) \searrow T \searrow \{\ast\} \quad \text{for } t \geq \max l(E), \text{ and}$$

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \quad \text{whenever } l(e) \notin (t, u] \text{ for all } e \in E.$$



# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.


## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case:*

$$\text{Rips}_t(X) \searrow T_t \quad \text{for all } t \in \mathbb{R}, \text{ with}$$

$$\text{Rips}_t(X) \searrow T \searrow \{\ast\} \quad \text{for } t \geq \max l(E), \text{ and}$$

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \quad \text{whenever } l(e) \notin (t, u] \text{ for all } e \in E.$$



# Tree metrics beyond the generic case

Why is Ripser particularly fast on genetic distances (tree-like)?

- Consider a weighted finite tree  $T = (V, E)$ , viewed as a metric space  $X$ .
- Choose an arbitrary root and extend the rooted tree partial order to a total order.


## Theorem (B, Roll 2022)

*The apparent pairs gradient for this total order induces the same collapses as in the generic case:*

$$\text{Rips}_t(X) \searrow T_t \quad \text{for all } t \in \mathbb{R}, \text{ with}$$

$$\text{Rips}_t(X) \searrow T \searrow \{\ast\} \quad \text{for } t \geq \max l(E), \text{ and}$$

$$\text{Rips}_u(X) \searrow \text{Rips}_t(X) \quad \text{whenever } l(e) \notin (t, u] \text{ for all } e \in E.$$



# Origins

When was persistent homology discovered first?

# When was persistent homology discovered first?

ANNALS OF MATHEMATICS  
Vol. 41, No. 2, April, 1940

## RANK AND SPAN IN FUNCTIONAL TOPOLOGY

BY MARSTON MORSE

(Received August 9, 1939)

### 1. Introduction.

The analysis of functions  $F$  on metric spaces  $M$  of the type which appear in variational theories is made difficult by the fact that the critical limits, such as absolute minima, relative minima, minimax values etc., are in general infinite in number. These limits are associated with relative  $k$ -cycles of various dimensions and are classified as 0-limits, 1-limits etc. The number of  $k$ -limits suitably counted is called the  $k^{\text{th}}$  type number  $m_k$  of  $F$ . The theory seeks to establish relations between the numbers  $m_k$  and the connectivities  $p_k$  of  $M$ . The numbers  $p_k$  are finite in the most important applications. It is otherwise with the numbers  $m_k$ .

The theory has been able to proceed provided one of the following hypotheses is satisfied. The critical limits cluster at most at  $+\infty$ ; the critical points are

# When was persistent homology discovered first?

Web Images More... Sign in

Google Scholar 9 results (0.02 sec) My Citations ▾

All citations Rank and span in functional topology

Articles  Search within citing articles

Case law Exact homomorphism sequences in homology theory ed.ac.uk [PDF]

JL Kelley, E Pitcher - Annals of Mathematics, 1947 - JSTOR

The developments of this paper stem from the attempts of one of the authors to deduce relations between homology groups of a complex and homology groups of a complex which is its image under a simplicial map. Certain relations were deduced (see [EP 1] and [EP 2] ...)

Any time Since 2016 Cited by 46 Related articles All 3 versions Cite Save More

Since 2015 Since 2012 Custom range... Marston Morse and his mathematical works ams.org [PDF]

R Bott - Bulletin of the American Mathematical Society, 1980 - ams.org

American Mathematical Society. Thus Morse grew to maturity just at the time when the subject of Analysis Situs was being shaped by such masters as Poincaré, Veblen, LEJ Brouwer, GD Birkhoff, Lefschetz and Alexander, and it was Morse's genius and destiny to ...

Sort by relevance Sort by date Cited by 24 Related articles All 4 versions Cite Save More

Unstable minimal surfaces of higher topological structure

include citations M Morse, CB Tompkins - Duke Math. J., 1941 - projecteuclid.org

1. Introduction. We are concerned with extending the calculus of variations in the large to multiple integrals. The problem of the existence of minimal surfaces of unstable type contains many of the typical difficulties, especially those of a topological nature. Having studied this ...

Cited by 19 Related articles All 2 versions Cite Save


[PDF] Persistence in discrete Morse theory psu.edu [PDF]

U Bauer - 2011 - Citeseer

# Motivation and application: minimal surfaces

## Problem (Plateau's problem)

*Find an immersed disk of least area spanned by a given closed Jordan curve.*




(from Dierkes et al.: *Minimal Surfaces*, 2010)

## Motivation and application: minimal surfaces

### Problem (Plateau's problem)

*Find an immersed disk of least area spanned by a given closed Jordan curve.*



(from Dierkes et al.: *Minimal Surfaces*, 2010)


Solution by Douglas (1930):

- identifies minimal surfaces with critical points of the *Douglas functional*

## Existence of unstable minimal surfaces

Theorem (Morse, Tompkins 1939; Shiffman 1939)

*Assume that a given curve bounds two separate stable minimal surfaces.*




## Existence of unstable minimal surfaces

Theorem (Morse, Tompkins 1939; Shiffman 1939)

Assume that a given curve bounds two separate stable minimal surfaces.

Then that curve also bounds an unstable minimal surface (a critical point that is not a local minimum).



## Q-tame persistence modules

### Definition (Chazal et al. 2009)

A persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  is *q-tame* if all structure maps  $M_s \rightarrow M_t$  ( $s < t$ ) have finite rank.

## Q-tame persistence modules

### Definition (Chazal et al. 2009)

A persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  is *q-tame* if all structure maps  $M_s \rightarrow M_t$  ( $s < t$ ) have finite rank.

- A q-tame persistence module still admits a barcode (disregarding the endpoints).

## Q-tame persistence modules

### Definition (Chazal et al. 2009)

A persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  is *q-tame* if all structure maps  $M_s \rightarrow M_t$  ( $s < t$ ) have finite rank.

- A q-tame persistence module still admits a barcode (disregarding the endpoints).

Morse's main technical result, in modern language:

- The minimal surface functional has q-tame persistent homology.

## Q-tameness from local connectivity


### Theorem (Morse, 1937)

*If the sublevel set filtration off:  $X \rightarrow \mathbb{R}$  is compact and weakly locally connected, then it has  $q$ -tame persistent Vietoris homology.*

# Q-tameness from local connectivity

Theorem (Morse, 1937; incorrect)


If the sublevel set filtration off:  $X \rightarrow \mathbb{R}$  is compact and weakly locally connected, then it has  $q$ -tame persistent Vietoris homology.



## Q-tameness from local connectivity

Theorem (Morse, 1937; incorrect)

*If the sublevel set filtration off:  $X \rightarrow \mathbb{R}$  is compact and weakly locally connected, then it has  $q$ -tame persistent Vietoris homology.*



Theorem (B, Medina-Mardones, Schmahl 2021)

*If the sublevel set filtration off:  $X \rightarrow \mathbb{R}$  is compact and homologically locally connected, then it has  $q$ -tame persistent homology.*

# Simplification

# Topological simplification of functions

Consider the following problem:


## Problem (Topological simplification)

*Given a function  $f$  and a real number  $\delta \geq 0$ , find a function  $f_\delta$  with the minimal number of critical points subject to  $\|f_\delta - f\|_\infty \leq \delta$ .*

# Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points




(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

# Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points



(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

For a Morse function:

- critical points correspond to endpoints of barcode intervals

# Combining persistence and Morse theory

By stability of persistence barcodes:

## Proposition

*The critical points of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

# Combining persistence and Morse theory

By stability of persistence barcodes:

## Proposition

*The critical points of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

## Theorem (B, Lange, Wardetzky, 2011)

*Let  $f$  be a function on a surface and let  $\delta > 0$ .*

*Canceling all pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$*

- *satisfying  $\|f_\delta - f\|_\infty \leq \delta$  and*
- *achieving the lower bound on the number of critical points.*

# Combining persistence and Morse theory

By stability of persistence barcodes:

## Proposition

*The critical points of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*


## Theorem (B, Lange, Wardetzky, 2011)


*Let  $f$  be a function on a surface and let  $\delta > 0$ .*


*Canceling all pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$*


- *satisfying  $\|f_\delta - f\|_\infty \leq \delta$  and*
- *achieving the lower bound on the number of critical points.*


Does not generalize to higher-dimensional manifolds!














Thanks for your attention!