

# The structure of persistence

(An introspection)

Ulrich Bauer

TUM

Apr 24, 2019


TopApp workshop 2019, IST Austria


# Part 1:


# 1-parameter persistence


(Barcodes & persistence diagrams)


Joint work with Michael Lesnick (Albany)


















# What is persistent homology?



# What is persistent homology?



# Interval decompositions and persistence modules

## Theorem (Crawley-Boewey 2015)

*Any pointwise finite-dimensional (pfd) persistence module (a diagram  $M : \mathbb{R} \rightarrow \mathbf{vect}$ ) has an essentially unique decomposition as a direct sum of indecomposable interval modules, isomorphic to*

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \underbrace{\mathbb{K} \rightarrow \cdots \rightarrow \mathbb{K}}_{\text{supported by an interval } I \subseteq \mathbb{R}} \rightarrow 0 \rightarrow \cdots$$

- ▶ The corresponding collection (multiset) of intervals is the *persistence barcode* of  $M$ .

# Interval decompositions and persistence modules

## Theorem (Crawley-Boewey 2015)

Any pointwise finite-dimensional (pfd) persistence module (a diagram  $M : \mathbb{R} \rightarrow \mathbf{vect}$ ) has an essentially unique decomposition as a direct sum of indecomposable interval modules, isomorphic to

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \underbrace{\mathbb{K} \rightarrow \cdots \rightarrow \mathbb{K}}_{\text{supported by an interval } I \subseteq \mathbb{R}} \rightarrow 0 \rightarrow \cdots$$

- ▶ The corresponding collection (multiset) of intervals is the *persistence barcode* of  $M$ .
- ▶ The points in the *persistence diagram* are the endpoints of the intervals in the barcode.

# Interval decompositions and persistence modules

## Theorem (Crawley-Boewey 2015)

Any pointwise finite-dimensional (pfd) persistence module (a diagram  $M : \mathbb{R} \rightarrow \mathbf{vect}$ ) has an essentially unique decomposition as a direct sum of indecomposable interval modules, isomorphic to

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \underbrace{\mathbb{K} \rightarrow \cdots \rightarrow \mathbb{K}}_{\text{supported by an interval } I \subseteq \mathbb{R}} \rightarrow 0 \rightarrow \cdots$$

- ▶ The corresponding collection (multiset) of intervals is the *persistence barcode* of  $M$ .
- ▶ The points in the *persistence diagram* are the endpoints of the intervals in the barcode.
- ▶ This is not a diagram in the sense of category theory (functor)!

# Persistence and stability: the big picture

point cloud

$$P \subset \mathbb{R}^d$$

Hausdorff distance

# Persistence and stability: the big picture

point cloud

$$P \subset \mathbb{R}^d$$

Hausdorff distance




distance

function

$$f : \mathbb{R}^d \rightarrow \mathbb{R}$$


$L_\infty$  distance

# Persistence and stability: the big picture






# Persistence and stability: the big picture



# Persistence and stability: the big picture




# The category of matchings


Consider the category **Mch** (a subcategory of the category **Rel** of sets and relations) with

- ▶ objects: sets,
- ▶ morphisms: matchings (partial bijections).

Composition:



(Co)kernel/(co)image:




# The category of matchings


Consider the category **Mch** (a subcategory of the category **Rel** of sets and relations) with

- ▶ objects: sets,
- ▶ morphisms: matchings (partial bijections).

Composition:



(Co)kernel/(co)image:



**Mch** is a *Puppe-exact* (*p-exact*) category:


- ▶ it has a zero object ( $\emptyset$ )
- ▶ it has all (co)kernels
- ▶ every mono (epi) is (co)kernel
- ▶ every morphism  $f : A \rightarrow B$  has an epi-mono factorization  $A \twoheadrightarrow \operatorname{im} f \hookrightarrow B$

# The category of matchings


Consider the category **Mch** (a subcategory of the category **Rel** of sets and relations) with

- ▶ objects: sets,
- ▶ morphisms: matchings (partial bijections).

Composition:



(Co)kernel/(co)image:



**Mch** is a *Puppe-exact* (*p-exact*) category:


- ▶ it has a zero object ( $\emptyset$ )
- ▶ it has all (co)kernels
- ▶ every mono (epi) is (co)kernel
- ▶ every morphism  $f : A \rightarrow B$  has an epi-mono factorization  $A \twoheadrightarrow \text{im } f \hookrightarrow B$

but not additive:


- ▶ it does not have all (co)products

## From barcodes to matching diagrams (and back)

- ▶ A barcode (collection of intervals) can be read as a diagram  $\mathbb{R} \rightarrow \mathbf{Mch}$ :





$t \mapsto \{\text{intervals in barcode containing } t\}$



$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$

# From barcodes to matching diagrams (and back)


- ▶ A barcode (collection of intervals) can be read as a diagram  $\mathbb{R} \rightarrow \mathbf{Mch}$ :



$t \mapsto \{\text{intervals in barcode containing } t\}$



$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$

- ▶ A matching diagram defines a barcode:



# From barcodes to matching diagrams (and back)


- ▶ A barcode (collection of intervals) can be read as a diagram  $\mathbb{R} \rightarrow \mathbf{Mch}$ :



$t \mapsto \{\text{intervals in barcode containing } t\}$

$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$

- ▶ A matching diagram defines a barcode:





- ▶ intervals formed by equivalence classes of matched elements




# From barcodes to matching diagrams (and back)

- ▶ A barcode (collection of intervals) can be read as a diagram  $\mathbb{R} \rightarrow \mathbf{Mch}$ :



$t \mapsto \{\text{intervals in barcode containing } t\}$        $(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$

- ▶ A matching diagram defines a barcode:



- ▶ intervals formed by equivalence classes of matched elements

Turn this into an equivalence of categories  $\mathbf{Barc} \simeq \mathbf{Mch}^{\mathbb{R}}$

# A category of barcodes

## Proposition

The functor category  $\mathbf{Mch}^{\mathbb{R}}$  is equivalent to  $\mathbf{Barc}$ , the category with

- ▶ *objects: barcodes (as a disjoint union of intervals),*
- ▶ *morphisms: overlap matchings of barcodes  $U \rightarrowtail V$ :*

# A category of barcodes

## Proposition

The functor category  $\mathbf{Mch}^{\mathbb{R}}$  is equivalent to  $\mathbf{Barc}$ , the category with

- ▶ objects: barcodes (as a disjoint union of intervals),
- ▶ morphisms: overlap matchings of barcodes  $U \rightarrowtail V$ :  
if  $I \in U$  is matched to  $J \in V$ , then  $I$  overlaps  $J$  above:
  - ▶  $I$  bounds  $J$  above (every  $s \in J$  is bounded above by some  $t \in I$ ),
  - ▶  $J$  bounds  $I$  below,
  - ▶  $I \cap J \neq \emptyset$ .



# A category of barcodes

## Proposition

The functor category  $\mathbf{Mch}^{\mathbb{R}}$  is equivalent to  $\mathbf{Barc}$ , the category with

- ▶ *objects*: barcodes (as a disjoint union of intervals),
- ▶ *morphisms*: overlap matchings of barcodes  $U \rightarrowtail V$ :  
if  $I \in U$  is matched to  $J \in V$ , then  $I$  overlaps  $J$  above:
  - ▶  $I$  bounds  $J$  above (every  $s \in J$  is bounded above by some  $t \in I$ ),
  - ▶  $J$  bounds  $I$  below,
  - ▶  $I \cap J \neq \emptyset$ .
- ▶ *composition of overlap matchings*:  $\tau \bullet \sigma = \{(I, K) \in \tau \circ \sigma \mid I \text{ overlaps } K \text{ above}\}$   
(where  $\tau \circ \sigma$  is the standard composition of matchings)



$(I, K) \in \tau \bullet \sigma$  (overlap)



$(I, K) \notin \tau \bullet \sigma$  (no overlap)


## Bottleneck distance as an interleaving distance

- ▶  $\delta$ -matching between barcodes  $U, V$ :
  - ▶ matched intervals have  $\delta$ -close endpoints
  - ▶ unmatched intervals are  $2\delta$ -trivial (shorter than  $2\delta$ )

# Bottleneck distance as an interleaving distance

- ▶  $\delta$ -matching between barcodes  $U, V$ :
  - ▶ matched intervals have  $\delta$ -close endpoints
  - ▶ unmatched intervals are  $2\delta$ -trivial (shorter than  $2\delta$ )


*Bottleneck distance:*  $d_B(U, V) = \inf\{\delta \mid \exists \delta\text{-matching } U \rightarrow V\}$



# Bottleneck distance as an interleaving distance

- ▶  $\delta$ -matching between barcodes  $U, V$ :
  - ▶ matched intervals have  $\delta$ -close endpoints
  - ▶ unmatched intervals are  $2\delta$ -trivial (shorter than  $2\delta$ )

Bottleneck distance:  $d_B(U, V) = \inf\{\delta \mid \exists \delta\text{-matching } U \rightarrow V\}$




- ▶  $\delta$ -interleaving between diagrams  $X, Y$  indexed over  $\mathbb{R}$  (in any category):  
 natural transformations  $f_t : X_t \rightarrow Y_{t+\delta}, g_t : Y_t \rightarrow X_{t+\delta}$  yielding commutative diagrams

$$\begin{array}{ccccc}
 X_{t-\delta} & \longrightarrow & X_t & \longrightarrow & X_{t+\delta} \\
 & \searrow & & \nearrow & \\
 & & & & \\
 & \nearrow & & \searrow & \\
 Y_{t-\delta} & \longrightarrow & Y_t & \longrightarrow & Y_{t+\delta}
 \end{array}
 \quad \forall t \in \mathbb{R}.$$

# Bottleneck distance as an interleaving distance

- ▶  $\delta$ -matching between barcodes  $U, V$ :
  - ▶ matched intervals have  $\delta$ -close endpoints
  - ▶ unmatched intervals are  $2\delta$ -trivial (shorter than  $2\delta$ )



Bottleneck distance:  $d_B(U, V) = \inf\{\delta \mid \exists \delta\text{-matching } U \dashv\dashv V\}$

- ▶  $\delta$ -interleaving between diagrams  $X, Y$  indexed over  $\mathbb{R}$  (in any category):  
 natural transformations  $f_t : X_t \rightarrow Y_{t+\delta}, g_t : Y_t \rightarrow X_{t+\delta}$  yielding commutative diagrams


$$\begin{array}{ccccc}
 X_{t-\delta} & \longrightarrow & X_t & \longrightarrow & X_{t+\delta} \\
 & \searrow & & \nearrow & \\
 & & & & \\
 & \nearrow & & \searrow & \\
 & & & & \\
 Y_{t-\delta} & \longrightarrow & Y_t & \longrightarrow & Y_{t+\delta}
 \end{array}
 \quad \forall t \in \mathbb{R}.$$

Interleaving distance:  $d_B(U, V) = \inf\{\delta \mid \exists \delta\text{-interleaving } X \leftrightarrow Y\}$



# Bottleneck distance as an interleaving distance

- ▶  $\delta$ -matching between barcodes  $U, V$ :
  - ▶ matched intervals have  $\delta$ -close endpoints
  - ▶ unmatched intervals are  $2\delta$ -trivial (shorter than  $2\delta$ )



Bottleneck distance:  $d_B(U, V) = \inf\{\delta \mid \exists \delta\text{-matching } U \dashv\dashv V\}$

- ▶  $\delta$ -interleaving between diagrams  $X, Y$  indexed over  $\mathbb{R}$  (in any category):  
natural transformations  $f_t : X_t \rightarrow Y_{t+\delta}, g_t : Y_t \rightarrow X_{t+\delta}$  yielding commutative diagrams

$$\begin{array}{ccccc}
 X_{t-\delta} & \longrightarrow & X_t & \longrightarrow & X_{t+\delta} \\
 & \searrow & & \nearrow & \\
 & & & & \\
 & \nearrow & & \searrow & \\
 & & & & \\
 Y_{t-\delta} & \longrightarrow & Y_t & \longrightarrow & Y_{t+\delta}
 \end{array} \quad \forall t \in \mathbb{R}.$$

Interleaving distance:  $d_B(U, V) = \inf\{\delta \mid \exists \delta\text{-interleaving } X \leftrightarrow Y\}$

## Proposition

$d_I = d_B$  (using the equivalence  $\mathbf{Barc} \simeq \mathbf{Mch}^{\mathbb{R}}$ ).

## Non-functoriality of persistence barcodes

Can a pfd persistence module  $M : \mathbf{vect}^{\mathbb{R}}$  be mapped to its barcode  $B(M) : \mathbf{Mch}^{\mathbb{R}}$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$  (or  $\mathbf{vect}^{\mathbb{R}} \rightarrow \mathbf{Mch}^{\mathbb{R}}$ )?

- ▶ This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.

## Non-functoriality of persistence barcodes

Can a pfd persistence module  $M : \mathbf{vect}^{\mathbb{R}}$  be mapped to its barcode  $B(M) : \mathbf{Mch}^{\mathbb{R}}$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$  (or  $\mathbf{vect}^{\mathbb{R}} \rightarrow \mathbf{Mch}^{\mathbb{R}}$ )?

- ▶ This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



# Non-functoriality of persistence barcodes

Can a pfd persistence module  $M : \mathbf{vect}^{\mathbb{R}}$  be mapped to its barcode  $B(M) : \mathbf{Mch}^{\mathbb{R}}$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$  (or  $\mathbf{vect}^{\mathbb{R}} \rightarrow \mathbf{Mch}^{\mathbb{R}}$ )?

- ▶ This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



## Proposition

*There is no functor  $\mathbf{vect} \rightarrow \mathbf{Mch}$  sending every vector space  $V$  to a set of cardinality  $\dim V$  (equivalently: sending a linear map  $f$  to a matching of cardinality  $\text{rank } f$ ).*

# Non-functoriality of persistence barcodes

Can a pfd persistence module  $M : \mathbf{vect}^{\mathbb{R}}$  be mapped to its barcode  $B(M) : \mathbf{Mch}^{\mathbb{R}}$  by a functor  $B : \mathbf{vect} \rightarrow \mathbf{Mch}$  (or  $\mathbf{vect}^{\mathbb{R}} \rightarrow \mathbf{Mch}^{\mathbb{R}}$ )?

- ▶ This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



## Proposition

*There is no functor  $\mathbf{vect} \rightarrow \mathbf{Mch}$  sending every vector space  $V$  to a set of cardinality  $\dim V$  (equivalently: sending a linear map  $f$  to a matching of cardinality  $\text{rank } f$ ).*

But: there is a barcode functor for subcategories of monos/epis of persistence modules  $\mathbf{vect}^{\mathbb{R}}$ :

# Structure of persistence sub-/quotient modules

## Proposition

*Let  $M \twoheadrightarrow N$  be an epimorphism.*

*Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that if  $J$  is mapped to  $I$ , then*

- ▶  *$I$  and  $J$  are aligned below, and*
- ▶  *$I$  bounds  $J$  above.*

*This construction is functorial.*

*Dually, there is an injection  $B(M) \hookrightarrow B(N)$  for monomorphisms  $M \hookrightarrow N$ .*



# Persistence sub-/quotient modules and their matching diagrams

Structure of persistence sub-/quotient modules, rephrased for  $\mathbf{Mch}^{\mathbb{R}}$ :

## Proposition


*There is a functor from epimorphisms of persistence modules to epimorphisms of matching diagrams.*

*Dually, there is a functor from monos to monos.*



## Algebraic stability via induced matchings


Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):






## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




Induced  $\delta$ -matching of barcodes:


$$\text{—————} B(N)$$

# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




Induced  $\delta$ -matching of barcodes:




# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):




Induced  $\delta$ -matching of barcodes:




# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



Induced  $\delta$ -matching of barcodes:



$B(\text{im } N(-\delta, \delta))$


$B(\text{im } f)$

$B(N(\delta))$


$B(N)$

# Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



Induced  $\delta$ -matching of barcodes:



## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \operatorname{im} f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:

## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \operatorname{im} f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:

- compose the functorial injections  $B(M) \leftarrow B(\operatorname{im} f) \hookrightarrow B(N)$  from before to a matching

$$\chi(f) : B(M) \rightarrowtail B(N).$$





## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \operatorname{im} f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:

- compose the functorial injections  $B(M) \hookleftarrow B(\operatorname{im} f) \hookrightarrow B(N)$  from before to a matching

$$\chi(f) : B(M) \dashrightarrow B(N).$$

### Theorem

Assume that  $\ker f$  is  $\delta$ -trivial. If  $I$  is matched to  $J$ , then

- (i)  $I$  overlaps  $J$ , and  $J$  overlaps  $I(\delta)$ .
- (ii) Any unmatched interval of  $B(M)$  is  $\delta$ -trivial.

There is a dual statement for  $\operatorname{coker} f$   $\delta$ -trivial.



# The categorified induced matching theorem

Induced matching theorem, rephrased in  $\mathbf{Mch}^{\mathbb{R}}$ :

## Theorem

*If  $f : M \rightarrow N$  has  $\delta$ -trivial (co)kernel, then so does the induced matching  $\chi(f) : B(M) \rightarrowtail B(N)$ .*




# The categorified induced matching theorem

Induced matching theorem, rephrased in  $\mathbf{Mch}^{\mathbb{R}}$ :

## Theorem

*If  $f : M \rightarrow N$  has  $\delta$ -trivial (co)kernel, then so does the induced matching  $\chi(f) : B(M) \rightarrow B(N)$ .*



Note:

- ▶ We always have  $B(\operatorname{im} f) = \operatorname{im} \chi(f)$  by construction.
- ▶ But  $\ker \chi(f)$  may differ from  $B(\ker f)$ .
- ▶ The induced matching may strictly decrease the triviality of the kernel.

## A general criterion for $\delta$ -trivial (co)kernels

### Lemma

*For a natural transformation  $f : M \rightarrow N$  between diagrams  $M, N : \mathbf{R} \rightarrow \mathbf{A}$  in a Puppe-exact  $\mathbf{A}$ , consider the epi-mono factorization*

$$f : M \xrightarrow{q} \operatorname{im} f \xhookrightarrow{i} N.$$

## A general criterion for $\delta$ -trivial (co)kernels

### Lemma

*For a natural transformation  $f : M \rightarrow N$  between diagrams  $M, N : \mathbf{R} \rightarrow \mathbf{A}$  in a Puppe-exact  $\mathbf{A}$ , consider the epi-mono factorization*

$$f : M \xrightarrow{q} \operatorname{im} f \xrightarrow{i} N.$$

*Moreover, let  $s : M \rightarrow M(\delta)$  be given by the internal morphisms  $\{M_t \rightarrow M_{t+\delta}\}_{t \in \mathbb{R}}$ .*

## A general criterion for $\delta$ -trivial (co)kernels

### Lemma

For a natural transformation  $f : M \rightarrow N$  between diagrams  $M, N : \mathbf{R} \rightarrow \mathbf{A}$  in a Puppe-exact  $\mathbf{A}$ , consider the epi-mono factorization

$$f : M \xrightarrow{q} \operatorname{im} f \xrightarrow{i} N.$$

Moreover, let  $s : M \rightarrow M(\delta)$  be given by the internal morphisms  $\{M_t \rightarrow M_{t+\delta}\}_{t \in \mathbb{R}}$ .

By a diagram chase, the following are equivalent:

- (i)  $\ker f$  is  $\delta$ -trivial;
- (ii) the image epimorphism  $M \twoheadrightarrow \operatorname{im} s$  factors (through  $q$ ) as

$$M \xrightarrow{q} \operatorname{im} f \twoheadrightarrow \operatorname{im} s.$$

A dual statement holds for  $\operatorname{coker} f$ .

# The induced matching theorem for (monos and) epis

## Lemma

*Let  $f : M \twoheadrightarrow N$  be an epimorphism. If  $\ker f$  is  $\delta$ -trivial, then so is  $\ker \chi(f)$ . (Dually for monos.)*

# The induced matching theorem for (monos and) epis

## Lemma

*Let  $f : M \twoheadrightarrow N$  be an epimorphism. If  $\ker f$  is  $\delta$ -trivial, then so is  $\ker \chi(f)$ . (Dually for monos.)*

## Proof outline.

As before, let  $s : M \rightarrow M(\delta)$  be the shift morphism.



# The induced matching theorem for (monos and) epis

## Lemma

*Let  $f : M \twoheadrightarrow N$  be an epimorphism. If  $\ker f$  is  $\delta$ -trivial, then so is  $\ker \chi(f)$ . (Dually for monos.)*

## Proof outline.

As before, let  $s : M \rightarrow M(\delta)$  be the shift morphism.

- ▶ Since  $\ker f$  is  $\delta$ -trivial, the image epimorphism  $M \twoheadrightarrow \operatorname{im} s$  factors through  $f$  as

$$M \xrightarrow{f} N \twoheadrightarrow \operatorname{im} s.$$

# The induced matching theorem for (monos and) epis

## Lemma

*Let  $f : M \twoheadrightarrow N$  be an epimorphism. If  $\ker f$  is  $\delta$ -trivial, then so is  $\ker \chi(f)$ . (Dually for monos.)*

## Proof outline.

As before, let  $s : M \rightarrow M(\delta)$  be the shift morphism.

- ▶ Since  $\ker f$  is  $\delta$ -trivial, the image epimorphism  $M \twoheadrightarrow \operatorname{im} s$  factors through  $f$  as

$$M \xrightarrow{f} N \twoheadrightarrow \operatorname{im} s.$$

- ▶ By functoriality on epis, the canonical epimorphism  $B(M) \twoheadrightarrow \operatorname{im} \chi(s)$  factors through  $\chi(f)$ :

$$B(M) \xrightarrow{\chi(f)} B(N) \twoheadrightarrow \operatorname{im} \chi(s).$$

# The induced matching theorem for (monos and) epis

## Lemma

*Let  $f : M \twoheadrightarrow N$  be an epimorphism. If  $\ker f$  is  $\delta$ -trivial, then so is  $\ker \chi(f)$ . (Dually for monos.)*

## Proof outline.

As before, let  $s : M \rightarrow M(\delta)$  be the shift morphism.

- ▶ Since  $\ker f$  is  $\delta$ -trivial, the image epimorphism  $M \twoheadrightarrow \operatorname{im} s$  factors through  $f$  as

$$M \xrightarrow{f} N \twoheadrightarrow \operatorname{im} s.$$

- ▶ By functoriality on epis, the canonical epimorphism  $B(M) \twoheadrightarrow \operatorname{im} \chi(s)$  factors through  $\chi(f)$ :

$$B(M) \xrightarrow{\chi(f)} B(N) \twoheadrightarrow \operatorname{im} \chi(s).$$

- ▶ Equivalently,  $\ker \chi(f)$  is  $\delta$ -trivial.



# The induced matching theorem

## Theorem

*If  $f : M \rightarrow N$  has  $\delta$ -trivial (co)kernel, then so does  $\chi(f)$ .*

# The induced matching theorem

## Theorem

*If  $f : M \rightarrow N$  has  $\delta$ -trivial (co)kernel, then so does  $\chi(f)$ .*

## Proof outline.

- ▶ Consider epi-mono factorization

$$f : M \xrightarrow{q} \operatorname{im} f \xrightarrow{i} N.$$

# The induced matching theorem

## Theorem

*If  $f : M \rightarrow N$  has  $\delta$ -trivial (co)kernel, then so does  $\chi(f)$ .*

## Proof outline.

- ▶ Consider epi-mono factorization

$$f : M \xrightarrow{q} \operatorname{im} f \xrightarrow{i} N.$$

- ▶ If  $\ker q = \ker f$  is  $\delta$ -trivial, then so is  $\ker \chi(q) = \ker \chi(f)$ .

# The induced matching theorem

## Theorem

*If  $f : M \rightarrow N$  has  $\delta$ -trivial (co)kernel, then so does  $\chi(f)$ .*

## Proof outline.

- ▶ Consider epi-mono factorization

$$f : M \xrightarrow{q} \operatorname{im} f \xrightarrow{i} N.$$

- ▶ If  $\ker q = \ker f$  is  $\delta$ -trivial, then so is  $\ker \chi(q) = \ker \chi(f)$ .
- ▶ Dually for  $\operatorname{coker} i = \operatorname{coker} f$ .



# The algebraic stability theorem (categorically)

## Theorem

*Two pfd persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if and only if their barcodes  $B(M)$  and  $B(N)$  are  $\delta$ -interleaved. In particular,  $d_I(M, N) = d_I(B(M), B(N))$ .*



# The algebraic stability theorem (categorically)

## Theorem

*Two pfd persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if and only if their barcodes  $B(M)$  and  $B(N)$  are  $\delta$ -interleaved. In particular,  $d_I(M, N) = d_I(B(M), B(N))$ .*

## Proof outline.

Forward direction:

- ▶ A  $\delta$ -interleaving morphism  $f : M \rightarrow N(\delta)$  has  $2\delta$ -trivial kernel and cokernel.

# The algebraic stability theorem (categorically)

## Theorem

*Two pfd persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if and only if their barcodes  $B(M)$  and  $B(N)$  are  $\delta$ -interleaved. In particular,  $d_I(M, N) = d_I(B(M), B(N))$ .*

## Proof outline.

Forward direction:

- ▶ A  $\delta$ -interleaving morphism  $f : M \rightarrow N(\delta)$  has  $2\delta$ -trivial kernel and cokernel.
- ▶ By the induced matching theorem, the same is true for  $\chi(f) : B(M) \rightarrow B(N(\delta))$ .

# The algebraic stability theorem (categorically)

## Theorem

*Two pfd persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if and only if their barcodes  $B(M)$  and  $B(N)$  are  $\delta$ -interleaved. In particular,  $d_I(M, N) = d_I(B(M), B(N))$ .*

## Proof outline.

Forward direction:

- ▶ A  $\delta$ -interleaving morphism  $f : M \rightarrow N(\delta)$  has  $2\delta$ -trivial kernel and cokernel.
- ▶ By the induced matching theorem, the same is true for  $\chi(f) : B(M) \rightarrow B(N(\delta))$ .
- ▶ This is equivalent to  $B(M)$  and  $B(N)$  being  $\delta$ -interleaved in  $\mathbf{Mch}^{\mathbb{R}}$ .

# The algebraic stability theorem (categorically)

## Theorem

*Two pfd persistence modules  $M$  and  $N$  are  $\delta$ -interleaved if and only if their barcodes  $B(M)$  and  $B(N)$  are  $\delta$ -interleaved. In particular,  $d_I(M, N) = d_I(B(M), B(N))$ .*

## Proof outline.

Forward direction:

- ▶ A  $\delta$ -interleaving morphism  $f : M \rightarrow N(\delta)$  has  $2\delta$ -trivial kernel and cokernel.
- ▶ By the induced matching theorem, the same is true for  $\chi(f) : B(M) \rightarrow B(N(\delta))$ .
- ▶ This is equivalent to  $B(M)$  and  $B(N)$  being  $\delta$ -interleaved in  $\mathbf{Mch}^{\mathbb{R}}$ .

Converse direction:

- ▶ Apply the free functor  $\mathbf{Mch} \rightarrow \mathbf{Vect}$ .



## Barcodes from scratch

- ▶ Goal: construct barcode/matching diagram of persistence module *without* decomposition
- ▶ At each index in the matching diagram, the set should be natural numbers  $\{1, \dots, n\}$

## Barcodes from scratch

- ▶ Goal: construct barcode/matching diagram of persistence module *without* decomposition
- ▶ At each index in the matching diagram, the set should be natural numbers  $\{1, \dots, n\}$




Approach:

- ▶ Order intervals in barcode (lexicographically) by birth (increasing), then death (decreasing)

## Barcodes from scratch

- ▶ Goal: construct barcode/matching diagram of persistence module *without* decomposition
- ▶ At each index in the matching diagram, the set should be natural numbers  $\{1, \dots, n\}$




Approach:

- ▶ Order intervals in barcode (lexicographically) by birth (increasing), then death (decreasing)
- ▶ At each index  $t$ , enumerate the intervals containing  $t$  in that order

## Barcodes from scratch

- ▶ Goal: construct barcode/matching diagram of persistence module *without* decomposition
- ▶ At each index in the matching diagram, the set should be natural numbers  $\{1, \dots, n\}$




### Approach:

- ▶ Order intervals in barcode (lexicographically) by birth (increasing), then death (decreasing)
- ▶ At each index  $t$ , enumerate the intervals containing  $t$  in that order
- ▶ Between any two indices, match numbers according to their associated barcode intervals



## Barcodes from scratch

- ▶ Goal: construct barcode/matching diagram of persistence module *without* decomposition
- ▶ At each index in the matching diagram, the set should be natural numbers  $\{1, \dots, n\}$



### Approach:

- ▶ Order intervals in barcode (lexicographically) by birth (increasing), then death (decreasing)
- ▶ At each index  $t$ , enumerate the intervals containing  $t$  in that order
- ▶ Between any two indices, match numbers according to their associated barcode intervals

## A rank formula for barcodes


$M$ : persistence module,  $D$ : corresponding barcode (ordered by birth, then death).



## A rank formula for barcodes

$M$ : persistence module,  $D$ : corresponding barcode (ordered by birth, then death).


- ▶ Which  $j$  are in  $D_t$ ? This is  $\{j \mid j \leq \dim M_t\}$ .



## A rank formula for barcodes

$M$ : persistence module,  $D$ : corresponding barcode (ordered by birth, then death).


- ▶ Which  $j$  are in  $D_t$ ? This is  $\{j \mid j \leq \dim M_t\}$ .
- ▶ Which  $j \in D_t$  are matched to some  $i \in D_s$ ?  
This is  $\{j \mid j \leq \text{rank } M_{s,t}\}$ .



## A rank formula for barcodes

$M$ : persistence module,  $D$ : corresponding barcode (ordered by birth, then death).


- ▶ Which  $j$  are in  $D_t$ ? This is  $\{j \mid j \leq \dim M_t\}$ .
- ▶ Which  $j \in D_t$  are matched to some  $i \in D_s$ ?  
This is  $\{j \mid j \leq \text{rank } M_{s,t}\}$ .
- ▶ Given  $j \in D_t$ , to which  $i \in D_s$  is it matched?  
The difference  $i - j$  is the number of bars that
  - (a) are born before the  $j$ th interval at index  $t$ , and
  - (b) die between  $s$  and  $t$ :



## A rank formula for barcodes

$M$ : persistence module,  $D$ : corresponding barcode (ordered by birth, then death).


- ▶ Which  $j$  are in  $D_t$ ? This is  $\{j \mid j \leq \dim M_t\}$ .
- ▶ Which  $j \in D_t$  are matched to some  $i \in D_s$ ?  
This is  $\{j \mid j \leq \text{rank } M_{s,t}\}$ .
- ▶ Given  $j \in D_t$ , to which  $i \in D_s$  is it matched?  
The difference  $i - j$  is the number of bars that
  - (a) are born before the  $j$ th interval at index  $t$ , and
  - (b) die between  $s$  and  $t$ :
- ▶ Given  $j \in D_t$ , what are the lower bounds for the  $j$ th interval at index  $t$ ? This is  $\{r < t \mid \text{rank } M_{r,t} < j\}$ .



## A rank formula for barcodes

$M$ : persistence module,  $D$ : corresponding barcode (ordered by birth, then death).


- ▶ Which  $j$  are in  $D_t$ ? This is  $\{j \mid j \leq \dim M_t\}$ .
- ▶ Which  $j \in D_t$  are matched to some  $i \in D_s$ ?  
This is  $\{j \mid j \leq \text{rank } M_{s,t}\}$ .
- ▶ Given  $j \in D_t$ , to which  $i \in D_s$  is it matched?  
The difference  $i - j$  is the number of bars that
  - are born before the  $j$ th interval at index  $t$ , and
  - die between  $s$  and  $t$ :
- ▶ Given  $j \in D_t$ , what are the lower bounds for the  $j$ th interval at index  $t$ ? This is  $\{r < t \mid \text{rank } M_{r,t} < j\}$ .
- ▶ Together, this yields
$$i - j = \max \{ \text{rank } M_{r,s} - \text{rank } M_{r,t} \mid r < s, \text{rank } M_{r,t} < j \}.$$



## A rank formula for barcodes

$M$ : persistence module,  $D$ : corresponding barcode (ordered by birth, then death).

- ▶ Which  $j$  are in  $D_t$ ? This is  $\{j \mid j \leq \dim M_t\}$ .
- ▶ Which  $j \in D_t$  are matched to some  $i \in D_s$ ?  
This is  $\{j \mid j \leq \text{rank } M_{s,t}\}$ .
- ▶ Given  $j \in D_t$ , to which  $i \in D_s$  is it matched?  
The difference  $i - j$  is the number of bars that
  - (a) are born before the  $j$ th interval at index  $t$ , and
  - (b) die between  $s$  and  $t$ :
- ▶ Given  $j \in D_t$ , what are the lower bounds for the  $j$ th interval at index  $t$ ? This is  $\{r < t \mid \text{rank } M_{r,t} < j\}$ .
- ▶ Together, this yields
$$i - j = \max \{ \text{rank } M_{r,s} - \text{rank } M_{r,t} \mid r < s, \text{rank } M_{r,t} < j \}.$$




This specifies the barcode of  $M$  (as a matching diagram) based on ranks only.



# Functoriality

The previous construction extends to a functor of epimorphisms  $M \rightarrow N$  from persistence modules to matching diagrams.




# Functoriality

The previous construction extends to a functor of epimorphisms  $M \rightarrow N$  from persistence modules to matching diagrams.

- ▶ For the record:  $i \in D(M)_t$  is matched to  $j \in D(N)_t$  iff

$$\max \{ \text{rank } M_{s,t} - \text{rank } N_{s,t} \mid s < t, \text{rank } N_{s,t} < j \}.$$




# Functoriality

The previous construction extends to a functor of epimorphisms  $M \rightarrow N$  from persistence modules to matching diagrams.

- ▶ For the record:  $i \in D(M)_t$  is matched to  $j \in D(N)_t$  iff

$$\max \{ \text{rank } M_{s,t} - \text{rank } N_{s,t} \mid s < t, \text{rank } N_{s,t} < j \}.$$

- ▶ Dually for monomorphisms




# Functoriality

The previous construction extends to a functor of epimorphisms  $M \rightarrow N$  from persistence modules to matching diagrams.

- ▶ For the record:  $i \in D(M)_t$  is matched to  $j \in D(N)_t$  iff

$$\max \{ \text{rank } M_{s,t} - \text{rank } N_{s,t} \mid s < t, \text{rank } N_{s,t} < j \}.$$

- ▶ Dually for monomorphisms
- ▶ Epi-mono factorization yields an induced matching




# Functoriality

The previous construction extends to a functor of epimorphisms  $M \rightarrow N$  from persistence modules to matching diagrams.

- ▶ For the record:  $i \in D(M)_t$  is matched to  $j \in D(N)_t$  iff

$$\max \{ \text{rank } M_{s,t} - \text{rank } N_{s,t} \mid s < t, \text{rank } N_{s,t} < j \}.$$

- ▶ Dually for monomorphisms
- ▶ Epi-mono factorization yields an induced matching
- ▶ Coincides with previous induced matching  
(under the equivalence  $\mathbf{Barc} \simeq \mathbf{Mch}^{\mathbb{R}}$ )



# Functoriality

The previous construction extends to a functor of epimorphisms  $M \rightarrow N$  from persistence modules to matching diagrams.

- ▶ For the record:  $i \in D(M)_t$  is matched to  $j \in D(N)_t$  iff

$$\max \{ \text{rank } M_{s,t} - \text{rank } N_{s,t} \mid s < t, \text{rank } N_{s,t} < j \}.$$

- ▶ Dually for monomorphisms
- ▶ Epi-mono factorization yields an induced matching
- ▶ Coincides with previous induced matching  
(under the equivalence  $\mathbf{Barc} \simeq \mathbf{Mch}^{\mathbb{R}}$ )

Obtain induced matching and algebraic stability theorems without an interval decomposition

