

Topological Data Analysis: An Introduction to Persistent Homology


Part 2: complexes and computation


Ulrich Bauer





Sep 5, 2019


Machine learning summer school 2019
Skoltech, Moscow
























































Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?

Connecting Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have


$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?
- Are they related by a sequence of *simplicial collapses*?

Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.




Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex that has a unique proper coface.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.

- A *free face* of K is a simplex that has a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face and its proper coface.



Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.



- A *free face* of K is a simplex that has a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face and its proper coface.
- L is homotopy-equivalent to K . In particular, they have the same homology.

Simplicial collapses

Definition (Whitehead 1938)

Let K be a simplicial complex.



- A *free face* of K is a simplex that has a unique proper coface.
- We can *collapse* K to a subcomplex L by removing a free face and its proper coface.
- L is homotopy-equivalent to K . In particular, they have the same homology.


If there is a sequence of such elementary collapses from K to M , we say that K *collapses* to M (written $K \searrow M$).

Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2017)

Čech, Delaunay–Čech, Delaunay, and Wrap complexes are homotopy equivalent through a sequence of collapses

$$\text{Cech}_r X \searrow \text{Cech}_r X \cap \text{Del} X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$




Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2017)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent through a sequence of collapses

$$\text{Cech}_r X \searrow \text{Cech}_r X \cap \text{Del} X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$




- Extends to weighted Delaunay triangulations
- All collapses are induced by a single *discrete gradient field*
- This yields explicit isomorphisms in persistent homology

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.



Discrete Morse theory



Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory


Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



•₁

Discrete Morse theory


Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory



Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory



Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory



Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory



Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes



Discrete Morse theory



Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- level sets $f^{-1}(t)$ form a discrete vector field (*gradient of f*)



Discrete Morse theory

Definition (Forman 1998)



A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, with L a facet of U .

- indicated by an arrow from L to U
- The singletons are called *critical simplices*.

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:


- sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- level sets $f^{-1}(t)$ form a discrete vector field (*gradient of f*)

If $f^{-1}(t) = \{Q\}$ then t is a *critical value*.



Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.



Collapses from Morse functions and gradients


Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.

Corollary

K is homotopy-equivalent to a CW complex
with one d -cell for each critical d -simplex.



Collapses from Morse functions and gradients


Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.

Corollary

K is homotopy-equivalent to a CW complex
with one d -cell for each critical d -simplex.



Let V be a discrete gradient field on a simplicial complex K , and let L be a subcomplex of K .

Corollary

If $K \setminus L$ is a union of pairs of V , then $K \searrow L$.

We say that V induces the collapse $K \searrow L$.



Collapses from Morse functions and gradients


Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

If $(s, t]$ contains no critical value of f , then $K_t \searrow K_s$.

Corollary

K is homotopy-equivalent to a CW complex
with one d -cell for each critical d -simplex.




Let V be a discrete gradient field on a simplicial complex K , and let L be a subcomplex of K .

Corollary

If $K \setminus L$ is a union of pairs of V , then $K \searrow L$.


We say that V induces the collapse $K \searrow L$.



Čech and Delaunay functions

Čech and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)


Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)




Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)

Two radius functions:

- Čech radius function $f_{\text{Čech}}(Q)$:
radius of smallest enclosing sphere of Q




$\check{\text{C}}\text{ech}$ and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Consider *abstract simplices* (nonempty subsets $Q \subseteq X$)

Two radius functions:


- *$\check{\text{C}}\text{ech radius function } f_{\text{Cech}}(Q)$:*
radius of *smallest enclosing sphere* of Q
- *$\text{Delaunay radius function } f_{\text{Del}}(Q)$:*
radius of *smallest empty circumsphere*
 - defined only on $\text{Del}(X)$



Čech and Delaunay complexes from functions

For any radius r :


- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$



Čech and Delaunay complexes from functions

For any radius r :


- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$
- $\text{DelCech}_r(X) = \text{Del}(X) \cap \text{Cech}_r(X)$
 - restriction of Čech complex to Delaunay simplices



Čech and Delaunay complexes from functions

For any radius r :


- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$
- $\text{DelCech}_r(X) = \text{Del}(X) \cap \text{Cech}_r(X)$
 - restriction of Čech complex to Delaunay simplices
- $\text{Del}_r(X) = f_{\text{Del}}^{-1}(-\infty, r]$
 - simplices with smallest empty circumradius $\leq r$



Čech and Delaunay complexes from functions

For any radius r :

- $\text{Cech}_r(X) = f_{\text{Cech}}^{-1}(-\infty, r]$
 - simplices with smallest enclosing radius $\leq r$
- $\text{DelCech}_r(X) = \text{Del}(X) \cap \text{Cech}_r(X)$
 - restriction of Čech complex to Delaunay simplices
- $\text{Del}_r(X) = f_{\text{Del}}^{-1}(-\infty, r]$
 - simplices with smallest empty circumradius $\leq r$





Unfortunately...

Neither the Čech nor the Delaunay functions
are discrete Morse functions!

Unfortunately...

Neither the Čech nor the Delaunay functions
are discrete Morse functions!

- Example: two simplices Q, Q' with $f_{\text{Čech}}(Q) = f_{\text{Čech}}(Q')$
where neither is a face of the other:




Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex K is a partition of the simplices into *intervals* of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$

- indicated by an arrow from L to U



Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)



A *generalized discrete vector field* on a simplicial complex K is a partition of the simplices into *intervals* of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$

- indicated by an arrow from L to U

A *generalized discrete Morse function* $f : K \rightarrow \mathbb{R}$ satisfies:


- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes (for all $t \in \mathbb{R}$)
- the level sets $f^{-1}(t)$ form a generalized vector field (the *discrete gradient* of f)



Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical face interval $[L, U] \in V$:




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical face interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical face interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical face interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.



Therefore the collapsing theorems also hold for generalized discrete Morse functions.

Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*


Morse theory of Čech and Delaunay complexes

Proposition

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

The following are equivalent:

- Q is a critical simplex of $f_{\text{Čech}}$
- Q is a critical simplex of f_{Del}
- $f_{\text{Čech}}(Q) = f_{\text{Del}}(Q)$
- Q is a centered Delaunay simplex
(containing the circumcenter in the interior)




$\check{\text{C}}$ ech face intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $R \subseteq X$ has the same smallest enclosing sphere S iff



Q


Čech face intervals

Lemma


Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $R \subseteq X$ has the same smallest enclosing sphere S iff

On $S \subseteq R$



Q



On S


Čech face intervals

Lemma


Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $R \subseteq X$ has the same smallest enclosing sphere S iff


$$\text{On } S \subseteq R \subseteq \text{Encl } S.$$



Q



$\text{On } S$



$\text{Encl } S$


$\check{\text{C}}\text{ech}$ face intervals

Lemma


Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .

Then $R \subseteq X$ has the same smallest enclosing sphere S iff


$$R \in [\text{On } S, \text{Encl } S].$$



Q



$\text{On } S$



$\text{Encl } S$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.

Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.


Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.



Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$






Delaunay face intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $R \subseteq X$ has the same smallest empty circumsphere S iff

$$R \in [\text{Front } S, \text{On } S].$$



Selective Delaunay complexes

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{ll} \underset{r,z}{\text{minimize}} & r \\ \text{subject to} & \|z - q\| \leq r, \quad q \in Q, \\ & \|z - e\| \geq r, \quad e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{ll} \underset{r,z}{\text{minimize}} & r \\ \text{subject to} & \|z - q\| \leq r, \quad q \in Q, \\ & \|z - e\| \geq r, \quad e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{ll} \text{minimize}_{r,z} & r^2 \\ \text{subject to} & \|z - q\|^2 \leq r^2, \quad q \in Q, \\ & \|z - e\|^2 \geq r^2, \quad e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{lll} \text{minimize}_{a,z} & \|z\|^2 - a & (r^2 = \|z\|^2 - a) \\ \text{subject to} & \|z - q\|^2 \leq \|z\|^2 - a, & q \in Q, \\ & \|z - e\|^2 \geq \|z\|^2 - a, & e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{ll} \text{minimize}_{a,z} & \|z\|^2 - a \\ \text{subject to} & \|z\|^2 - 2\langle z, q \rangle + \|q\|^2 \leq \|z\|^2 - a, \quad q \in Q, \\ & \|z\|^2 - 2\langle z, e \rangle + \|e\|^2 \geq \|z\|^2 - a, \quad e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

$$\begin{array}{ll} \underset{a,z}{\text{minimize}} & \|z\|^2 - a \\ \text{subject to} & 2\langle z, q \rangle - a - \|q\|^2 \geq 0, \quad \forall q \in Q, \\ & 2\langle z, e \rangle - a - \|e\|^2 \leq 0, \quad \forall e \in E. \end{array}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem of the form

$$\begin{array}{ll} \underset{x}{\text{minimize}} & f(x) \\ \text{subject to} & g_j(x) \geq 0, \quad \forall j \in J, \\ & g_k(x) = 0, \quad \forall k \in K, \\ & g_l(x) \leq 0, \quad \forall l \in L. \end{array}$$

where the function f is convex and g_i are affine ($i \in I = J \cup K \cup L$).

The Karush–Kuhn–Tucker optimality conditions

Consider an optimization problem of the form

$$\begin{array}{ll} \underset{x}{\text{minimize}} & f(x) \\ \text{subject to} & g_j(x) \geq 0, \quad \forall j \in J, \\ & g_k(x) = 0, \quad \forall k \in K, \\ & g_l(x) \leq 0, \quad \forall l \in L. \end{array}$$

where the function f is convex and g_i are affine ($i \in I = J \cup K \cup L$).

Theorem (Karush 1939, Kuhn–Tucker 1951)

A feasible x is optimal iff there exist $(\lambda_i)_{i \in I}$ such that

$$\nabla f(x) + \sum_{i \in I} \lambda_i \nabla g_i(x) = 0, \quad (\text{stationarity})$$

$$\lambda_i g_i(x) = 0, \quad \forall i \in I, \quad (\text{complementary slackness})$$

$$\lambda_j \leq 0, \quad \forall j \in J, \quad (\text{dual feasibility})$$

$$\lambda_l \geq 0, \quad \forall l \in L.$$

KKT conditions for the smallest sphere problem

The KKT conditions for our sphere optimization problem are:

Proposition

*A sphere S enclosing Q and excluding E is minimal iff
its center z can be written as an affine combination*

$$z = \sum_{x \in Q \cup E} \lambda_x x, \quad 1 = \sum_{x \in Q \cup E} \lambda_x$$

such that

- $\lambda_x = 0$ whenever x does not lie on S ,
- $\lambda_x \leq 0$ whenever $x \in E \setminus Q$, and
- $\lambda_x \geq 0$ whenever $x \in Q \setminus E$.

Čech and Delaunay face intervals from KKT

The Karush–Kuhn–Tucker optimality conditions yield:

Proposition (Geometric KKT conditions)

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $Q \in [\text{Front } S, \text{Encl } S]$ (i.e., $\text{Front } S \subseteq Q \subseteq \text{Encl } S$), and
- $E \in [\text{Back } S, \text{Excl } S]$ (i.e., $\text{Back } S \subseteq E \subseteq \text{Excl } S$).

Čech and Delaunay face intervals from KKT

The Karush–Kuhn–Tucker optimality conditions yield:

Proposition (Geometric KKT conditions)

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $Q \in [\text{Front } S, \text{Encl } S]$ (i.e., $\text{Front } S \subseteq Q \subseteq \text{Encl } S$), and
- $E \in [\text{Back } S, \text{Excl } S]$ (i.e., $\text{Back } S \subseteq E \subseteq \text{Excl } S$).

Corollary

- The Čech face intervals are of the form $[\text{On } S, \text{Encl } S]$.

$\check{\text{C}}\text{ech}$ and Delaunay face intervals from KKT

The Karush–Kuhn–Tucker optimality conditions yield:

Proposition (Geometric KKT conditions)


A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $Q \in [\text{Front } S, \text{Encl } S]$ (i.e., $\text{Front } S \subseteq Q \subseteq \text{Encl } S$), and
- $E \in [\text{Back } S, \text{Excl } S]$ (i.e., $\text{Back } S \subseteq E \subseteq \text{Excl } S$).


Corollary

- The $\check{\text{C}}\text{ech}$ face intervals are of the form $[\text{On } S, \text{Encl } S]$.
- The Delaunay face intervals are of the form $[\text{Front } S, \text{On } S]$.


The Delaunay gradient




The Delaunay gradient




The Delaunay gradient




The Delaunay gradient




The Delaunay gradient



The Delaunay gradient



The Delaunay gradient



Wrap complexes

Wrap complexes

Generalize and simplify the surface reconstruction method *Wrap* (Edelsbrunner 1995)

Wrap complexes

Generalize and simplify the surface reconstruction method *Wrap* (Edelsbrunner 1995)




Wrap complexes

Generalize and simplify the surface reconstruction method *Wrap* (Edelsbrunner 1995)




Wrap complexes

Generalize and simplify the surface reconstruction method *Wrap* (Edelsbrunner 1995)




Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.



Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.




Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Wrap complexes

Consider the Delaunay gradient V_D of $X \subset \mathbb{R}^D$.



Define $\text{Wrap}_r(X)$ as the smallest subcomplex of $\text{Del } X$ that

- contains all critical simplices with circumradius $\leq r$ and
- is a union of face intervals of V_D .

Corollary

The Delaunay face intervals induce a collapse $\text{Del}_r \searrow \text{Wrap}_r$.

Vietoris–Rips complexes

Vietoris–Rips complexes

Consider a finite metric space (X, d) .

The *Vietoris–Rips complex* is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid \text{diam } S \leq t\}$$

- all edges with pairwise distance $\leq t$
- all possible higher simplices

Vietoris–Rips complexes

Consider a finite metric space (X, d) .


The *Vietoris–Rips complex* is the simplicial complex


$$\text{Rips}_t(X) = \{S \subseteq X \mid \text{diam } S \leq t\}$$


- all edges with pairwise distance $\leq t$
- all possible higher simplices


For large t , $\text{Rips}_t(X)$ is the full simplex with vertices X

- Number of d -simplices is $\binom{|X|}{d+1}$
- Computation is one of the most important challenges in applied topology!









Another example data set

coil100-obj14-grid.pdf

A software for computing Vietoris–Rips persistence barcodes

- around 1000 lines of C++ code, no external dependencies
- support for
 - coefficients in a prime field $\mathbb{Z}/p\mathbb{Z}$
 - sparse distance matrices (for distance threshold)
- open source (<http://ripser.org>)
- online version (<http://live.ripser.org>)
- 2016 ATMCS Best New Software Award (joint with RIVET by M. Lesnick and M. Wright)

The four special ingredients

Main improvements to the standard algorithm:

- Clearing inessential columns [Chen, Kerber 2011]
- Computing cohomology [de Silva et al. 2011]
- Implicit matrix reduction
- Apparent and emergent pairs

The four special ingredients

Main improvements to the standard algorithm:

- Clearing inessential columns [Chen, Kerber 2011]
- Computing cohomology [de Silva et al. 2011]
- Implicit matrix reduction
- Apparent and emergent pairs

Lessons from previous work:

- Clearing and cohomology yield considerable speedup,
- but only when *both* are used in conjunction!

Matrix reduction

Computing homology

Computing homology $H_* = Z_*/B_*$:

- compute basis for boundaries $B_* = \text{im } \partial_*$
- extend to basis for cycles $Z_* = \ker \partial_*$
- new (non-boundary) basis cycles generate quotient Z_*/B_*

Computing homology

Computing homology $H_* = Z_*/B_*$:

- compute basis for boundaries $B_* = \text{im } \partial_*$
- extend to basis for cycles $Z_* = \ker \partial_*$
- new (non-boundary) basis cycles generate quotient Z_*/B_*

Computing *persistent* homology $H_* = Z_*/B_*$ (for a simplexwise filtration $K_i \subseteq K$):

- compute *filtered* basis for boundaries $B_* = \text{im } \partial_*$
- extend to basis for cycles $Z_* = \ker \partial_*$
- all basis cycles generate *persistent* homology

Persistence by matrix reduction

Given:

- D : boundary matrix for a simplexwise filtration $(K_i)_i$

Wanted:

- persistent homology barcodes in dimensions $d = 0, \dots, k$

Notation:

- M_j : column j of M ; m_{ij} : entry in row i and column j
- PivotIndex $M_j = \min\{i : m_{kj} = 0 \text{ for all } k > i\}$.

Persistence by matrix reduction

Given:

- D : boundary matrix for a simplexwise filtration $(K_i)_i$

Wanted:

- persistent homology barcodes in dimensions $d = 0, \dots, k$

Notation:

- M_j : column j of M ; m_{ij} : entry in row i and column j
- PivotIndex $M_j = \min\{i : m_{kj} = 0 \text{ for all } k > i\}$.

Computation: barcode is obtained by *matrix reduction* of D

- $R = D \cdot V$ reduced (non-zero columns have distinct pivots)
- V is regular upper triangular

Compatible basis cycles

For a reduced boundary matrix $R = D \cdot V$, call

$$I_b = \{i : R_i = 0\}$$

birth indices,

$$I_d = \{j : R_j \neq 0\}$$

death indices,

$$I_e = I_b \setminus \text{PivotIndices } R$$

essential indices.

(note: all pivot indices are birth indices).

Compatible basis cycles

For a reduced boundary matrix $R = D \cdot V$, call

$$I_b = \{i : R_i = 0\}$$

birth indices,

$$I_d = \{j : R_j \neq 0\}$$

death indices,

$$I_e = I_b \setminus \text{PivotIndices } R$$

essential indices.

(note: all pivot indices are birth indices).

- Persistence intervals: $\{[i, j) \mid i = \text{PivotIndex } R_j\} \cup \{[i, \infty) \mid i \in I_e\}$

Compatible basis cycles

For a reduced boundary matrix $R = D \cdot V$, call

$$I_b = \{i : R_i = 0\}$$

birth indices,

$$I_d = \{j : R_j \neq 0\}$$

death indices,


$$I_e = I_b \setminus \text{PivotIndices } R$$

essential indices.

(note: all pivot indices are birth indices).

- Persistence intervals: $\{[i, j) \mid i = \text{PivotIndex } R_j\} \cup \{[i, \infty) \mid i \in I_e\}$
- Columns with non-essential birth indices ($\text{PivotIndices } R$) are not used at all!

Matrix reduction



	1	2	3	4	5	6	7
1			1		1		
2			1			1	
3							1
4				1	1		
5						1	
6							1
7							


$\underbrace{\hspace{10em}}_R$

$$= D \cdot \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{matrix} \\ \begin{matrix} 1 & 1 & & & & & \\ 2 & & 1 & & & & \\ 3 & & & 1 & & & \\ 4 & & & & 1 & & \\ 5 & & & & & 1 & \\ 6 & & & & & & 1 \\ 7 & & & & & & \end{matrix} \\ \underbrace{\hspace{10em}}_V \end{matrix}$$

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



	1	2	3	4	5	6	7
1			1	1			
2					1		
3							1
4				1	1		
5						1	
6							1
7							

$\underbrace{\hspace{10em}}$ _R

$= D \cdot$


	1	2	3	4	5	6	7
1	1						
2		1					
3			1				
4				1			
5					1		
6						1	
7							1

$\underbrace{\hspace{10em}}$ _V

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



	1	2	3	4	5	6	7
1			1		1		
2			1			1	
3							1
4				1	1		
5						1	
6						1	
7							

$\underbrace{\hspace{10em}}$ _R

$= D \cdot$


	1	2	3	4	5	6	7
1	1						
2		1					
3			1				
4				1			
5					1		
6						1	
7							1

$\underbrace{\hspace{10em}}$ _V

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



	1	2	3	4	5	6	7
1			1	1	1	1	
2			1		1		
3							1
4				1	0		
5						1	
6						1	
7							

$\underbrace{\hspace{10em}}_R$

$= D \cdot$


	1	2	3	4	5	6	7
1	1						
2		1					
3			1				
4				1			
5					1	1	
6						1	
7							1

$\underbrace{\hspace{10em}}_V$

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction




$$\begin{array}{c} \begin{array}{|c|c|c|c|c|c|c|c|} \hline & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \hline 1 & & & 1 & 1 & 1 & 1 \\ \hline 2 & & & & & & 1 \\ \hline 3 & & & & & & & 1 \\ \hline 4 & & & & & 1 & & \\ \hline 5 & & & & & & & 1 \\ \hline 6 & & & & & & & 1 \\ \hline 7 & & & & & & & \\ \hline \end{array} \\ \underbrace{\hspace{10em}}_R \end{array} = D \cdot \begin{array}{c} \begin{array}{|c|c|c|c|c|c|c|c|} \hline & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \hline 1 & 1 & & & & & & \\ \hline 2 & & 1 & & & & & \\ \hline 3 & & & 1 & & & & \\ \hline 4 & & & & 1 & & & \\ \hline 5 & & & & & 1 & 1 & \\ \hline 6 & & & & & & 1 & \\ \hline 7 & & & & & & & 1 \\ \hline \end{array} \\ \underbrace{\hspace{10em}}_V \end{array}$$

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



	1	2	3	4	5	6	7
1			1	1	1	0	
2			1			0	
3							1
4				1			
5						1	
6						1	
7						1	

$\underbrace{\hspace{10em}}$
 R

$= D \cdot$


	1	2	3	4	5	6	7
1	1						
2		1					
3			1				1
4				1			
5					1	1	
6						1	
7							1

$\underbrace{\hspace{10em}}$
 V

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction




$$\begin{array}{c} \begin{array}{|c|c|c|c|c|c|c|c|} \hline & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \hline 1 & & & 1 & & 1 & & \\ \hline 2 & & & 1 & & & & \\ \hline 3 & & & & & & 1 & \\ \hline 4 & & & & & 1 & & \\ \hline 5 & & & & & & 1 & \\ \hline 6 & & & & & & 1 & \\ \hline 7 & & & & & & 1 & \\ \hline \end{array} \\ \underbrace{\hspace{10em}}_R \end{array} = D \cdot \begin{array}{c} \begin{array}{|c|c|c|c|c|c|c|c|} \hline & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \hline 1 & 1 & & & & & & \\ \hline 2 & & 1 & & & & & \\ \hline 3 & & & 1 & & & 1 & \\ \hline 4 & & & & 1 & & & \\ \hline 5 & & & & & 1 & 1 & \\ \hline 6 & & & & & & 1 & \\ \hline 7 & & & & & & & 1 \\ \hline \end{array} \\ \underbrace{\hspace{10em}}_V \end{array}$$

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



	1	2	3	4	5	6	7
1			1		1		
2			1				
3							1
4				1			
5						1	
6						1	
7							

$\underbrace{\hspace{10em}}$ _R

$= D \cdot$


	1	2	3	4	5	6	7
1	1						
2		1					
3			1			1	
4				1			
5					1	1	
6						1	
7							1

$\underbrace{\hspace{10em}}$ _V

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



	1	2	3	4	5	6	7
1			1		1		
2			1				
3						1	
4					1		
5						1	
6						1	
7							

$\underbrace{\hspace{10em}}$ _R

$= D \cdot$


	1	2	3	4	5	6	7
1	1						
2		1					
3			1			1	
4				1			
5					1	1	
6						1	
7							1

$\underbrace{\hspace{10em}}$ _V

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



	1	2	3	4	5	6	7
1			1		1		
2			1				
3						1	
4					1		
5						1	
6						1	
7							

$\underbrace{\hspace{10em}}$
 R

$= D \cdot$


	1	2	3	4	5	6	7
1	1						
2		1					
3			1			1	
4				1			
5					1	1	
6						1	
7							1

$\underbrace{\hspace{10em}}$
 V

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Matrix reduction



$$\begin{array}{c} \begin{array}{|c|c|c|c|c|c|c|c|} \hline & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \hline 1 & & & 1 & & 1 & & \\ \hline 2 & & & & 1 & & & \\ \hline 3 & & & & & & & 1 \\ \hline 4 & & & & & 1 & & \\ \hline 5 & & & & & & & 1 \\ \hline 6 & & & & & & & 1 \\ \hline 7 & & & & & & & \\ \hline \end{array} \\ \underbrace{\hspace{10em}}_R \end{array} = D \cdot \begin{array}{c} \begin{array}{|c|c|c|c|c|c|c|} \hline & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \hline 1 & 1 & & & & & & \\ \hline 2 & & 1 & & & & & \\ \hline 3 & & & 1 & & & & 1 \\ \hline 4 & & & & 1 & & & \\ \hline 5 & & & & & 1 & & 1 \\ \hline 6 & & & & & & 1 & \\ \hline 7 & & & & & & & 1 \\ \hline \end{array} \\ \underbrace{\hspace{10em}}_V \end{array}$$

Algorithm (over $\mathbb{Z}/2\mathbb{Z}$):

- while $\exists k < j$ with PivotIndex $R_k = \text{PivotIndex } R_j$
 - add R_k to R_j , add V_k to V_j

Clearing

Clearing non-essential positive columns

Recall:

- Columns with indices PivotIndices R (non-essential birth indices) not used at all

Idea [Chen, Kerber 2011]:

- Don't reduce those columns
- Instead, *clear* them to 0
 - Use the fact that every pivot index is a birth index: $i = \text{PivotIndex } R_j$ implies $R_i = 0$
- To ensure that column i is cleared before it is reduced:
 - Reduce boundary matrices individually, in decreasing dimension

Clearing non-essential positive columns

Recall:

- Columns with indices PivotIndices R (non-essential birth indices) not used at all

Idea [Chen, Kerber 2011]:

- Don't reduce those columns
- Instead, *clear* them to 0
 - Use the fact that every pivot index is a birth index: $i = \text{PivotIndex } R_j$ implies $R_i = 0$
- To ensure that column i is cleared before it is reduced:
 - Reduce boundary matrices individually, in decreasing dimension

Note:

- reducing *birth* columns typically harder than death columns:
 - $O(j^2)$ (birth) vs. $O((j - i)^2)$ (death)
- with clearing: need only reduce *essential* columns

Counting homology column reductions

Consider $k + 1$ -skeleton of $n - 1$ -simplex (Rips filtration)

Number of columns in boundary matrix:

$$\sum_{d=1}^{k+1} \underbrace{\binom{n}{d+1}}_{\text{total}} = \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d}}_{\text{death}} + \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d+1}}_{\text{birth}}$$

Counting homology column reductions

Consider $k + 1$ -skeleton of $n - 1$ -simplex (Rips filtration)

Number of columns in boundary matrix:

$$\begin{aligned} \sum_{d=1}^{k+1} \underbrace{\binom{n}{d+1}}_{\text{total}} &= \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d}}_{\text{death}} + \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d+1}}_{\text{birth}} \\ &= \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d}}_{\text{death}} + \underbrace{\binom{n-1}{k+2}}_{\text{essential}} + \sum_{d=1}^k \underbrace{\binom{n-1}{d+1}}_{\text{cleared}} \end{aligned}$$

Counting homology column reductions

Consider $k + 1$ -skeleton of $n - 1$ -simplex (Rips filtration)

Number of columns in boundary matrix:

$$\begin{aligned} \sum_{d=1}^{k+1} \underbrace{\binom{n}{d+1}}_{\text{total}} &= \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d}}_{\text{death}} + \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d+1}}_{\text{birth}} \\ &= \sum_{d=1}^{k+1} \underbrace{\binom{n-1}{d}}_{\text{death}} + \underbrace{\binom{n-1}{k+2}}_{\text{essential}} + \sum_{d=1}^k \underbrace{\binom{n-1}{d+1}}_{\text{cleared}} \end{aligned}$$

$$k = 2, n = 192: \quad 56\,050\,096 = 1161\,471 + \cancel{53\,727\,345} + \cancel{1161\,280}$$

- Clearing didn't help much!

Cohomology

Cohomology and clearing

The persistence barcode can also be computed using *cohomology* (the vector space dual of homology)

- Persistent homology and persistent cohomology have the same barcode.

Cohomology and clearing

The persistence barcode can also be computed using *cohomology* (the vector space dual of homology)

- Persistent homology and persistent cohomology have the same barcode.
- Persistent cohomology can be computed by reducing the transposed boundary matrix.

Cohomology and clearing

The persistence barcode can also be computed using *cohomology* (the vector space dual of homology)

- Persistent homology and persistent cohomology have the same barcode.
- Persistent cohomology can be computed by reducing the transposed boundary matrix.
- Cohomology allows for clearing starting from dimension 0

Cohomology and clearing

The persistence barcode can also be computed using *cohomology* (the vector space dual of homology)

- Persistent homology and persistent cohomology have the same barcode.
- Persistent cohomology can be computed by reducing the transposed boundary matrix.
- Cohomology allows for clearing starting from dimension 0
- Persistence in dimension 0 has special algorithms
 - Kruskal's algorithm for minimum spanning tree
 - union-find data structure

Counting cohomology column reductions

Consider K : $k + 1$ -skeleton of $n - 1$ -simplex, Rips filtration

- Number of columns in coboundary matrix:

$$\sum_{d=0}^k \underbrace{\binom{n}{d+1}}_{\text{total}} = \sum_{d=0}^k \underbrace{\binom{n-1}{d+1}}_{\text{death}} + \sum_{d=0}^k \underbrace{\binom{n-1}{d}}_{\text{birth}}$$

Counting cohomology column reductions

Consider K : $k + 1$ -skeleton of $n - 1$ -simplex, Rips filtration

- Number of columns in coboundary matrix:

$$\begin{aligned} \sum_{d=0}^k \underbrace{\binom{n}{d+1}}_{\text{total}} &= \sum_{d=0}^k \underbrace{\binom{n-1}{d+1}}_{\text{death}} + \sum_{d=0}^k \underbrace{\binom{n-1}{d}}_{\text{birth}} \\ &= \sum_{d=0}^k \underbrace{\binom{n-1}{d+1}}_{\text{death}} + \underbrace{\binom{n-1}{0}}_{\text{essential}} + \sum_{d=1}^k \underbrace{\binom{n-1}{d}}_{\text{cleared}} \end{aligned}$$

Counting cohomology column reductions

Consider K : $k + 1$ -skeleton of $n - 1$ -simplex, Rips filtration

- Number of columns in coboundary matrix:

$$\begin{aligned} \sum_{d=0}^k \underbrace{\binom{n}{d+1}}_{\text{total}} &= \sum_{d=0}^k \underbrace{\binom{n-1}{d+1}}_{\text{death}} + \sum_{d=0}^k \underbrace{\binom{n-1}{d}}_{\text{birth}} \\ &= \sum_{d=0}^k \underbrace{\binom{n-1}{d+1}}_{\text{death}} + \underbrace{\binom{n-1}{0}}_{\text{essential}} + \sum_{d=1}^k \underbrace{\binom{n-1}{d}}_{\text{cleared}} \end{aligned}$$

- $k = 2, n = 192: 1179\,808 = 1161\,471 + 1 + 18\,336$

Observations

For a typical input:

- V has very few off-diagonal entries
- most death columns of D are already reduced

Observations

For a typical input:

- V has very few off-diagonal entries
- most death columns of D are already reduced

Previous example ($k = 2, n = 192$):

- Only $79 + 42 + 1 = 122$ out of $192 + 18\,145 + 1\,143\,135 = 1161\,471$ columns are actually modified in matrix reduction

Implicit matrix reduction

Implicit matrix reduction

Standard approach:

- Boundary matrix D for filtration-ordered basis
 - Explicitly generated and stored in memory

Implicit matrix reduction

Standard approach:

- Boundary matrix D for filtration-ordered basis
 - Explicitly generated and stored in memory
- Matrix reduction: store only reduced matrix R
 - Initialize matrix as D , reduce to R by column operations

Implicit matrix reduction

Standard approach:

- Boundary matrix D for filtration-ordered basis
 - Explicitly generated and stored in memory
- Matrix reduction: store only reduced matrix R
 - Initialize matrix as D , reduce to R by column operations

Approach for Ripser:

- Boundary matrix D for lexicographically ordered basis
 - Columns recomputed on the fly

Implicit matrix reduction

Standard approach:

- Boundary matrix D for filtration-ordered basis
 - Explicitly generated and stored in memory
- Matrix reduction: store only reduced matrix R
 - Initialize matrix as D , reduce to R by column operations

Approach for Ripser:

- Boundary matrix D for lexicographically ordered basis
 - Columns recomputed on the fly
- Matrix reduction: store only coefficient matrix V
 - Columns of $R_j = D \cdot V_j$ recomputed on the fly
 - Typically, V is much sparser and smaller than R

Apparent pairs

Simplexwise filtrations

Call a filtration $K_\bullet = (K_i)_i$ of simplicial complexes *simplexwise* if in the filtration, the simplices enter one at a time.

Simplexwise filtrations

Call a filtration $K_\bullet = (K_i)_i$ of simplicial complexes *simplexwise* if in the filtration, the simplices enter one at a time.

Note:

- This is required for computation.
- The *Vietoris–Rips filtration* (indexed by \mathbb{R}) is not simplexwise: several simplices appear simultaneously.
- To compute Vietoris–Rips persistence, we have to break ties.
- This creates (many) persistence pairs with persistence 0 (as an artifact of tie-breaking).

Apparent pairs

Definition

In a simplexwise filtration, a pair of simplices (σ, τ) is an *apparent* pair if

- σ is the youngest facet of τ , and
- τ is the oldest cofacet of σ .

Apparent pairs

Definition

In a simplexwise filtration, a pair of simplices (σ, τ) is an *apparent* pair if

- σ is the youngest facet of τ , and
- τ is the oldest cofacet of σ .

Proposition

The apparent pairs are persistence pairs. Their corresponding columns in the boundary matrix are already reduced.

Apparent pairs

Definition

In a simplexwise filtration, a pair of simplices (σ, τ) is an *apparent* pair if


- σ is the youngest facet of τ , and
- τ is the oldest cofacet of σ .

Proposition

The apparent pairs are persistence pairs. Their corresponding columns in the boundary matrix are already reduced.

Theorem

For the Vietoris–Rips filtration, in dimension 1, the 0-persistence pairs are the apparent pairs.



Ripser Live users from 770 locations

<http://ulrich-bauer.org/complexes-skoltech.pdf>



Ulrich Bauer.

Ripser: efficient computation of Vietoris–Rips persistence barcodes.

Submitted, Aug. 2019. arXiv:1908.02518

Software available at <http://ripser.org>.



Ulrich Bauer and Herbert Edelsbrunner.

The Morse theory of Čech and Delaunay complexes.

Transactions of the American Mathematical Society, 369(5):3741–3762, 2017.



Ulrich Bauer and Michael Lesnick.

Induced matchings and the algebraic stability of persistence barcodes.

J. Comput. Geom., 6(2):162–191, 2015.



Jan Reininghaus, Stefan Huber, Ulrich Bauer, and Roland Kwitt.

A Stable Multi-Scale Kernel for Topological Machine Learning.

In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2015.