

# Homological reconstruction and simplification in $\mathbb{R}^3$

*Ulrich Bauer<sup>1</sup> Dominique Attali<sup>2</sup> Olivier Devillers<sup>3</sup>  
Marc Glisse<sup>3</sup> André Lieutier<sup>4</sup>*

<sup>1</sup>IST Austria

<sup>2</sup>GIPSA-lab

<sup>3</sup>GEOMETRICA (INRIA)


<sup>4</sup>Dassault Systèmes


June 18, 2013


Symposium on Computational Geometry 2013




















- ▶ Homology transforms topological spaces to groups



- ▶ Homology transforms topological spaces to groups  
... and continuous maps into group homomorphisms




- ▶ Homology transforms topological spaces to groups  
... and continuous maps into group homomorphisms
- ▶ *Persistent homology group:*  $\text{im } H_*(\text{yellow} \hookrightarrow \text{blue})$





- ▶ Homology transforms topological spaces to groups  
... and continuous maps into group homomorphisms
- ▶ *Persistent homology group:*  $\text{im } H_*(\text{yellow} \hookrightarrow \text{blue})$
- ▶ Under mild sampling conditions on :


$$\text{im } H_*(\text{yellow} \hookrightarrow \text{blue}) \sim H_*(\text{doughnut})$$


[Edelsbrunner et al. 2005, Chazal et al. 2005]











# Homological reconstruction

(Throughout this talk: homology with field coefficients)

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological reconstruction* of  $(K, L)$ .

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)

*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological reconstruction* of  $(K, L)$ .

- ▶ If  $L \subset Y \subset K$  then  $\beta(Y) \geq \text{rank } H_*(L \hookrightarrow K)$

# Homological reconstruction

(Throughout this talk: homology with field coefficients)

## Problem (Homological reconstruction)


*Given a simplicial pair  $(K, L)$ , find  $X$  with  $L \subset X \subset K$  such that  $H_*(L \hookrightarrow X)$  is surjective and  $H_*(X \hookrightarrow K)$  is injective.*

Such an  $X$  is called a *homological reconstruction* of  $(K, L)$ .

- ▶ If  $L \subset Y \subset K$  then  $\beta(Y) \geq \text{rank } H_*(L \hookrightarrow K)$
- ▶ Equality iff  $Y$  is a homological reconstruction


# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist




# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist




# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist




# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist




# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist




- ▶ A homological reconstruction may exist, but not as a subcomplex of  $K$




# Do homological reconstructions always exist?

- ▶ Homological reconstructions do not always exist



- ▶ A homological reconstruction may exist, but not as a subcomplex of  $K$



# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

## Theorem

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

## Theorem

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

Outline of proof: reduction from 3-SAT

- ▶ Given a formula  $\Phi$  in 3-CNF, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

## Theorem

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

Outline of proof: reduction from 3-SAT

- ▶ Given a formula  $\Phi$  in 3-CNF, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

## Theorem

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

Outline of proof: reduction from 3-SAT

- ▶ Given a formula  $\Phi$  in 3-CNF, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

## Theorem

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

Outline of proof: reduction from 3-SAT

- ▶ Given a formula  $\Phi$  in 3-CNF, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard

## Theorem

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

Outline of proof: reduction from 3-SAT

- ▶ Given a formula  $\Phi$  in 3-CNF, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment
  - (b)  $(K, L)$  has a reconstruction

# Homological reconstruction in $\mathbb{R}^3$ is NP-hard


## Theorem

*Let  $(K, L)$  be a simplicial pair in  $\mathbb{R}^3$ . Deciding whether  $(K, L)$  has a homological reconstruction is NP-hard.*

Outline of proof: reduction from 3-SAT


- ▶ Given a formula  $\Phi$  in 3-CNF, construct a simplicial pair  $(K, L)$  with trivial persistent homology group  $H_*(L \hookrightarrow K)$
- ▶ Reconstruction:  $X$  with trivial homology and  $L \subset X \subset K$
- ▶ For this construction, the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment
  - (b)  $(K, L)$  has a reconstruction
  - (c)  $(K, L)$  has a reconstruction as a subcomplex of  $K$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  can contain at most one of the edges  $\text{True}_i, \text{False}_i$

# Reduction from 3-SAT: the variable gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  can contain at most one of the edges  $\text{True}_i, \text{False}_i$
- ▶ edges  $\text{True}_i, \text{False}_i$  correspond to possible values of variable  $x_i$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget




- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$

# Reduction from 3-SAT: the clause gadget





- ▶ red: contained in  $L$ , blue:  $K \setminus L$
- ▶  $X$  must contain one of the edges  $a, b, c$
- ▶ edges  $a, b, c$  correspond to literals in a clause

# Reduction from 3-SAT: an example




$(t \vee \neg v \vee \neg w)$




- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals

# Reduction from 3-SAT: an example




- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals

# Reduction from 3-SAT: an example




- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals

# Reduction from 3-SAT: an example





- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals
- ▶ We can transform a satisfying assignment into a homological reconstruction of  $(K, L)$  and vice versa

# Reduction from 3-SAT: an example



- ▶ For every clause: identify edges  $a, b, c$  of clause gadget with edges of variable gadgets corresponding to the literals
- ▶ We can transform a satisfying assignment into a homological reconstruction of  $(K, L)$  and vice versa





# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .
- ▶ For any  $X$  with  $L \subset X \subset K$ , there is  $g$  with  $G_{\leq t} = X$ .

# Sublevel set simplification

Let  $F_{\leq t} = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

## Problem (Sublevel set simplification)

*Given a PL function  $f : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$   
minimizing  $\beta(G_{\leq t}) = \text{rank } H_*(G_{\leq t})$ .*

Let  $K = F_{\leq t+\delta}$ ,  $L = F_{\leq t-\delta}$ .

- ▶ For any  $g$ , we have  $L \subset G_{\leq t} \subset K$ .
- ▶ For any  $X$  with  $L \subset X \subset K$ , there is  $g$  with  $G_{\leq t} = X$ .

Thus we are looking for  $X$  with  $L \subset X \subset K$  minimizing  $\beta(X)$ .

- ▶ Lower bound  $\beta(G_{\leq t}) \geq \text{rank } H_*(L \hookrightarrow K)$
- ▶ *Sublevel set reconstruction:* a function  $g$  achieving the lower bound

## Theorem

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

## Theorem

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Outline of proof:

- ▶ Construct  $(K, L)$  from  $\Phi$  as before.

Recall that the following are equivalent:

- (a)  $\Phi$  has a satisfying assignment
- (b)  $(K, L)$  has a reconstruction
- (c)  $(K, L)$  has a reconstruction as a subcomplex  $X \subset K$

## Theorem

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Outline of proof:

- ▶ Construct  $(K, L)$  from  $\Phi$  as before.

Recall that the following are equivalent:

- (a)  $\Phi$  has a satisfying assignment
- (b)  $(K, L)$  has a reconstruction
- (c)  $(K, L)$  has a reconstruction as a subcomplex  $X \subset K$

- ▶ Construct a function  $f$  with  $F_{\leq t+\delta} = K$  and  $F_{\leq t-\delta} = L$ . The following are equivalent to the above:

## Theorem

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Outline of proof:

- ▶ Construct  $(K, L)$  from  $\Phi$  as before.

Recall that the following are equivalent:

- (a)  $\Phi$  has a satisfying assignment
- (b)  $(K, L)$  has a reconstruction
- (c)  $(K, L)$  has a reconstruction as a subcomplex  $X \subset K$

- ▶ Construct a function  $f$  with  $F_{\leq t+\delta} = K$  and  $F_{\leq t-\delta} = L$ . The following are equivalent to the above:

- (d)  $f$  has a simplexwise linear sublevel set reconstruction  $g$

## Theorem

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Outline of proof:

- ▶ Construct  $(K, L)$  from  $\Phi$  as before.

Recall that the following are equivalent:

- (a)  $\Phi$  has a satisfying assignment
- (b)  $(K, L)$  has a reconstruction
- (c)  $(K, L)$  has a reconstruction as a subcomplex  $X \subset K$

- ▶ Construct a function  $f$  with  $F_{\leq t+\delta} = K$  and  $F_{\leq t-\delta} = L$ . The following are equivalent to the above:

- (d)  $f$  has a simplexwise linear sublevel set reconstruction  $g$
- (e)  $f$  has a sublevel set reconstruction  $g$

## Theorem

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Outline of proof:

- ▶ Construct  $(K, L)$  from  $\Phi$  as before.

Recall that the following are equivalent:

- (a)  $\Phi$  has a satisfying assignment
- (b)  $(K, L)$  has a reconstruction
- (c)  $(K, L)$  has a reconstruction as a subcomplex  $X \subset K$

- ▶ Construct a function  $f$  with  $F_{\leq t+\delta} = K$  and  $F_{\leq t-\delta} = L$ . The following are equivalent to the above:

- (d)  $f$  has a simplexwise linear sublevel set reconstruction  $g$
- (e)  $f$  has a sublevel set reconstruction  $g$

- ▶ (c)  $\implies$  (d): construct  $g$  with  $G_{\leq t} = X$

## Theorem

*Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.*

Outline of proof:

- ▶ Construct  $(K, L)$  from  $\Phi$  as before.

Recall that the following are equivalent:

- (a)  $\Phi$  has a satisfying assignment
- (b)  $(K, L)$  has a reconstruction
- (c)  $(K, L)$  has a reconstruction as a subcomplex  $X \subset K$

- ▶ Construct a function  $f$  with  $F_{\leq t+\delta} = K$  and  $F_{\leq t-\delta} = L$ . The following are equivalent to the above:

- (d)  $f$  has a simplexwise linear sublevel set reconstruction  $g$
- (e)  $f$  has a sublevel set reconstruction  $g$

- ▶ (c)  $\implies$  (d): construct  $g$  with  $G_{\leq t} = X$
- ▶ (d)  $\implies$  (e)  $\implies$  (b): trivial

# Level set simplification

Let  $F_t = f^{-1}(t)$  denote the  $t$ -level set of  $f$ .

## Problem (Level set simplification)

*Given a PL function  $f : \mathbb{S}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\beta(G_t)$ .*

# Level set simplification

Let  $F_t = f^{-1}(t)$  denote the  $t$ -level set of  $f$ .

## Problem (Level set simplification)

*Given a PL function  $f : \mathbb{S}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\beta(G_t)$ .*

- ▶  $\beta(G_t) = \beta(G_{\leq t}) + \beta(G_{<t})$   
(Mayer-Vietoris, Alexander duality)

# Level set simplification

Let  $F_t = f^{-1}(t)$  denote the  $t$ -level set of  $f$ .

## Problem (Level set simplification)

Given a PL function  $f : \mathbb{S}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\beta(G_t)$ .

- ▶  $\beta(G_t) = \beta(G_{\leq t}) + \beta(G_{<t})$   
(Mayer-Vietoris, Alexander duality)
- ▶  $\beta(G_t) \geq \text{rank } H_*(F_{\leq t-\delta} \hookrightarrow F_{\leq t+\delta}) + \text{rank } H_*(F_{<t-\delta} \hookrightarrow F_{<t+\delta})$

# Level set simplification

Let  $F_t = f^{-1}(t)$  denote the  $t$ -level set of  $f$ .

## Problem (Level set simplification)

Given a PL function  $f : \mathbb{S}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,  
find a PL function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\beta(G_t)$ .

- ▶  $\beta(G_t) = \beta(G_{\leq t}) + \beta(G_{<t})$   
(Mayer-Vietoris, Alexander duality)
- ▶  $\beta(G_t) \geq \text{rank } H_*(F_{\leq t-\delta} \hookrightarrow F_{\leq t+\delta}) + \text{rank } H_*(F_{<t-\delta} \hookrightarrow F_{<t+\delta})$
- ▶ *Level set reconstruction:* a simplification  $g$  achieving the lower bound on  $\beta(G_t)$

## Theorem

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

## Theorem

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Outline of proof:

- ▶ Recall that the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment
  - (b,c)  $(K, L)$  has a reconstruction (as a subcomplex  $X \subset K$ )
  - (d,e)  $f$  has a (simplexwise linear) sublevel set reconstruction  $g$

## Theorem

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Outline of proof:

- ▶ Recall that the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment
  - (b,c)  $(K, L)$  has a reconstruction (as a subcomplex  $X \subset K$ )
  - (d,e)  $f$  has a (simplexwise linear) sublevel set reconstruction  $g$
- ▶ Extend  $f$  to  $\tilde{f} : \mathbb{S}^3 \rightarrow \mathbb{R}$ . Equivalent to the above:
  - (f)  $\tilde{f}$  has a simplexwise linear level set reconstruction  $\tilde{g}$
  - (g)  $\tilde{f}$  has a level set reconstruction  $\tilde{g}$

## Theorem

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Outline of proof:

- ▶ Recall that the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment
  - (b,c)  $(K, L)$  has a reconstruction (as a subcomplex  $X \subset K$ )
  - (d,e)  $f$  has a (simplexwise linear) sublevel set reconstruction  $g$
- ▶ Extend  $f$  to  $\tilde{f} : \mathbb{S}^3 \rightarrow \mathbb{R}$ . Equivalent to the above:
  - (f)  $\tilde{f}$  has a simplexwise linear level set reconstruction  $\tilde{g}$
  - (g)  $\tilde{f}$  has a level set reconstruction  $\tilde{g}$
- ▶ (g)  $\implies$  (e):  $\tilde{g}$  has regular value  $t$ , so  $\beta(\tilde{G}_t) = 2\beta(\tilde{G}_{\leq t})$ .  
Hence  $\tilde{g}$  also minimizes  $\beta(\tilde{G}_{\leq t})$

## Theorem

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Outline of proof:

- ▶ Recall that the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment
  - (b,c)  $(K, L)$  has a reconstruction (as a subcomplex  $X \subset K$ )
  - (d,e)  $f$  has a (simplexwise linear) sublevel set reconstruction  $g$
- ▶ Extend  $f$  to  $\tilde{f} : \mathbb{S}^3 \rightarrow \mathbb{R}$ . Equivalent to the above:
  - (f)  $\tilde{f}$  has a simplexwise linear level set reconstruction  $\tilde{g}$
  - (g)  $\tilde{f}$  has a level set reconstruction  $\tilde{g}$
- ▶ (g)  $\implies$  (e):  $\tilde{g}$  has regular value  $t$ , so  $\beta(\tilde{G}_t) = 2\beta(\tilde{G}_{\leq t})$ .  
Hence  $\tilde{g}$  also minimizes  $\beta(\tilde{G}_{\leq t})$
- ▶ (b)  $\implies$  (f): from  $X$ , construct  $\tilde{g}$  with  $G_{\leq t} = X$  and regular value  $t$ . Then  $\tilde{g}$  minimizes both  $\beta(\tilde{G}_{\leq t})$  and  $\beta(\tilde{G}_t) = 2\beta(\tilde{G}_{\leq t})$

## Theorem

*Level set simplification in  $\mathbb{S}^3$  is NP-hard.*

Outline of proof:

- ▶ Recall that the following are equivalent:
  - (a)  $\Phi$  has a satisfying assignment
  - (b,c)  $(K, L)$  has a reconstruction (as a subcomplex  $X \subset K$ )
  - (d,e)  $f$  has a (simplexwise linear) sublevel set reconstruction  $g$
- ▶ Extend  $f$  to  $\tilde{f} : \mathbb{S}^3 \rightarrow \mathbb{R}$ . Equivalent to the above:
  - (f)  $\tilde{f}$  has a simplexwise linear level set reconstruction  $\tilde{g}$
  - (g)  $\tilde{f}$  has a level set reconstruction  $\tilde{g}$
- ▶ (g)  $\implies$  (e):  $\tilde{g}$  has regular value  $t$ , so  $\beta(\tilde{G}_t) = 2\beta(\tilde{G}_{\leq t})$ .  
Hence  $\tilde{g}$  also minimizes  $\beta(\tilde{G}_{\leq t})$
- ▶ (b)  $\implies$  (f): from  $X$ , construct  $\tilde{g}$  with  $G_{\leq t} = X$  and regular value  $t$ . Then  $\tilde{g}$  minimizes both  $\beta(\tilde{G}_{\leq t})$  and  $\beta(\tilde{G}_t) = 2\beta(\tilde{G}_{\leq t})$
- ▶ (f)  $\implies$  (g): trivial

# Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a, b]$ .

## Definition (Well group)

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

# Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a, b]$ .

## Definition (Well group)

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

- ▶ Intuition: a  $\delta$ -robust version of level set homology  $H_*(F_t)$

# Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a, b]$ .

## Definition (Well group)

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

- ▶ Intuition: a  $\delta$ -robust version of level set homology  $H_*(F_t)$
- ▶  $\beta(G_t) \geq \text{rank } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]}) \geq \text{rank } W_*(f, t, \delta)$

# Well groups [Edelsbrunner, Morozov, Patel 2011]

Let  $f$  be a PL function and  $t \in \mathbb{R}$ ,  $\delta \geq 0$ . Let  $F_{[a,b]} = f^{-1}[a, b]$ .

## Definition (Well group)

The  $(t, \delta)$ -well group of  $f$  is

$$W_*(f, t, \delta) = \bigcap_{g: \|g-f\|_\infty \leq \delta} \text{im } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]})$$

- ▶ Intuition: a  $\delta$ -robust version of level set homology  $H_*(F_t)$
- ▶  $\beta(G_t) \geq \text{rank } H_*(G_t \hookrightarrow F_{[t-\delta, t+\delta]}) \geq \text{rank } W_*(f, t, \delta)$
- ▶ *Realization* of well group: a function  $g$  with  $\|g - f\|_\infty \leq \delta$  achieving the lower bound  $\beta(G_t) = \text{rank } W_*(f, t, \delta)$ .

# Well group realization is level set reconstruction

## Theorem

*Let  $f$  be a PL function on  $\mathbb{S}^n$  with  $t \pm \delta \in \text{int}(\text{im } f)$ .*

*A PL function  $g$  realizes the well group  $W_*(f, t, \delta)$  if and only if it is a  $(t, \delta)$ -level set reconstruction of  $f$ .*

# Well group realization is level set reconstruction

## Theorem

*Let  $f$  be a PL function on  $\mathbb{S}^n$  with  $t \pm \delta \in \text{int}(\text{im } f)$ .*

*A PL function  $g$  realizes the well group  $W_*(f, t, \delta)$  if and only if it is a  $(t, \delta)$ -level set reconstruction off.*

## Corollary

*Realization of a well group on  $\mathbb{S}^3$  is NP-hard.*

Thanks for your attention!

