

# Algebraic perspectives of Persistence

## The stability of persistence barcodes

Ulrich Bauer

TUM

April 24, 2017



||







||2




?































# What is persistent homology?





# What is persistent homology?





# What is persistent homology?





# What is persistent homology?





# What is persistent homology?



# What is persistent homology?



# What is persistent homology?



# What is persistent homology?

# What is persistent homology?

Persistent homology is the homology of a filtration

# What is persistent homology?

Persistent homology is the homology of a filtration

- ▶ A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - ▶  $\mathbf{R}$  is the poset category of  $(\mathbb{R}, \leq)$
  - ▶ A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - ▶ An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$

# What is persistent homology?

Persistent homology is the homology of a filtration

- ▶ A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - ▶  $\mathbf{R}$  is the poset category of  $(\mathbb{R}, \leq)$
  - ▶ A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - ▶ An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- ▶ Consider homology with coefficients in a field (often  $\mathbb{Z}_2$ )  
 $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$

# What is persistent homology?

Persistent homology is the homology of a filtration

- ▶ A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - ▶  $\mathbf{R}$  is the poset category of  $(\mathbb{R}, \leq)$
  - ▶ A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - ▶ An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- ▶ Consider homology with coefficients in a field (often  $\mathbb{Z}_2$ )  
 $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- ▶ Persistent homology is a diagram  $M : \mathbf{R} \rightarrow \mathbf{Vect}$   
*(persistence module)*

# Homology inference

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $X$  with  $H_*(X) \cong H_*(\Omega)$ .*

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $X$  with  $H_*(X) \cong H_*(\Omega)$ .*

Idea:

- ▶ approximate the shape by a thickening  $B_\delta(P)$  covering  $\Omega$

# Homology inference

Given: finite sample  $P \subset \Omega$  of unknown shape  $\Omega \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(\Omega)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $X$  with  $H_*(X) \cong H_*(\Omega)$ .*

Idea:

- ▶ approximate the shape by a thickening  $B_\delta(P)$  covering  $\Omega$

Requires strong assumptions:


# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $M$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset M$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- $\delta < \sqrt{3/20} \text{reach}(M)$ .

Then  $H_*(M) \cong H_*(B_{2\delta}(P))$ .





# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $\Omega \subset \mathbb{R}^d$ . Let  $P \subset \Omega$ ,  $\delta > 0$  be such that

- $B_\delta(P)$  covers  $\Omega$ , and
- the inclusions  $\Omega \hookrightarrow B_\delta(\Omega) \hookrightarrow B_{2\delta}(\Omega)$  preserve homology.

Then  $H_*(\Omega) \cong \text{im } H_*(B_\delta(P) \hookrightarrow B_{2\delta}(P))$ .



# Stability

# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)


*Let  $\|f - g\|_\infty = \delta$ . Then there exists a matching between the intervals of the persistence barcodes of  $f$  and  $g$  such that*

# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $\|f - g\|_\infty = \delta$ . Then there exists a matching between the intervals of the persistence barcodes of  $f$  and  $g$  such that

- matched intervals have endpoints within distance  $\leq \delta$ , and




# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $\|f - g\|_\infty = \delta$ . Then there exists a matching between the intervals of the persistence barcodes of  $f$  and  $g$  such that

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .




# Persistence and stability: the big picture

Data


point cloud

$P \subset \mathbb{R}^d$


# Persistence and stability: the big picture




# Persistence and stability: the big picture



# Persistence and stability: the big picture



# Persistence and stability: the big picture



# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  and  $G_t = g^{-1}(-\infty, t]$ .

# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  and  $G_t = g^{-1}(-\infty, t]$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  
 $\delta$ -interleaved:

$$\begin{array}{ccccccc} F_t & \hookrightarrow & F_{t+\delta} & \hookrightarrow & F_{t+2\delta} & \cdots & \rightarrow \\ \swarrow \searrow & & \swarrow \searrow & & \swarrow \searrow & & \\ G_t & \hookrightarrow & G_{t+\delta} & \hookrightarrow & G_{t+2\delta} & \cdots & \rightarrow \end{array}$$

# Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  and  $G_t = g^{-1}(-\infty, t]$ .


Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:

$$\begin{array}{ccccccc} F_t & \hookrightarrow & F_{t+\delta} & \hookrightarrow & F_{t+2\delta} & \cdots & \rightarrow \\ \swarrow \searrow & & \swarrow \searrow & & \swarrow \searrow & & \\ G_t & \hookrightarrow & G_{t+\delta} & \hookrightarrow & G_{t+2\delta} & \cdots & \rightarrow \end{array}$$


Applying homology (functor) preserves commutativity

- persistent homology of  $f, g$  yields  $\delta$ -interleaved persistence modules  $\mathbf{R} \rightarrow \mathbf{Vect}$


# Geometric interleavings




# Geometric interleavings




# Geometric interleavings




# Geometric interleavings



# Geometric interleavings



# Geometric interleavings



# Algebraic stability of persistence barcodes

**Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)**

*If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:*


- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .

# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .



# Barcodes as diagrams


# The matching category

A *matching*  $\sigma : S \nrightarrow T$  is a bijection  $S' \rightarrow T'$ , where  $S' \subseteq S$ ,  $T' \subseteq T$ .




# The matching category

A *matching*  $\sigma : S \not\rightarrow T$  is a bijection  $S' \rightarrow T'$ , where  $S' \subseteq S$ ,  $T' \subseteq T$ .



Composition of matchings  $\sigma : S \not\rightarrow T$  and  $\tau : T \not\rightarrow U$ :



# The matching category

A *matching*  $\sigma : S \not\rightarrow T$  is a bijection  $S' \rightarrow T'$ , where  $S' \subseteq S$ ,  $T' \subseteq T$ .



Composition of matchings  $\sigma : S \not\rightarrow T$  and  $\tau : T \not\rightarrow U$ :




Matchings form a category **Mch**

- objects: sets
- morphisms: matchings

# Barcodes as matching diagrams


We can regard a barcode  $B$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :



# Barcodes as matching diagrams

We can regard a barcode  $B$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :


- For each real number  $t$ , let  $B_t$  be those intervals of  $B$  that contain  $t$ , and



# Barcodes as matching diagrams

We can regard a barcode  $B$  as a functor  $\mathbf{R} \rightarrow \mathbf{Mch}$ :

- For each real number  $t$ , let  $B_t$  be those intervals of  $B$  that contain  $t$ , and
- for each  $s \leq t$ , define the matching  $B_s \rightarrow B_t$  to be the identity on  $B_s \cap B_t$ .



# Stability via functoriality?

$$\begin{array}{ccc} F_t & \xrightarrow{\hspace{2cm}} & F_{t+2\delta} \\ \searrow & \nearrow & \searrow \\ G_{t+\delta} & \xrightarrow{\hspace{2cm}} & G_{t+3\delta} \end{array}$$

# Stability via functoriality?

$$\begin{array}{ccc} H_*(F_t) & \longrightarrow & H_*(F_{t+2\delta}) \\ & \searrow & \nearrow \\ & H_*(G_{t+\delta}) & \longrightarrow H_*(G_{t+3\delta}) \end{array}$$

# Stability via functoriality?

$$\begin{array}{ccc} B(H_*(F_t)) & \rightarrow & B(H_*(F_{t+2\delta})) \\ \searrow & & \nearrow \\ & B(H_*(G_{t+\delta})) & \rightarrow B(H_*(G_{t+3\delta})) \end{array}$$

# Stability via functoriality?

$$\begin{array}{ccc} B(H_*(F_t)) & \rightarrow & B(H_*(F_{t+2\delta})) \\ & \searrow & \nearrow \\ & B(H_*(G_{t+\delta})) & \rightarrow B(H_*(G_{t+3\delta})) \end{array}$$



# Non-functoriality of the persistence barcode

Theorem (B, Lesnick 2014)

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}^R$  sending each persistence module to its barcode.*

# Non-functoriality of the persistence barcode

## Theorem (B, Lesnick 2014)

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}^R$  sending each persistence module to its barcode.*

## Proposition

*There exists no functor  $\mathbf{Vect} \rightarrow \mathbf{Mch}$  sending each vector space of dimension  $d$  to a set of cardinality  $d$ .*

# Non-functoriality of the persistence barcode

## Theorem (B, Lesnick 2014)

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}^R$  sending each persistence module to its barcode.*

## Proposition

*There exists no functor  $\mathbf{Vect} \rightarrow \mathbf{Mch}$  sending each vector space of dimension  $d$  to a set of cardinality  $d$ .*

- ▶ Such a functor would necessarily send a linear map of rank  $r$  to a matching of cardinality  $r$ .

# Non-functoriality of the persistence barcode

## Theorem (B, Lesnick 2014)

*There exists no functor  $\mathbf{Vect}^R \rightarrow \mathbf{Mch}^R$  sending each persistence module to its barcode.*

## Proposition

*There exists no functor  $\mathbf{Vect} \rightarrow \mathbf{Mch}$  sending each vector space of dimension  $d$  to a set of cardinality  $d$ .*

- ▶ Such a functor would necessarily send a linear map of rank  $r$  to a matching of cardinality  $r$ .
- ▶ In particular, there is no natural choice of basis for vector spaces