

INDECOMPOSABLES IN MULTI-PARAMETER PERSISTENCE

Ulrich Bauer (TUM)

Applied Topology Seminar, Oxford University

November 24, 2023

joint work with:

Magnus Botnan / Steffen Oppermann / Johan Steen / Luis Scoccola / Ben Flahar




CLUSTERING FUNCTIONS

X : finite set

Clustering function φ :

maps a metric $d : X \times X \rightarrow \mathbb{R}$ (distance matrix)
to a partition of X .





KLEINBERG's AXIOMS

Desirable properties

- scale invariance :

$$\varphi(d) = \varphi(t \cdot d)$$



KLEINBERG's AXIOMS



Desirable properties

- scale invariance :

$$\varphi(d) = \varphi(t \cdot d)$$

- richness :

every partition is obtained from some d .



KLEINBERG's AXIOMS

Desirable properties

- scale invariance :

$$\varphi(d) = \varphi(t \cdot d)$$

- richness :




every partition is obtained from some d .

- consistency :

decreasing d within clusters /

increasing d across clusters

does not change the result.



KLEINBERG'S IMPOSSIBILITY THEOREM

Thm [Kleinberg 2002] No clustering function satisfies

- scale-invariant,
- richness , and
- consistency .

KLEINBERG'S IMPOSSIBILITY THEOREM

Thm [Kleinberg 2002] No clustering function satisfies


- scale-invariant,
- richness , and
- consistency .

Motivates the use of a scale parameter
⇒ hierarchical clustering

CLUSTERING FROM CONNECTED COMPONENTS

proximity graph

- filter edges by proximity





CLUSTERING FROM CONNECTED COMPONENTS

proximity graph

- filter edges by proximity



Π_0 (connected components)



single-linkage
clustering

CLUSTERING FROM CONNECTED COMPONENTS

proximity graph

- filter edges by proximity





π_0 (connected components)




H_0 (homology in deg. 0 with coeffs in K)

$$H_0 = F \circ \pi_0$$



single-linkage
clustering



persistent homology


HIERARCHICAL CLUSTERING : EXISTENCE & UNIQUENESS

Then [Carlsson, Mémoli 2010] single-linkage clustering is the **unique** hierarchical clustering method satisfying
[... certain axioms similar to Kleinberg's].

But ...

CHAINING EFFECT

Single-linkage clustering is sensitive to outliers




→ not used much in practice!

2 - PARAMETER CLUSTERING

density - proximity graph


- filters points by density
- filters edges by proximity



2 - PARAMETER CLUSTERING

density - proximity graph

- filters points by density
- filters edges by proximity




Π_0 (connected components)


2 - PARAMETER CLUSTERING

density - proximity graph


- filters points by density
- filters edges by proximity



π_0 (connected components)



H_0 (homology in deg. 0
with coeffs in K)



TRICHOTOMY OF REPRESENTATION TYPES

Given a finite indexing poset P . $(\mathcal{U}$: algebra-closed)

What are the indecomposable diagrams with shape P ?

TRICHOTOMY OF REPRESENTATION TYPES

Given a finite indexing poset P . $(\mathcal{U}$: algebra. closed)

What are the indecomposable diagrams with shape P ?

3 cases (representation types): [Drozd 1977]

② A finite list.

finite type

TRICHOTOMY OF REPRESENTATION TYPES

Given a finite indexing poset P . $(\mathcal{U}$: algebra. closed)

What are the indecomposable diagrams with shape P ?

3 cases (representation types): [Drozd 1977]

Ⓐ A finite list.

finite type

Ⓑ 1-param. families.

tame

TRICHOTOMY OF REPRESENTATION TYPES

Given a finite indexing poset P . (\mathbb{K} : algebra · closed)

What are the indecomposable diagrams with shape P ?

3 cases (representation types): [Drozd 1977]

(a) A finite list.

finite type

(b) 1-param. families.



tame

(c) It's complicated.

wild

(as complicated as modules over any finite-dim. algebra;
including undecidable problems)

REPRESENTATION TYPES OF COMMUTATIVE GRIDS




- finite type
 - tame
 - wild
- } for $(m-1)(n-1)$ { < = > } 4

[Leszczyński '94,
Łukowroński '2000]


GRID DIAGRAMS FROM CLUSTERING

Consider again 2-parameter clustering (proximity / density)




GRID DIAGRAMS FROM CLUSTERING

Consider again 2-parameter clustering (proximity / density)




This yields diagrams of the form




Horizontal maps are surjective !

GRID DIAGRAMS FROM CLUSTERING

Consider again 2-parameter clustering (proximity / density)



This yields diagrams of the form




Horizontal maps are surjective !

Does this simplify the picture ?

EPIMORPHISMS

Lemma $\text{Rep}^{\rightarrow}(m, 2)$ is finite type.



$$\begin{array}{c} H_0 \\ \sim \end{array} \left(\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix} \right) \begin{array}{c} K^3 \xrightarrow{(1,1,1)} K \xrightarrow{(1,1)} K \\ \uparrow \quad \uparrow \\ K^2 \xrightarrow{(1,1)} K^2 \xrightarrow{(1,1)} K \end{array}$$

$$\cong \begin{array}{ccc} K & \xrightarrow{\quad} & K & \xrightarrow{\quad} & K \\ \uparrow & & \uparrow & & \uparrow \\ K & \xrightarrow{\quad} & K & \xrightarrow{\quad} & K \end{array} \oplus \begin{array}{ccc} K & \xrightarrow{\quad} & 0 & \xrightarrow{\quad} & 0 \\ \uparrow & & \uparrow & & \uparrow \\ K & \xrightarrow{\quad} & K & \xrightarrow{\quad} & 0 \end{array} \oplus \begin{array}{ccc} K & \xrightarrow{\quad} & 0 & \xrightarrow{\quad} & 0 \\ \uparrow & & \uparrow & & \uparrow \\ 0 & \xrightarrow{\quad} & 0 & \xrightarrow{\quad} & 0 \end{array}$$

- $\text{Rep}(m, n)$: all commutative dgms over $m \times n$ grid
- $\text{Rep}^{\rightarrow}(m, n)$: epis in horizontal direction
- $\text{Rep}^{\uparrow\rightarrow}(m, n)$: epis in both directions.

EPIC GRIDS & WILD THINGS

Thm [B, Botnan, Oppermann, Steen 20]

$$\begin{array}{ccc} \text{Rep}^{\xrightarrow{\dagger}}(m, n) & \sim & \text{Rep}^{\dagger}(m, n-1) \\ \} & & \} \text{ same representation type} \\ \text{Rep}^{\xrightarrow{\dagger}}(m-1, n) & \sim & \text{Rep}(m-1, n-1) \end{array}$$

Corollary $\text{Rep}^{\xrightarrow{\dagger}}(m, n)$ is

- finite type
 - tame
 - wild
- } for $(m-1)(n-2)$ { $\begin{matrix} < \\ = \\ > \end{matrix}$ } 4 .

BEHIND THE SCENES


equivalence of categories

$$\frac{\text{Rep}^{\rightarrow}(m, n)}{\text{Rep}^{\tilde{\rightarrow}}(m, n)} \simeq \text{Rep}(m, n-1)$$

additive quotient $\frac{A}{B}$:

identify morphisms in A
whose difference factors
through B

indecomposables are of the form



\Rightarrow finite type

THE INSTABILITY OF DECOMPOSITIONS

How useful are indecomposables for TDA?

THE INSTABILITY OF DECOMPOSITIONS

How useful are indecomposables for TDA?

Thm [B, Scoccola 22] For $n > 1$, among the finitely presented n -parameter persistence modules the indecomposables are dense in interleaving distance.

THE INSTABILITY OF DECOMPOSITIONS

How useful are indecomposables for TDA?

Thm [B, Scoccola 22] For $n > 1$, among the finitely presented n -parameter persistence modules the indecomposables are dense in interleaving distance.

For every $\epsilon > 0$, the ϵ -indecomposables

$$A \oplus B$$

\nearrow \nwarrow
indecomposable $d_I(B, 0) < \epsilon$

form an open & dense subset.

- ϵ -indecomposability is a generic property

THE INSTABILITY OF DECOMPOSITIONS

Proof.

(a) $\forall \varepsilon, M : R^{\text{rect}} \rightarrow \text{vect f.p.}, \varepsilon\text{-indecomposable}$

$\exists \delta :$

$d_I(M, N) < \delta \Rightarrow N \text{ is } \varepsilon\text{-indecomposable.}$

THE INSTABILITY OF DECOMPOSITIONS

Proof.

(a) $\forall \varepsilon, M : \mathbb{R}^n \rightarrow \text{vect f.p.}, \varepsilon\text{-indecomposable}$

$\exists \delta :$

$d_I(M, N) < \delta \Rightarrow N \text{ is } \varepsilon\text{-indecomposable.}$

(b) Indecomposables can be "tacked together" with an arbitrarily small change in interleaving distance.

THE IDEA OF TACKING INDECOMPOSABLES

$$\begin{array}{ccc} k & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$

\oplus

$$\begin{array}{ccc} 0 & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$

\oplus

$$\begin{array}{ccc} 0 & \rightarrow & 0 \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$

\equiv

$$\begin{array}{ccc} k & \rightarrow & k^2 & \rightarrow & k^3 \\ \uparrow & & \uparrow & & \uparrow \\ 0 & \rightarrow & k & \rightarrow & k^2 \\ \uparrow & & \uparrow & & \uparrow \\ 0 & \rightarrow & 0 & \rightarrow & k \end{array}$$

THE IDEA OF TACKING INDECOMPOSABLES

$$\begin{array}{ccc} k & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$

\oplus

$$\begin{array}{ccc} 0 & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$

\oplus


$$\begin{array}{ccc} 0 & \rightarrow & 0 \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$

$$\begin{array}{ccc} k & \rightarrow & k^2 \\ \uparrow & & \uparrow \\ 0 & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$




$$\begin{array}{ccc} k & \xrightarrow{(1)} & k^2 \\ \uparrow & \uparrow^{(0)} & \uparrow \\ 0 & \rightarrow & k \\ \uparrow & & \uparrow \\ 0 & \rightarrow & 0 \end{array}$$


THE TACKING GADGET



THE TACKING GADGET




THE TACKING GADGET




TACKLING INDECOMPOSABLES


(0.)




(1.)



(2.)



(3.)



$$d_I(A \oplus B, M) < \delta \quad \text{for } \delta > 0 \text{ arbitrary}$$

THIN-DECOMPOSABLES ARE NOWHERE DENSE

Can we work with classes of simpler indecomposables?
(e.g. thin : pointwise $\dim \leq 1$)


THIN-DECOMPOSABLES ARE NOWHERE DENSE

Can we work with classes of simpler indecomposables?
(e.g. thin : pointwise $\dim \leq 1$)

Theorem [B, Scoccola '23] Let \mathcal{F} be a class of
indecomposable ($n=2$)-parameter persistence modules
with pointwise dimension bounded by some constant.
Then the \mathcal{F} -decomposable persistence modules are
nowhere dense.


THE MAYER - VIETORIS PYRAMID

[Carlsson et al. 2009] $f: X \rightarrow \mathbb{R}$




THE MAYER - VIETORIS PYRAMID

[Carlsson et al. 2009] $f: X \rightarrow \mathbb{R}$




THE MAYER - VIETORIS PYRAMID

[Carlsson et al. 2009] $f: X \rightarrow \mathbb{R}$




THE MAYER - VIETORIS PYRAMID

[Carlsson et al. 2009] $f: X \rightarrow \mathbb{R}$




THE MAYER - VIETORIS PYRAMID

[Carlsson et al. 2009] $f: X \rightarrow \mathbb{R}$




THE MAYER - VIETORIS PYRAMID

[Carlsson et al. 2009] $f: X \rightarrow \mathbb{R}$



THE MAYER - VIETORIS PYRAMID

[Carlsson et al. 2009] $f: X \rightarrow \mathbb{R}$



RISC OF A FUNCTION

[B, Bothan, Fluhr 2021] Assume $h(f)$ p.f.d.

Prop. $h(f)$ is a 2-parameter persistence module supported on a strip $M \subseteq \mathbb{R}^2$.

RISC OF A FUNCTION

[B, Bothan, Fluhr 2021] Assume $h(f)$ p.f.d.

Prop. $h(f)$ is a 2-parameter persistence module supported on a strip $M \subseteq \mathbb{R}^2$.

Prop. $h(f)$ is cohomological
(equivalently: middle-exact:

$$\begin{array}{ccc} A & \rightarrow & B \\ \downarrow & & \downarrow \\ C & \rightarrow & D \end{array}$$

$$\rightsquigarrow A \rightarrow B \oplus C \rightarrow D \text{ exact})$$

RISC OF A FUNCTION

[B, Bothan, Fluhr 2021] Assume $h(f)$ p.f.d.

Prop. $h(f)$ is a 2-parameter persistence module supported on a strip $M \subseteq \mathbb{R}^2$.


Prop. $h(f)$ is cohomological
(equivalently: middle-exact:

$$\begin{array}{ccc} A & \rightarrow & B \\ \downarrow & & \downarrow \\ C & \rightarrow & D \end{array} \quad \rightsquigarrow \quad A \rightarrow B \oplus C \rightarrow D \text{ exact})$$

Prop. $h(f)$ is sequentially continuous
(for sequences moving up/left in M)


DECOMPOSITION OF COHOMOLOGICAL FUNCTORS

Theorem [BBF '21] Any cohomological sequentially continuous p.f.d. functor $M \rightarrow \text{vect}$ decomposes into rectangle summands.



DECOMPOSITION OF COHOMOLOGICAL FUNCTORS


Theorem [BBF '21] Any cohomological sequentially continuous p.f.d. functor $M \rightarrow \text{vect}$ decomposes into rectangle summands.



(see also : [Botnan, Lebarici, Ondot '20])

DECOMPOSITION OF COHOMOLOGICAL FUNCTORS

Theorem [BBF '21] Any cohomological sequentially continuous p.f.d. functor $M \rightarrow \text{vect}$ decomposes into rectangle summands.



(see also : [Botnan, Lebarici, Ondot '20])

Proposition $M \rightarrow \text{vect}$ q-tame &
sequentially continuous
 \Rightarrow p.f.d.

INDUCED MORPHISMS & INTERLEAVINGS

- A map

$$\varphi : \begin{array}{ccc} X & \longrightarrow & Y \\ f \searrow & & \swarrow g \\ & R & \end{array} \quad \text{induces}$$

a morphism in RISC

$$h(\varphi) : h(f) \rightarrow h(g).$$

INDUCED MORPHISMS & INTERLEAVINGS

- A map

$$\varphi : \begin{array}{ccc} X & \longrightarrow & Y \\ f \searrow & & \swarrow g \\ & R & \end{array} \quad \text{induces}$$

a morphism in RISC

$$h(\varphi) : h(f) \rightarrow h(g).$$

- Two functions $f, g : X \rightarrow R$ with $\|f - g\|_\infty = \delta$ induce a δ -interleaving between $h(f)$ and $h(g)$.

INDUCED MORPHISMS & INTERLEAVINGS

- A map


$$\varphi: \begin{array}{ccc} X & \longrightarrow & Y \\ f \searrow & & \swarrow g \\ & R & \end{array} \quad \text{induces}$$

a morphism in RISC


$$h(\varphi): h(f) \rightarrow h(g).$$

- Two functions $f, g: X \rightarrow \mathbb{R}$ with $\|f - g\|_\infty = \delta$ induce a δ -interleaving between $h(f)$ and $h(g)$.
- These induced morphisms are richer than those in standard persistent homology!


WHAT STANDARD PERSISTENCE CAN'T SEE (BUT R1SC CAN)



WHAT STANDARD PERSISTENCE CAN'T SEE (BUT R1SC CAN)





WHAT STANDARD PERSISTENCE CAN'T SEE (BUT R1SC CAN)



- f has persistent reduced homology only in degree 0,
 g only in degree 1.
- Hence the induced map is zero.

WHAT STANDARD PERSISTENCE CAN'T SEE (BUT R1SC CAN)



- f has persistent reduced homology only in degree 0,
 g only in degree 1.
- Hence the induced map is zero.

But $h(\varphi)$ is nonzero!

EXTENDED & LEVEL SET PERSISTENCE

Extended persistence [Cohen-Steiner, Edelsbrunner, Harer '08]

$$\cdots \hookrightarrow H_*(f^{-1}(-\infty, s]) \hookrightarrow \cdots \hookrightarrow H_*(f^{-1}(-\infty, t]) \hookrightarrow \cdots \hookrightarrow H_*(X) \rightarrow$$

$$\cdots \hookrightarrow H_*(X, f^{-1}[v, \infty)) \hookrightarrow \cdots \hookrightarrow H_*(X, f^{-1}[u, \infty)) \hookrightarrow \cdots$$

EXTENDED & LEVEL SET PERSISTENCE

Extended persistence [Cohen-Steiner, Edelsbrunner, Harer '09]

$$\cdots \hookrightarrow H_*(f^{-1}(-\infty, s]) \hookrightarrow \cdots \hookrightarrow H_*(f^{-1}(-\infty, t]) \hookrightarrow \cdots \hookrightarrow H_*(X) \rightarrow$$

$$\cdots \hookrightarrow H_*(X, f^{-1}[v, \infty)) \hookrightarrow \cdots \hookrightarrow H_*(X, f^{-1}[u, \infty)) \hookrightarrow \cdots$$

Theorem [Carlsson, de Silva, Morozov '09] The persistence diagrams of extended persistence and level set persistence are in bijective correspondence.

- Some features appear in different degrees across this correspondence

FUNCTORIAL EQUIVALENCE

Can we make the correspondence of extended and level set persistence functorial (an equivalence of categories)?

- Does the correspondence extend to morphisms of persistence modules in a natural way?

FUNCTORIAL EQUIVALENCE

Can we make the correspondence of extended and level set persistence functorial (an equivalence of categories)?

- Does the correspondence extend to morphisms of persistence modules in a natural way?

Not in the standard way

(persistent homology as graded persistence modules)

FUNCTORIAL EQUIVALENCE

Can we make the correspondence of extended and level set persistence functorial (an equivalence of categories)?

- Does the correspondence extend to morphisms of persistence modules in a natural way?


Not in the standard way

(persistent homology as graded persistence modules)

Yes using R1SC

(level set persistence in a derived category)

R1SC \cong DERIVED LEVEL SET PERSISTENCE



R1SC \cong DERIVED LEVEL SET PERSISTENCE

Derived
Level
Set
Persistence

$$f : X \rightarrow \mathbb{R}$$

$$Rf_* (h_X) : D^+ (\mathrm{Sh}(\mathbb{R}))$$

$$h(f) : M \rightarrow \text{rect}$$

Relative
Inter-level
Set
Cohomology

R1SC \cong DERIVED LEVEL SET PERSISTENCE

Derived
Level
Set
Persistence

Relative
Inter-level
Set
Cohomology

$$\begin{array}{ccc} f : X \rightarrow R & & \\ \downarrow & & \downarrow \\ Rf_*(h_X) : D^+(\text{Sh}(R)) & \xrightarrow{\cong} & h(f) : M \rightarrow \text{vect} \\ (\text{tame}) & & (\text{cohomological, bounded above}) \end{array}$$

R1SC \cong DERIVED LEVEL SET PERSISTENCE

Derived
Level
Set
Persistence

$$\begin{array}{ccc} & \text{(X locally contractible)} & \\ f : X \rightarrow R & \text{(tame)} & \\ \downarrow & \text{\textcircled{\text{G}}\textcircled{\text{G}}} & \downarrow \\ Rf_*(h_X) : D^+(\mathrm{Sh}(R)) & \xrightarrow{\cong} & h(f) : M \rightarrow \mathrm{vect} \\ \text{(tame)} & & \text{(cohomological, bounded above)} \end{array}$$

Relative
Inter-level
Set
Cohomology

R1SC \cong DERIVED LEVEL SET PERSISTENCE

Derived
Level
Set
Persistence

$$\begin{array}{ccc} & (\times \text{ locally contractible}) & \\ f: X \rightarrow R & \swarrow & \searrow \\ Rf_*(h_X): & D^+(\mathrm{Sh}(R)) & \xrightarrow{\cong} M \rightarrow \mathrm{vect} \\ (\text{tame}) & & (\text{cohomological, bounded above}) \end{array}$$

\cong

Relative
Inter-level
Set
Cohomology

Theorem [B, Fluhr '22] DLSP \cong R1SC.

R1SC \cong DERIVED LEVEL SET PERSISTENCE

Derived
Level
Set
Persistence

$$\begin{array}{ccc} & (\times \text{ locally contractible}) & \\ f: X \rightarrow R & \swarrow & \searrow \\ Rf_*(h_X): & D^+(\mathrm{Sh}(R)) & \xrightarrow{\cong} M \rightarrow \mathrm{vect} \\ & (\text{tame}) & \\ & & h(f): \\ & & \text{(cohomological, bounded above)} \end{array}$$

Theorem [B, Fluhr '22] DLSP \cong R1SC.

Depends crucially on morphisms across degrees!

CATEGORIFICATION OF EXTENDED PERSISTENCE

categorification: replacing set-theoretic notions by categorical ones

- more structure (in particular : morphisms)

CATEGORIFICATION OF EXTENDED PERSISTENCE

categorification: replacing set-theoretic notions by categorical ones

- more structure (in particular : morphisms)

Standard example:

The Grothendieck group of an abelian category *

$$K_0(\mathcal{A}) = \{\text{iso classes of } \mathcal{A}\} / [A] + [C] \sim [B]$$

V.s.e.s $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$

CATEGORIFICATION OF EXTENDED PERSISTENCE

categorification: replacing set-theoretic notions by categorical ones

- more structure (in particular : morphisms)

Standard example:

The Grothendieck group of an abelian category *

$$K_0(\mathcal{A}) = \{\text{iso classes of } \mathcal{A}\} / [A] + [C] \sim [B] \quad \text{v.s.e.s}$$
$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

Abelian categorification of an abelian group G :

- abelian category \mathcal{A} & isomorphism $K_0(\mathcal{A}) \xrightarrow{\cong} G$

CATEGORIFICATION OF EXTENDED PERSISTENCE

categorification: replacing set-theoretic notions by categorical ones

- more structure (in particular : morphisms)

Standard example:

The Grothendieck group of an abelian category *

$$K_0(\mathcal{A}) = \{\text{iso classes of } \mathcal{A}\} / [A] + [C] \sim [B] \quad \forall \text{s.e.s} \\ 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

Abelian categorification of an abelian group G :

- abelian category \mathcal{A} & isomorphism $K_0(\mathcal{A}) \xrightarrow{\cong} G$

Categorify (the group of signed) extended persistence diagrams!

ABELIANIZATION OF COHOMOLOGICAL STRIP DIAGRAMS

Let J : category of RISC:
cohomological $M \rightarrow$ rect bounded above.

ABELIANIZATION OF COHOMOLOGICAL STRIP DIAGRAMS

Let J : category of RISC:

cohomological $M \rightarrow \text{vect}$ bounded above.

pres J : J -presentable objects in $M \rightarrow \text{vect}$:

$$P_1 \xrightarrow{f} P_0 \rightarrow M = \text{coker } f \quad (P_1, P_0 \in J)$$

ABELIANIZATION OF COHOMOLOGICAL STRIP DIAGRAMS

Let J : category of RISC:

cohomological $M \rightarrow \text{vect}$ bounded above.

$\text{pres } J$: J -presentable objects in $M \rightarrow \text{vect}$:

$$P_1 \xrightarrow{f} P_0 \rightarrow M = \text{coker } f \quad (P_1, P_0 \in J)$$

Key facts

- $\text{pres } J$ is the abelianization of J

ABELIANIZATION OF COHOMOLOGICAL STRIP DIAGRAMS

Let J : category of RISC:

cohomological $M \rightarrow \text{vect}$ bounded above.

$\text{pres } J$: J -presentable objects in $M \rightarrow \text{vect}$:

$$P_1 \xrightarrow{f} P_0 \rightarrow M = \text{coker } f \quad (P_1, P_0 \in J)$$

Key facts

- $\text{pres } J$ is the abelianization of J
- J are the projectives & injectives of $\text{pres } J$

ABELIANIZATION OF COHOMOLOGICAL STRIP DIAGRAMS

Let J : category of RISC:

cohomological $M \rightarrow \text{vect}$ bounded above.

$\text{pres } J$: J -presentable objects in $M \rightarrow \text{vect}$:

$$P_1 \xrightarrow{f} P_0 \rightarrow M = \text{coker } f \quad (P_1, P_0 \in J)$$

Key facts

- $\text{pres } J$ is the abelianization of J
- J are the projectives & injectives of $\text{pres } J$
- The "extended persistence map" on J extends to a "signed extended persistence map" on $\text{pres } J$

ABELIANIZATION OF COHOMOLOGICAL STRIP DIAGRAMS

Let J : category of RISC:

cohomological $M \rightarrow \text{vect}$ bounded above.

$\text{pres } J$: J -presentable objects in $M \rightarrow \text{vect}$:

$$P_1 \xrightarrow{f} P_0 \rightarrow M = \text{coker } f \quad (P_1, P_0 \in J)$$

Key facts

- $\text{pres } J$ is the abelianization of J
- J are the projectives & injectives of $\text{pres } J$
- The "extended persistence map" on J extends to a "signed extended persistence map" on $\text{pres } J$
- This map is additive \Rightarrow well-defined on K_0

R1SC CATEGORIFIES EXTENDED PERSISTENCE

Theorem [B, Flahr '22] The signed persistence map categorifies extended persistence diagrams:

$$K_0(\text{pres } \mathcal{J}) \xrightarrow{\cong} \{\text{signed ext. PDs}\}$$

TAKE-HOME MESSAGES

Indecomposables in multi-parameter persistence:

- can be complicated, even in H_0

TAKE-HOME MESSAGES

Indecomposables in multi-parameter persistence:

- can be complicated, even in H_0
- are close to every persistence module

TAKE-HOME MESSAGES

Indecomposables in multi-parameter persistence:

- can be complicated, even in H_0
- are close to every persistence module
- clarify the structure of interleval set persistence

TAKE-HOME MESSAGES

Indecomposables in multi-parameter persistence:

- can be complicated, even in H_0
- are close to every persistence module
- clarify the structure of interleval set persistence
- illuminate the role of derived categories in persistence

TAKE-HOME MESSAGES

Indecomposables in multi-parameter persistence:

- can be complicated, even in H_0
- are close to every persistence module
- clarify the structure of interleval set persistence
- illuminate the role of derived categories in persistence
- lead to the categorification of extended persistence