

The Morse theory of Čech and Delaunay complexes

Ulrich Bauer


TUM

April 27, 2017


Oberwolfach workshop
Computational geometry

Joint work with Herbert Edelsbrunner (IST Austria)


Connect the dots: topology from geometry




Connect the dots: topology from geometry




Connect the dots: topology from geometry




Connect the dots: topology from geometry




Connect the dots: topology from geometry



Connect the dots: topology from geometry



Connect the dots: topology from geometry



Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)



Čech and Delaunay functions


$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X
(abstract simplices)

Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X
(abstract simplices)




Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X
(abstract simplices)

Two functions on simplices $Q \subseteq X$:

- Čech function $f_C(Q)$:
radius of smallest enclosing sphere of Q




$\check{\text{C}}\text{ech}$ and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X
(abstract simplices)

Two functions on simplices $Q \subseteq X$:

- *$\check{\text{C}}\text{ech function}$* $f_C(Q)$:
radius of smallest enclosing sphere of Q
- *Delaunay function* $f_D(Q)$:
radius of smallest empty circumsphere




Čech and Delaunay functions

$X \subset \mathbb{R}^d$: finite point set (in general position)

Simplices $\Delta(X)$: nonempty subsets of X
(abstract simplices)

Two functions on simplices $Q \subseteq X$:

- *Čech function* $f_C(Q)$:
radius of smallest enclosing sphere of Q
- *Delaunay function* $f_D(Q)$:
radius of smallest empty circumsphere
 - defined only if Q has an empty circumsphere: $Q \in \text{Del}(X)$



Čech and Delaunay complexes from functions

Define for any radius r :

Čech and Delaunay complexes from functions


Define for any radius r :

- Čech complex $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$

Čech and Delaunay complexes from functions

Define for any radius r :


- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$
 - restriction of Čech complex to Delaunay simplices



Čech and Delaunay complexes from functions

Define for any radius r :


- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$
 - restriction of Čech complex to Delaunay simplices
- *Delaunay complex* (α -shape, for $\alpha = r$) $\text{Del}_r = f_D^{-1}(-\infty, r]$
 - all simplices having an empty circumsphere of radius $\leq r$



Čech and Delaunay complexes from functions

Define for any radius r :

- *Čech complex* $\text{Cech}_r = f_C^{-1}(-\infty, r]$
 - all simplices having an enclosing sphere of radius $\leq r$
- *Delaunay–Čech complex* $\text{DelCech}_r = \text{Del} \cap \text{Cech}_r$
 - restriction of Čech complex to Delaunay simplices
- *Delaunay complex* (α -shape, for $\alpha = r$) $\text{Del}_r = f_D^{-1}(-\infty, r]$
 - all simplices having an empty circumsphere of radius $\leq r$



Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

Properties of Čech and Delaunay complexes

By the *Nerve theorem* (Borsuk 1947):

$$\text{Del}_r(X) \simeq \text{Cech}_r(X) \simeq B_r(X).$$

But we also have

$$\text{Del}_r(X) \subseteq \text{DelCech}_r(X) \subseteq \text{Cech}_r(X).$$

- Are all three complexes homotopy equivalent?
- Can we express the equivalence combinatorially?

Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014/17, SoCG/Trans. AMS)

$\check{\text{C}}\text{ech}$, Delaunay– $\check{\text{C}}\text{ech}$, Delaunay, and Wrap complexes are homotopy equivalent through a sequence of collapses

$$\text{Cech}_r X \searrow \text{Cech}_r X \cap \text{Del} X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$



Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014/17, SoCG/Trans. AMS)

Čech, Delaunay–Čech, Delaunay, and Wrap complexes are homotopy equivalent through a sequence of collapses

$$\text{Cech}_r X \searrow \text{Cech}_r X \cap \text{Del} X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$



- All collapses are induced by a single *discrete gradient field*
- Explicit chain maps inducing isomorphisms in homology
- Also works for weighted point sets

Main result: a sequence of collapses

Theorem (B, Edelsbrunner 2014/17, SoCG/Trans. AMS)

Čech, Delaunay–Čech, Delaunay, and Wrap complexes are homotopy equivalent through a sequence of collapses

$$\text{Cech}_r X \searrow \text{Cech}_r X \cap \text{Del} X \searrow \text{Del}_r X \searrow \text{Wrap}_r X.$$




- All collapses are induced by a single *discrete gradient field*
- Explicit chain maps inducing isomorphisms in homology
- Also works for weighted point sets

Discrete Morse theory

Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, where L is a face of U with codimension 1.




Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, where L is a face of U with codimension 1.


- indicated by an arrow from L to U



Discrete Morse theory

Definition (Forman 1998)

A *discrete vector field* on a simplicial complex is a partition of the simplices into singletons and pairs $\{L, U\}$, where L is a face of U with codimension 1.



- indicated by an arrow from L to U


The singletons are called *critical simplices*.

Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes



Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes

•
1

Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ form a discrete vector field (the *discrete gradient* of f)




Discrete Morse theory

Definition (Forman 1998)

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex is a *discrete Morse function* if for all $t \in \mathbb{R}$:


- the sublevel sets $K_t = f^{-1}(-\infty, t]$ are subcomplexes
- the level sets $f^{-1}(t)$ form a discrete vector field (the *discrete gradient* of f)



If $f^{-1}(t) = \{Q\}$ then t is a *critical value*.

Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

*If $(s, t]$ contains no critical value of f ,
then $K_t \searrow K_s$.*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

*If $(s, t]$ contains no critical value of f ,
then $K_t \searrow K_s$.*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

*If $(s, t]$ contains no critical value of f ,
then $K_t \searrow K_s$.*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)


*If $(s, t]$ contains no critical value of f ,
then $K_t \searrow K_s$.*



Let V be a discrete gradient field on a simplicial complex K ,
and let L be a subcomplex of K .

Corollary

*If $K \setminus L$ is the union of some pairs of V ,
then $K \searrow L$.*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)


*If $(s, t]$ contains no critical value of f ,
then $K_t \searrow K_s$.*



Let V be a discrete gradient field on a simplicial complex K ,
and let L be a subcomplex of K .

Corollary

*If $K \setminus L$ is the union of some pairs of V ,
then $K \searrow L$.*




Collapses from Morse functions and gradients

Let f be a discrete Morse function on a simplicial complex K .

Theorem (Forman 1998)

*If $(s, t]$ contains no critical value of f ,
then $K_t \searrow K_s$.*




Let V be a discrete gradient field on a simplicial complex K ,
and let L be a subcomplex of K .

Corollary

*If $K \setminus L$ is the union of some pairs of V ,
then $K \searrow L$.*


We say that V induces the collapse $K \searrow L$.




Unfortunately...

Neither the Čech nor the Delaunay functions
are discrete Morse functions!

- Example: two simplices Q, Q' with $f_C(Q) = f_C(Q')$
such that neither is a face of the other:



Q




Q'

Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$




- indicated by an arrow from L to U

Generalized discrete Morse theory

Definition (Chari 2000, Freij 2009)

A *generalized discrete vector field* on a simplicial complex is a partition of the simplices into intervals of the face poset:

$$[L, U] = \{Q : L \subseteq Q \subseteq U\}$$



- indicated by an arrow from L to U


The singletons are called *critical simplices*.

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes



Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes

•
1

Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes



Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes



Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:

- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes




Generalized discrete Morse theory

Definition

A function $f : K \rightarrow \mathbb{R}$ on a simplicial complex
is a *generalized discrete Morse function* if for $t \in \mathbb{R}$:


- the sublevel sets $K_t = f^{-1}(-\infty, t]$
are subcomplexes
- the level sets $f^{-1}(t)$ form a generalized
vector field (the *discrete gradient* of f)



Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.




Refining generalized vector fields

A generalized vector field V can be refined to a vector field.

For each non-critical interval $[L, U] \in V$:

- choose an arbitrary vertex $x \in U \setminus L$
- partition $[L, U]$ into pairs $(Q \setminus \{x\}, Q \cup \{x\})$ for all $Q \in [L, U]$.



Therefore the collapsing theorems also hold for generalized discrete Morse functions.

Čech and Delaunay intervals

Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*


Morse theory of Čech and Delaunay complexes

Proposition (B, Edelsbrunner 2014)

*The Čech function and the Delaunay function
are generalized discrete Morse functions.*

The following are equivalent for each simplex $Q \subseteq X$:


- *Q is a critical simplex of f_C*
- *Q is a critical simplex of f_D*
- $f_D(Q) = f_C(Q)$
- *Q is a centered Delaunay simplex
(containing the circumcenter in its interior)*



Čech intervals

Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff




Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff

On $S \subseteq Q'$



Q




On S

$\check{\text{C}}$ ech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$\text{On } S \subseteq Q' \subseteq \text{Encl } S.$$



Q



$\text{On } S$




$\text{Encl } S$

Čech intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest enclosing sphere S .
Then $Q' \subseteq X$ has the same smallest enclosing sphere S iff


$$Q' \in [\text{On } S, \text{Encl } S].$$



Q



$\text{On } S$



$\text{Encl } S$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex On $S \subseteq X$.

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.

Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.


Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$



The front and back faces of a simplex

Let S be the smallest circumsphere of a simplex $\text{On } S \subseteq X$.

Then the center z of S is an affine combination of those points:

$$z = \sum_{x \in \text{On } S} \mu_x x.$$

We define

$$\text{Front } S = \{x \in \text{On } S \mid \mu_x > 0\},$$

$$\text{Back } S = \{x \in \text{On } S \mid \mu_x < 0\}.$$




Delaunay intervals


Lemma

Let $Q \subseteq X$ be a simplex with smallest empty circumsphere S .
Then $Q' \subseteq X$ has the same smallest empty circumsphere S iff


$$\text{Front } S \subseteq Q' \subseteq \text{On } S.$$



Q



$\text{Front } S$



$\text{On } S$

Selective Delaunay complexes

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

minimize
 r,z

r

subject to

$$\begin{aligned}\|z - q\| &\leq r, & q \in Q, \\ \|z - e\| &\geq r, & e \in E.\end{aligned}$$

Here r is the radius of the sphere S , and z is the center of S .

Sphere optimization problems

Both Čech and Delaunay functions are defined using the smallest sphere S satisfying certain constraints:

minimize
 r,z

r

subject to

$$\begin{aligned}\|z - q\| &\leq r, & q \in Q, \\ \|z - e\| &\geq r, & e \in E.\end{aligned}$$

Here r is the radius of the sphere S , and z is the center of S .

- Čech function: choose $E = \emptyset$
- Delaunay function: choose $E = X$

Čech and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

Proposition

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of Q ,
- $\text{Front } S \subseteq Q \subseteq \text{Encl } S$, and
- $\text{Back } S \subseteq E \subseteq \text{Excl } S$.

$\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The *Karush–Kuhn–Tucker* optimality conditions yield:

Proposition

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $\text{Front } S \subseteq Q \subseteq \text{Encl } S$, and
- $\text{Back } S \subseteq E \subseteq \text{Excl } S$.

Corollary

The $\check{\text{C}}\text{ech}$ intervals are of the form $[\text{On } S, \text{Encl } S]$.

$\check{\text{C}}\text{ech}$ and Delaunay intervals from KKT

The Karush–Kuhn–Tucker optimality conditions yield:

Proposition

A sphere S enclosing Q and excluding E is minimal iff

- S is the smallest circumsphere of $\text{On } S$,
- $\text{Front } S \subseteq Q \subseteq \text{Encl } S$, and
- $\text{Back } S \subseteq E \subseteq \text{Excl } S$.

Corollary

The $\check{\text{C}}\text{ech}$ intervals are of the form $[\text{On } S, \text{Encl } S]$.

The Delaunay intervals are of the form $[\text{Front } S, \text{On } S]$.

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$
- *Selective Delaunay complex* $\text{Del}_r(X, E) = f_D^{-1}(-\infty, r]$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$
- *Selective Delaunay complex* $\text{Del}_r(X, E) = f_D^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2017)

Let $E \subseteq F \subseteq X$. Then

$$\text{Del}_r(X, E) \setminus \text{Del}_r(X, E) \cap \text{Del}(X, F) \setminus \text{Del}_r(X, F).$$

Selective Delaunay complexes

Define for finite point sets $X, E \subset \mathbb{R}^d$:

- *Selective Delaunay function* $f_E(Q)$:
radius of smallest sphere enclosing Q and excluding E
 - defined only if Q has an E -empty enclosing sphere:
 $Q \in \text{Del}(X, E)$
- *Selective Delaunay complex* $\text{Del}_r(X, E) = f_D^{-1}(-\infty, r]$

Theorem (B, Edelsbrunner 2017)

Let $E \subseteq F \subseteq X$. Then

$$\text{Del}_r(X, E) \setminus \text{Del}_r(X, E) \cap \text{Del}(X, F) \setminus \text{Del}_r(X, F).$$

Choosing $E = \emptyset$ and $F = X$ yields


$$\text{Cech}_r(X) \setminus \text{DelCech}_r(X) \setminus \text{Del}_r(X).$$

Connecting different Delaunay complexes

X




Y



Connecting different Delaunay complexes

$B_r(X)$




$B_r(Y)$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$




Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



$\text{Del}_r(X)$





$\text{Del}_r(X \cup Y)$



$\text{Del}_r(Y)$



Connecting different Delaunay complexes

$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



Connecting different Delaunay complexes


$$B_r(X) \longleftrightarrow B_r(X \cup Y) \longleftrightarrow B_r(Y)$$



Wrap complexes


Wrap complexes

Generalizes and greatly simplifies the surface reconstruction algorithm *Wrap* (Edelsbrunner 1995, Geomagic)



Wrap complexes


Consider the Delaunay function f_D of X .



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order



Wrap complexes

Consider the Delaunay function f_D of X .


- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order



Wrap complexes

Consider the Delaunay function f_D of X .

- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order




Wrap complexes

Consider the Delaunay function f_D of X .

- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$




Wrap complexes

Consider the Delaunay function f_D of X .

- Let $V_D = \{f_D^{-1}(t) : t \in \mathbb{R}\}$ be the Delaunay intervals
- The face relation induces a partial order on V_D
- Let $C \in \text{Crit}_r$ be a critical simplex with value $f_D(C) \leq r$
- Let $\downarrow C$ denote the descending set of C in that partial order

Define

$$\text{Wrap}_r = \bigcup_{C \in \text{Crit}_r} \downarrow C$$



The Delaunay intervals induce a collapse $\text{Del}_r \searrow \text{Wrap}_r$.

Wrapping up

- Čech and Delaunay complexes from Morse functions
- Explicit homotopy equivalence by simplicial collapses
- Simple definition and generalization of *Wrap* complexes