

# Persistent homology

## From theory to computation

Ulrich Bauer



Oct 26, 2019

50th anniversary workshop  
Institute for Numerical and Applied Mathematics  
University of Göttingen



112



?




112





?


# Persistent homology



















## Gramicidin: an antibiotic functioning as ion channel




## Gramicidin: an antibiotic functioning as ion channel




# Homology




# Homology




# Homology




# Homology




# Homology





# Homology




## What is persistent homology?




## What is persistent homology?





## What is persistent homology?





# What is persistent homology?





# What is persistent homology?



## What is persistent homology?



## What is persistent homology?



## What is persistent homology?

## What is persistent homology?

Persistent homology is the homology of a filtration.

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain commutative diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ :
  - $\mathbf{R}$  is the partially ordered set  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain commutative diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ :
  - $\mathbf{R}$  is the partially ordered set  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain commutative diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ :
  - $\mathbf{R}$  is the partially ordered set  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- Persistent homology is a commutative diagram  $M : \mathbf{R} \rightarrow \mathbf{Vect}$  (*persistence module*):
  - A vector space  $M_t$  for each  $t \in \mathbb{R}$
  - A linear map  $M_s \rightarrow M_t$  for each  $s \leq t \in \mathbb{R}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain commutative diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ :
  - $\mathbf{R}$  is the partially ordered set  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- Persistent homology is a commutative diagram  $M : \mathbf{R} \rightarrow \mathbf{Vect}$  (*persistence module*):
  - A vector space  $M_t$  for each  $t \in \mathbb{R}$
  - A linear map  $M_s \rightarrow M_t$  for each  $s \leq t \in \mathbb{R}$

In this talk, all vector spaces will be finite dimensional.

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

(in an essentially unique way).

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

(in an essentially unique way).

- The supporting intervals form the *persistence barcode*  $B(M)$ .

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

(in an essentially unique way).

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The barcode completely describes the persistence module (up to isomorphism).

# Stability


# Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that



# Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that

- matched intervals have endpoints within distance  $\leq \delta$ , and



# Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .




## Persistence and stability: the big picture

Data


point cloud

$$P \subset \mathbb{R}^d$$


## Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




## Geometric interleavings




## Geometric interleavings




## Geometric interleavings




## Geometric interleavings



## Geometric interleavings



## Geometric interleavings



## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:

$$\begin{array}{ccccccc} \dots & \rightarrow & F_t & \hookrightarrow & F_{t+\delta} & \hookrightarrow & F_{t+2\delta} & \dots \rightarrow \\ & & \swarrow \searrow & & \swarrow \searrow & & & \\ \dots & \rightarrow & G_t & \hookrightarrow & G_{t+\delta} & \hookrightarrow & G_{t+2\delta} & \dots \rightarrow \end{array} \quad \forall t \in \mathbb{R}.$$

## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:

$$\begin{array}{ccccccc} \dots & \rightarrow & F_t & \hookrightarrow & F_{t+\delta} & \hookrightarrow & F_{t+2\delta} & \dots \rightarrow \\ & & \swarrow \times \searrow & & \swarrow \times \searrow & & & \\ \dots & \rightarrow & G_t & \hookrightarrow & G_{t+\delta} & \hookrightarrow & G_{t+2\delta} & \dots \rightarrow \end{array} \quad \forall t \in \mathbb{R}.$$

Applying homology (a functor) preserves commutativity


- persistent homology of  $f, g$  yields  $\delta$ -interleaved persistence modules  $\mathbf{R} \rightarrow \mathbf{Vect}$

# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .

Equivalently: there exists a  $\delta$ -interleaving of their barcodes (as diagrams  $R \rightarrow Mch$ ).



## Structure of persistence submodules and quotients

### Proposition

Let  $M \twoheadrightarrow N$  be an epimorphism.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial (compatible with composition and identity).



# Structure of persistence submodules and quotients

## Proposition

Let  $M \twoheadrightarrow N$  be an epimorphism.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial (compatible with composition and identity).



Dually, there is an injection  $B(N) \hookrightarrow B(O)$  for monomorphisms  $N \hookrightarrow O$ .



## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:

## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:

- compose the injections  $B(M) \hookrightarrow B(\text{im } f) \hookrightarrow B(N)$  from before to a matching

$$\chi(f) : B(M) \not\rightarrow B(N).$$



## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:

- compose the injections  $B(M) \hookleftarrow B(\text{im } f) \hookrightarrow B(N)$  from before to a matching

$$\chi(f) : B(M) \not\rightarrow B(N).$$



When applied to a  $\delta$ -interleaving, this yields a  $\delta$ -matching of barcodes.

# Homology inference

## Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .*

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .*

- Approach: approximate the shape by a thickening  $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covering  $X$

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .*

- Approach: approximate the shape by a thickening  $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covering  $X$
- Requires strong assumptions:


## Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- the induced maps  $H_*(X \hookrightarrow X_\delta)$  and  $H_*(X_\delta \hookrightarrow X_{2\delta})$  are isomorphisms.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .





# Homological realization

This motivates the *homological realization problem*:

## Problem

Given a pair  $L \subseteq K$  of simplicial complexes, find a third complex  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ .



Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)


The homological realization problem is NP-hard, even in  $\mathbb{R}^3$ .

# Homological realization

This motivates the *homological realization problem*:

## Problem


Given a pair  $L \subseteq K$  of simplicial complexes, find a third complex  $X$  with  $L \subseteq X \subseteq K$  such that  $H_*(X) \cong \text{im } H_*(L \hookrightarrow K)$ .




Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)

The homological realization problem is NP-hard, even in  $\mathbb{R}^3$ .

# Simplification





## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}, \delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , the linear map  $H_*(F_{t-\delta} \hookrightarrow F_{t+\delta})$  factors through  $H_*(G_t)$ .
- This gives a lower bound

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}, \delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , the linear map  $H_*(F_{t-\delta} \hookrightarrow F_{t+\delta})$  factors through  $H_*(G_t)$ .
- This gives a lower bound

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

- But it is NP-hard to decide whether this bound can be achieved.

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , the linear map  $H_*(F_{t-\delta} \hookrightarrow F_{t+\delta})$  factors through  $H_*(G_t)$ .
- This gives a lower bound

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

- But it is NP-hard to decide whether this bound can be achieved.

### Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)

Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.











## Topological simplification of functions

Consider the following problem:


### Problem (Topological simplification)

*Given a function  $f$  and a real number  $\delta \geq 0$ , find a function  $f_\delta$  subject to  $\|f_\delta - f\|_\infty \leq \delta$  with the minimal number of critical points.*

## Persistence vs Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points




(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

## Persistence vs Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points



(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

Persistent homology:

- Relates homology of different sublevel set
- Identifies pairs of critical points (birth and death of homological feature) and quantifies their *persistence*

# Combining persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

# Combining persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

## Theorem (B, Lange, Wardetzky, 2011)

*Let  $f$  be a function on a surface and let  $\delta > 0$ .*

*Then canceling all persistence pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$  with*

$$\|f_\delta - f\|_\infty \leq \delta$$

*that achieves the lower bound on the number of critical points.*

# Combining persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

## Theorem (B, Lange, Wardetzky, 2011)

*Let  $f$  be a function on a surface and let  $\delta > 0$ .*

*Then canceling all persistence pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$  with*

$$\|f_\delta - f\|_\infty \leq \delta$$

*that achieves the lower bound on the number of critical points.*

- Does not generalize to higher-dimensional manifolds!

# Computation

## Vietoris–Rips complexes

Consider a finite metric space  $(X, d)$ .

The *Vietoris–Rips complex* is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid \text{diam } S \leq t\}$$

- all edges with pairwise distance  $\leq t$
- all possible higher simplices

## Vietoris–Rips complexes

Consider a finite metric space  $(X, d)$ .

The *Vietoris–Rips complex* is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid \text{diam } S \leq t\}$$

- all edges with pairwise distance  $\leq t$
- all possible higher simplices

For large  $t$ ,  $\text{Rips}_t(X)$  is the full simplex with  $n = |X|$  vertices

- Number of  $d$ -simplices is  $\binom{n}{d+1}$
- Computation is one of the most important challenges in applied topology!

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 596 seconds, 3.4 GB
- DIPHA (IST Austria): 53 seconds, 5.5 GB
- GUDHI (INRIA): 46 seconds, 2.6 GB

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 596 seconds, 3.4 GB
- DIPHA (IST Austria): 53 seconds, 5.5 GB
- GUDHI (INRIA): 46 seconds, 2.6 GB

Demo: [live.ripser.org](http://live.ripser.org)

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 596 seconds, 3.4 GB
- DIPHA (IST Austria): 53 seconds, 5.5 GB
- GUDHI (INRIA): 46 seconds, 2.6 GB

Demo: [live.ripser.org](http://live.ripser.org)

- Ripser: 1.1 seconds, 196 MB

Thanks for your attention!

