

# Persistent Homology

## From Theory to Computation

Ulrich Bauer

TUM

May 2, 2019

Mathematical Colloquium, Bielefeld University

Workshop Computational Applications of Quiver Representations: TDA and QPA



11



?




11





?


# Persistent homology






























## Gramicidin: an antibiotic functioning as ion channel





## Gramicidin: an antibiotic functioning as ion channel





















## What is persistent homology?





## What is persistent homology?





# What is persistent homology?




# What is persistent homology?




# What is persistent homology?



# What is persistent homology?



# What is persistent homology?



## What is persistent homology?

## What is persistent homology?

Persistent homology is the homology of a filtration.

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - $\mathbf{R}$  is the poset  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - $\mathbf{R}$  is the poset  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - $\mathbf{R}$  is the poset  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- Persistent homology is a diagram  $M : \mathbf{R} \rightarrow \mathbf{Vect}$  (*persistence module*)

# What is persistent homology?

Persistent homology is the homology of a filtration.

- A filtration is a certain diagram  $K : \mathbf{R} \rightarrow \mathbf{Top}$ 
  - $\mathbf{R}$  is the poset  $(\mathbb{R}, \leq)$
  - A topological space  $K_t$  for each  $t \in \mathbb{R}$
  - An inclusion map  $K_s \hookrightarrow K_t$  for each  $s \leq t \in \mathbb{R}$
- Apply homology  $H_* : \mathbf{Top} \rightarrow \mathbf{Vect}$
- Persistent homology is a diagram  $M : \mathbf{R} \rightarrow \mathbf{Vect}$  (*persistence module*)

In this talk, all vector spaces will be finite dimensional.

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

*Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules*

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

*(in an essentially unique way).*

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

*Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules*

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

*(in an essentially unique way).*

- The supporting intervals form the *persistence barcode*  $B(M)$ .

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

(in an essentially unique way).

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The barcode completely describes the persistence module (up to isomorphism).

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

(in an essentially unique way).

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The barcode completely describes the persistence module (up to isomorphism).
- This is why we use homology with coefficients in a field.

## Barcodes: the structure of persistence modules

Theorem (Crawley-Boevey 2015)

Any persistence module  $M : \mathbf{R} \rightarrow \mathbf{vect}$  (of finite dim. vector spaces over some field  $\mathbb{F}$ ) decomposes as a direct sum of interval modules

$$0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbb{F} \rightarrow \cdots \rightarrow \mathbb{F} \rightarrow 0 \rightarrow \cdots \rightarrow 0$$

(in an essentially unique way).

- The supporting intervals form the *persistence barcode*  $B(M)$ .
- The barcode completely describes the persistence module (up to isomorphism).
- This is why we use homology with coefficients in a field.
- We rarely have a similarly clean structure for other indexing posets, like  $\mathbf{R}^2 \rightarrow \mathbf{vect}$  (two-parameter persistence modules)

# Stability


## Stability of persistence barcodes for functions

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that



# Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that

- matched intervals have endpoints within distance  $\leq \delta$ , and



# Stability of persistence barcodes for functions


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $f, g : X \rightarrow \mathbb{R}$  with  $\|f - g\|_\infty = \delta$  (and some regularity assumptions).

Consider the persistence barcodes of (sublevel set filtrations of)  $f$  and  $g$ .

Then there exists a matching between their intervals such that

- matched intervals have endpoints within distance  $\leq \delta$ , and
- unmatched intervals have length  $\leq 2\delta$ .




## Persistence and stability: the big picture

Data


point cloud

$$P \subset \mathbb{R}^d$$


# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture




# Persistence and stability: the big picture



# Persistence and stability: the big picture




# Persistence and stability: the big picture



## The category of matchings

Consider the category **Mch** with

- objects: sets,
- morphisms: matchings (bijections between subsets).




# The category of matchings

Consider the category **Mch** with

- objects: sets,
- morphisms: matchings (bijections between subsets).

Composition:




# The category of matchings


Consider the category **Mch** with

- objects: sets,
- morphisms: matchings (bijections between subsets).

Composition:




(Co)kernel/(co)image:




## From barcodes to matching diagrams (and back)

- A barcode (collection of intervals) defines a diagram  $\mathbf{R} \rightarrow \mathbf{Mch}$ :





$t \mapsto \{\text{intervals in barcode containing } t\}$



$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$

## From barcodes to matching diagrams (and back)


- A barcode (collection of intervals) defines a diagram  $\mathbf{R} \rightarrow \mathbf{Mch}$ :



$t \mapsto \{\text{intervals in barcode containing } t\}$

$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$



- A matching diagram defines a barcode:



- intervals are formed by equivalence classes of matched elements

## From barcodes to matching diagrams (and back)


- A barcode (collection of intervals) defines a diagram  $\mathbf{R} \rightarrow \mathbf{Mch}$ :



$t \mapsto \{\text{intervals in barcode containing } t\}$

$(s \leq t) \mapsto \{\text{intervals containing both } s, t\}$


- A matching diagram defines a barcode:




- intervals are formed by equivalence classes of matched elements

We can turn barcodes into a category  $\mathbf{Barc} \simeq \mathbf{Mch}^{\mathbf{R}}$


## Geometric interleavings




## Geometric interleavings




## Geometric interleavings




## Geometric interleavings



## Geometric interleavings



## Geometric interleavings



## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:

$$\begin{array}{ccccccc} \dots & \rightarrow & F_t & \hookrightarrow & F_{t+\delta} & \hookrightarrow & F_{t+2\delta} & \dots & \rightarrow \\ & & \swarrow \searrow & & \swarrow \searrow & & & & \\ \dots & \rightarrow & G_t & \hookrightarrow & G_{t+\delta} & \hookrightarrow & G_{t+2\delta} & \dots & \rightarrow \end{array} \quad \forall t \in \mathbf{R}.$$

## Interleavings

Let  $\delta = \|f - g\|_\infty$ . Write  $F_t = f^{-1}(-\infty, t]$  for the  $t$ -sublevel set of  $f$ .

Then the sublevel set filtrations  $F, G : \mathbf{R} \rightarrow \mathbf{Top}$  are  $\delta$ -interleaved:

$$\begin{array}{ccccccc} \dots & \rightarrow & F_t & \hookrightarrow & F_{t+\delta} & \hookrightarrow & F_{t+2\delta} & \dots & \rightarrow \\ & & \swarrow \searrow & & \swarrow \searrow & & & & \\ \dots & \rightarrow & G_t & \hookrightarrow & G_{t+\delta} & \hookrightarrow & G_{t+2\delta} & \dots & \rightarrow \end{array} \quad \forall t \in \mathbb{R}.$$

Applying homology (a functor) preserves commutativity


- persistent homology of  $f, g$  yields  $\delta$ -interleaved persistence modules  $\mathbf{R} \rightarrow \mathbf{Vect}$

# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .




# Algebraic stability of persistence barcodes

Theorem (Chazal et al. 2009, 2012; B, Lesnick 2015)

If two persistence modules are  $\delta$ -interleaved,  
then there exists a  $\delta$ -matching of their barcodes:

- matched intervals have endpoints within distance  $\leq \delta$ ,
- unmatched intervals have length  $\leq 2\delta$ .

Equivalently: there exists a  $\delta$ -interleaving of their barcodes (as diagrams  $\mathbf{R} \rightarrow \mathbf{Mch}$ ).



## Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor

$B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.

## Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor

$B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



## Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor  
 $B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



### Proposition

*There is no functor  $\mathbf{vect} \rightarrow \mathbf{Mch}$  sending a  $d$ -dimensional vector space to a set with  $d$  elements.*

## Non-functoriality of persistence barcodes

Can a persistence module  $M$  be mapped to its barcode  $B(M)$  by a functor  
 $B : \mathbf{vect} \rightarrow \mathbf{Mch}$ ?

- This would preserve  $\delta$ -interleavings, and thus yield stability of persistence barcodes.



### Proposition

*There is no functor  $\mathbf{vect} \rightarrow \mathbf{Mch}$  sending a  $d$ -dimensional vector space to a set with  $d$  elements.*

But:

- there is a barcode functor for monos/epis of persistence modules  $\mathbf{vect}^R$ :

# Structure of persistence sub-/quotient modules

## Proposition

Let  $M \twoheadrightarrow N$  be an epimorphism.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial.



# Structure of persistence sub-/quotient modules

## Proposition

Let  $M \twoheadrightarrow N$  be an epimorphism.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial.



- Sort intervals with same left end by length, in both  $B(M)$  and  $B(N)$
- Match up intervals in order

# Structure of persistence sub-/quotient modules

## Proposition

Let  $M \twoheadrightarrow N$  be an epimorphism.

Then there is an injection of barcodes  $B(N) \hookrightarrow B(M)$  such that if  $J$  is mapped to  $I$ , then

- $I$  and  $J$  are aligned below, and
- $I$  bounds  $J$  above.

This construction is functorial.



- Sort intervals with same left end by length, in both  $B(M)$  and  $B(N)$
- Match up intervals in order

Dually, there is an injection  $B(M) \hookrightarrow B(N)$  for monomorphisms  $M \hookrightarrow N$ .

## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:

## Induced matchings

For  $f : M \rightarrow N$  a morphism of pfd persistence modules, the epi-mono factorization

$$M \twoheadrightarrow \text{im } f \hookrightarrow N$$

gives an *induced matching*  $\chi(f)$  between their barcodes:


- compose the functorial injections  $B(M) \hookleftarrow B(\text{im } f) \hookrightarrow B(N)$  from before to a matching

$$\chi(f) : B(M) \not\rightarrow B(N).$$



## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):



## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \twoheadrightarrow & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow \curvearrowleft & N_{t+\delta} \end{array}$$

## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \gg & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow \curvearrowright & N_{t+\delta} \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  with length  $> 2\delta$  to a similar interval in  $B(M)$



$B(N)$

## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \gg & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow & N_{t+\delta} \\ & & \curvearrowright & & \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  with length  $> 2\delta$  to a similar interval in  $B(M)$



$B(N(\delta))$

$B(N)$

## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \gg & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow \curvearrowright & N_{t+\delta} \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  with length  $> 2\delta$  to a similar interval in  $B(M)$




## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \gg & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow & N_{t+\delta} \\ & \swarrow & & \searrow & \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  with length  $> 2\delta$  to a similar interval in  $B(M)$




## Algebraic stability via induced matchings

Consider an interleaving  $f_t : M_t \rightarrow N_{t+\delta}$ ,  $g_t : N_t \rightarrow M_{t+\delta}$  ( $\forall t \in \mathbb{R}$ ):

$$\begin{array}{ccccc} & & M_t & & \\ & \nearrow g_{t-\delta} & & \searrow & \\ N_{t-\delta} & \longrightarrow \gg & \text{im } N_{t-\delta, t+\delta} & \hookleftarrow & N_{t+\delta} \\ & \swarrow & & \searrow & \end{array}$$

Construct an induced  $\delta$ -matching of barcodes:

- Match every interval in  $B(N)$  with length  $> 2\delta$  to a similar interval in  $B(M)$



# Homology inference

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .*

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .*

Approach:

- approximate the shape by a thickening  $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covering  $X$

# Homology inference

Given: finite sample  $P \subset X$  of unknown shape  $X \subset \mathbb{R}^d$

## Problem (Homology inference)

*Determine the homology  $H_*(X)$ .*

## Problem (Homological reconstruction)

*Construct a shape  $R$  with  $H_*(R) \cong H_*(X)$ .*

Approach:

- approximate the shape by a thickening  $P_\delta = \bigcup_{p \in P} B_\delta(p)$  covering  $X$

Requires strong assumptions:


## Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .




# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology reconstruction by thickening

Theorem (Niyogi, Smale, Weinberger 2006)

Let  $X$  be a submanifold of  $\mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ , and
- $\delta < \sqrt{3/20} \text{ reach}(X)$ .

Then  $H_*(X) \cong H_*(P_{2\delta})$ .





# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .



# Homology inference using persistence


Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.



# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccc} H_*(X) & H_*(X_\delta) & H_*(X_{2\delta}) \\ \searrow & \nearrow & \searrow \\ H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \searrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \swarrow & \nearrow \\ & & H_*(P_\delta) & & H_*(P_{2\delta}) \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \longleftrightarrow & H_*(X_\delta) & \hookrightarrow & H_*(X_{2\delta}) \\ \searrow & & \nearrow & & \downarrow \cong \\ H_*(P_\delta) & & & & H_*(P_{2\delta}) \\ & & \downarrow & & \swarrow \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

# Homology inference using persistence

Theorem (Cohen-Steiner, Edelsbrunner, Harer 2005)

Let  $X \subset \mathbb{R}^d$ . Let  $P \subset X$  and  $\delta > 0$  be such that

- $P_\delta$  covers  $X$ ,
- the induced map  $H_*(X \hookrightarrow X_\delta)$  is an isomorphism, and
- the induced map  $H_*(X_\delta \hookrightarrow X_{2\delta})$  is a monomorphism.

Then  $H_*(X) \cong \text{im } H_*(P_\delta \hookrightarrow P_{2\delta})$ .

Proof.

$$\begin{array}{ccccc} H_*(X) & \xhookleftarrow{\quad \cong \quad} & H_*(X_\delta) & \xhookrightarrow{\quad} & H_*(X_{2\delta}) \\ & \searrow & \nearrow & \downarrow \cong & \swarrow \nearrow \\ H_*(P_\delta) & & & H_*(P_{2\delta}) & \\ & \searrow & \downarrow & \nearrow & \\ & & \text{im } H_*(P_\delta \hookrightarrow P_{2\delta}) & & \end{array}$$

□

# Homological realization

This motivates the *homological realization problem*:

## Problem

*Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that*

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

*equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .*

# Homological realization


This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .



# Homological realization


This motivates the *homological realization problem*:

## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .

This is not always possible:



# Homological realization

This motivates the *homological realization problem*:


## Problem

Given a simplicial pair  $L \subseteq K$ , find  $X$  with  $L \subseteq X \subseteq K$  such that

$$H_*(L) \twoheadrightarrow H_*(X) \hookrightarrow H_*(K);$$

equivalently,  $\dim H_*(X) = \text{rank } H_*(L \hookrightarrow K)$ .


This is not always possible:




**Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)**

The homological realization problem is NP-hard, even in  $\mathbb{R}^3$ .

# Simplification





## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}$ ,  $\delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}, \delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , a lower bound is given by

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}, \delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .

- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , a lower bound is given by

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$

- But it is NP-hard to decide whether this bound can be achieved.

## Sublevel set simplification

Let  $F_t = f^{-1}(-\infty, t]$  denote the  $t$ -sublevel set of  $f$ .

### Problem (Sublevel set simplification)

Given a function  $f : X \subset \mathbb{R}^3 \rightarrow \mathbb{R}$  and  $t \in \mathbb{R}, \delta > 0$ ,

find a function  $g$  with  $\|g - f\|_\infty \leq \delta$  minimizing  $\dim H_*(G_t)$ .


- Since  $F_{t-\delta} \subseteq G_t \subseteq F_{t+\delta}$ , a lower bound is given by

$$\dim H_*(G_t) \geq \text{rank } H_*(F_{t-\delta} \hookrightarrow F_{t+\delta}).$$


- But it is NP-hard to decide whether this bound can be achieved.


### Theorem (Attali, B, Devillers, Glisse, Lieutier 2013)


Sublevel set simplification in  $\mathbb{R}^3$  is NP-hard.











# Topological simplification of functions

Consider the following problem:


## Problem (Topological simplification)

*Given a function  $f$  and a real number  $\delta \geq 0$ , find a function  $f_\delta$  subject to  $\|f_\delta - f\|_\infty \leq \delta$  with the minimal number of critical points.*

# Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points




(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

## Persistence and Morse theory

Morse theory (smooth or discrete):

- Relates critical points to homology of sublevel sets
- Provides a method for *cancelling* pairs of critical points



(from Milnor: *Lectures on the h-cobordism theorem*, 1965)

Persistent homology:

- Relates homology of different sublevel set
- Identifies pairs of critical points (birth and death of homological feature) and quantifies their *persistence*

# Persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

# Persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

## Theorem (B, Lange, Wardetzky, 2011)

*Let  $f$  be a function on a surface and let  $\delta > 0$ .*

*Then canceling all persistence pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$  with*

$$\|f_\delta - f\|_\infty \leq \delta$$

*that achieves the lower bound on the number of critical points.*

# Persistence and discrete Morse theory

By stability of persistence barcodes:

## Proposition

*The intervals in the barcode of  $f$  with persistence  $> 2\delta$  provide a lower bound on the number of critical points of any function  $g$  with  $\|g - f\|_\infty \leq \delta$ .*

## Theorem (B, Lange, Wardetzky, 2011)

*Let  $f$  be a function on a surface and let  $\delta > 0$ .*

*Then canceling all persistence pairs with persistence  $\leq 2\delta$  yields a function  $f_\delta$  with*

$$\|f_\delta - f\|_\infty \leq \delta$$

*that achieves the lower bound on the number of critical points.*

- Does not generalize to higher-dimensional manifolds!

# History

## When was persistent homology invented?



H. Edelsbrunner, D. Letscher, and A. Zomorodian

Topological persistence and simplification

*Foundations of Computer Science*, 2000

## When was persistent homology invented?

-  H. Edelsbrunner, D. Letscher, and A. Zomorodian  
Topological persistence and simplification  
*Foundations of Computer Science*, 2000
-  V. Robbins  
Computational Topology at Multiple Resolutions.  
PhD thesis, 2000

## When was persistent homology invented?

-  H. Edelsbrunner, D. Letscher, and A. Zomorodian  
Topological persistence and simplification  
*Foundations of Computer Science*, 2000
-  V. Robbins  
Computational Topology at Multiple Resolutions.  
PhD thesis, 2000
-  P. Frosini  
A distance for similarity classes of submanifolds of a Euclidean space  
*Bulletin of the Australian Mathematical Society*, 1990.

When was persistent homology invented first?

# When was persistent homology invented first?

ANNALS OF MATHEMATICS  
Vol. 41, No. 2, April, 1940

## RANK AND SPAN IN FUNCTIONAL TOPOLOGY

BY MARSTON MORSE

(Received August 9, 1939)

### 1. Introduction.

The analysis of functions  $F$  on metric spaces  $M$  of the type which appear in variational theories is made difficult by the fact that the critical limits, such as absolute minima, relative minima, minimax values etc., are in general infinite in number. These limits are associated with relative  $k$ -cycles of various dimensions and are classified as 0-limits, 1-limits etc. The number of  $k$ -limits suitably counted is called the  $k^{\text{th}}$  type number  $m_k$  of  $F$ . The theory seeks to establish relations between the numbers  $m_k$  and the connectivities  $p_k$  of  $M$ . The numbers  $p_k$  are finite in the most important applications. It is otherwise with the numbers  $m_k$ .

The theory has been able to proceed provided one of the following hypotheses is satisfied. The critical limits cluster at most at  $+\infty$ ; the critical points are isolated;<sup>1</sup> the problem is defined by analytic functions; the critical limits taken in their natural order are well-ordered. These conditions are not generally

# When was persistent homology invented first?

Web Images More... Sign in

Google Scholar 9 results (0.02 sec) My Citations ▾

All citations Articles Case law My library

Any time Since 2016 Since 2015 Since 2012 Custom range...

Sort by relevance Sort by date

include citations  Create alert

**Rank and span in functional topology**

Search within citing articles

**Exact homomorphism sequences in homology theory** ed.ac.uk [PDF]

JL Kelley, E Pitcher - Annals of Mathematics, 1947 - JSTOR

The developments of this paper stem from the attempts of one of the authors to deduce relations between homology groups of a complex and homology groups of a complex which is its image under a simplicial map. Certain relations were deduced (see [EP 1] and [EP 2] ...)

Cited by 46 Related articles All 3 versions Cite Save More

**Marston Morse and his mathematical works** ams.org [PDF]

R Bott - Bulletin of the American Mathematical Society, 1980 - ams.org

American Mathematical Society. Thus Morse grew to maturity just at the time when the subject of Analysis Situs was being shaped by such masters2 as Poincaré, Veblen, LEJ Brouwer, GD Birkhoff, Lefschetz and Alexander, and it was Morse's genius and destiny to ...

Cited by 24 Related articles All 4 versions Cite Save More

**Unstable minimal surfaces of higher topological structure**

M Morse, CB Tompkins - Duke Math. J., 1941 - projecteuclid.org

1. Introduction. We are concerned with extending the calculus of variations in the large to multiple integrals. The problem of the existence of minimal surfaces of unstable type contains many of the typical difficulties, especially those of a topological nature. Having studied this ...

Cited by 19 Related articles All 2 versions Cite Save

**[PDF] Persistence in discrete Morse theory** psu.edu [PDF]

U Bauer - 2011 - Citeseer

The goal of this thesis is to bring together two different theories about critical points of a scalar function and their relation to topology: Discrete Morse theory and Persistent homology. While the goals and fundamental techniques are different, there are certain ...

Cited by 8 Related articles All 4 versions Cite Save More

# When was persistent homology invented first?

BULLETIN (New Series) OF THE  
AMERICAN MATHEMATICAL SOCIETY  
Volume 3, Number 3, November 1980

## MARSTON MORSE AND HIS MATHEMATICAL WORKS

BY RAOUL BOTT<sup>1</sup>

**1. Introduction.** Marston Morse was born in 1892, so that he was 33 years old when in 1925 his paper *Relations between the critical points of a real-valued function of n independent variables* appeared in the Transactions of the American Mathematical Society. Thus Morse grew to maturity just at the time when the subject of Analysis Situs was being shaped by such masters<sup>2</sup> as Poincaré, Veblen, L. E. J. Brouwer, G. D. Birkhoff, Lefschetz and Alexander, and it was Morse's genius and destiny to discover one of the most beautiful and far-reaching relations between this fledgling and Analysis; a relation which is now known as *Morse Theory*.

In retrospect all great ideas take on a certain simplicity and inevitability, partly because they shape the whole subsequent development of the subject. And so to us, today, Morse Theory seems natural and inevitable. However one only has to glance at these early papers to see what a tour de force it was in the 1920's to go from the mini-max principle of Birkhoff to the Morse inequalities, let alone extend these inequalities to function spaces, so that by the early 30's Morse could establish the theorem that for any Riemann

## When was persistent homology invented first?

*inequalities pertain between the dimensions of the  $A_i$  and those of  $H(A_i)$ .* Thus the Morse inequalities already reflect a certain part of the “Spectral Sequence magic”, and a modern and tremendously general account of Morse’s work on rank and span in the framework of Leray’s theory was developed by Deheuvels [D] in the 50’s.

Unfortunately both Morse’s and Deheuvel’s papers are not easy reading. On the other hand there is no question in my mind that the papers [36] and [44] constitute another tour de force by Morse. Let me therefore illustrate rather than explain some of the ideas of the rank and span theory in a very simple and tame example.

In the figure which follows I have drawn a homeomorph of  $M = S^1$  in the plane, and I will be studying the height function  $F = y$  on  $M$ .






FIGURE 8

## Motivation and application: minimal surfaces






(from Dierkes et al.: *Minimal Surfaces*, 2010)

## Motivation and application: minimal surfaces



(from Dierkes et al.: *Minimal Surfaces*, 2010)

## Motivation and application: minimal surfaces

 $\Gamma_1$  $\Gamma_2$  $\Gamma_3$  $\dots$ 

(from Dierkes et al.: *Minimal Surfaces*, 2010)

## Existence of unstable minimal surfaces

Using persistent homology:

- Number of  $\epsilon$ -persistent critical points (minimal surfaces) is finite for any  $\epsilon > 0$
- Morse inequalities for  $\epsilon$ -persistent critical points

Theorem (Morse, Tompkins 1939)

*There is a  $C_1$  curve bounding an unstable minimal surface (an index 1 critical point of the area functional).*

# Computation

## Vietoris–Rips complexes

Consider a finite metric space  $(X, d)$ .

The *Vietoris–Rips complex* is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid \text{diam } S \leq t\}$$

- all edges with pairwise distance  $\leq t$
- all possible higher simplices

## Vietoris–Rips complexes

Consider a finite metric space  $(X, d)$ .

The *Vietoris–Rips complex* is the simplicial complex

$$\text{Rips}_t(X) = \{S \subseteq X \mid \text{diam } S \leq t\}$$

- all edges with pairwise distance  $\leq t$
- all possible higher simplices

For large  $t$ ,  $\text{Rips}_t(X)$  is the full simplex with vertices  $X$

- Number of  $d$ -simplices is  $\binom{|X|}{d+1}$
- Computation is one of the most important challenges in applied topology!

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB

Demo: [live.ripser.org](http://live.ripser.org)

## An example computation

Example data set:

- 192 points on  $\mathbb{S}^2$
- persistent homology barcodes up to dimension 2
- over 56 mio. simplices in 3-skeleton

Some previous software:

- javaplex (Stanford): 3200 seconds, 12 GB
- Dionysus (Duke): 615 seconds, 3.4 GB
- DIPHA (IST Austria): 50 seconds, 6 GB
- GUDHI (INRIA): 60 seconds, 3 GB


Demo: [live.ripser.org](http://live.ripser.org)

- Ripser: 1.2 seconds, 160 MB

## A software for computing Vietoris–Rips persistence barcodes

- around 1000 lines of C++ code, no external dependencies
- support for
  - coefficients in a prime field  $\mathbb{Z}/p\mathbb{Z}$
  - sparse distance matrices (for distance threshold)
- open source (<http://ripser.org>)
- online version (<http://live.ripser.org>)
- 2016 ATMCS Best New Software Award (joint with RIVET by M. Lesnick and M. Wright)

Thanks for your attention!



Ripser Live users from 770 locations