


## 22-R-VIB-TW-49



A new fish species discovered in the Fraser River has been observed to be attracted to swinging pendulums as depicted above. You want to observe this phenomenon so you take a spherical metal ball of radius  $r = 0.1$  m and mass 4 kg attached to a rope of length  $L = 1.5$  m and tie it to a buoy with a servomotor driving the pendulum with a moment of  $M(t) = 7 \cos(5t)$ . If the drag force of the ball is modelled by  $|F_d| = 4v$ , what is the amplitude of the pendulum's oscillations? (Use  $g = 9.81$  m/s<sup>2</sup> and assume that  $\sin \theta = \theta$ )

### Solution:

Let's begin by writing the sum of moments to get a differential equation in the form of Hooke's law

$$\sum M_A : I_A \alpha = -cvL - mgL \sin \theta + M_0 \cos(\omega_0 t)$$

$$I_A = \frac{2}{5}mr^2 + mL^2 = 9.016 \text{ [kg} \cdot \text{m}^2]$$

$$v = L\dot{\theta}$$

$$\sin \theta = \theta, \quad \alpha = \ddot{\theta}$$

$$I_A \ddot{\theta} + cL^2 \dot{\theta} + mgL\theta = M_0 \cos(\omega_0 t)$$

Skipping the derivation, we can use a given formula (from the textbook) to express the amplitude:

$$X' = \frac{F_0/k}{\sqrt{(1 - (\frac{\omega_0}{\omega_n})^2)^2 + (2(\frac{c}{c_c})(\frac{\omega_0}{\omega_n}))^2}}$$

Replacing the values of  $k$ ,  $m$ , and  $c$  with the corresponding values in our ODE, we get

$$\begin{aligned}
 \omega_n &= \sqrt{\frac{mgL}{I_A}} = 2.56 \text{ [rad]} \\
 c_c &= 2I_A\omega_n = 46 \text{ [kg} \cdot \text{rad} \cdot \text{m}^2\text{]} \\
 X' &= \frac{F_0/(mgL)}{\sqrt{(1 - (\frac{\omega_0}{\omega_n})^2)^2 + (2(\frac{c}{c_c})(\frac{\omega_0}{\omega_n}))^2}} = \frac{7/((4)(9.81)(1.5))}{\sqrt{(1 - (\frac{5}{2.56})^2)^2 + (2(\frac{4}{46})(\frac{5}{2.56}))^2}} \\
 &= 0.0417 \text{ [rad]}
 \end{aligned}$$