

Biochar Research at Russell Ranch


Deirdre E. Griffin¹, Kate M. Scow¹,
Emma Torbert², Tom Hollemans³, Israel Herrera²
¹Dept. of Land, Air and Water Resources, UC Davis
²Agricultural Sustainability Institute, UC Davis
³Wageningen University, Netherlands, Visiting Scholar

UCDAVIS
UNIVERSITY OF CALIFORNIA

Department of
LAND, AIR AND WATER RESOURCES
University of California, Davis
Climate Change - Sustainable Agriculture
Environmental Quality - Landscape Processes

How is biochar made?

- Plant residues or manure are burned at 200-900°C under either no O₂ (pyrolysis) or low O₂ (gasification).
- These processes yield a char product that can be applied to soil.
- Oil and gas by-products can also be used as fuel.



What are the properties of biochar?


- Biochar's properties can vary greatly depending on feedstock and production method
- However, common characteristics that may make it beneficial to the soil include:
 - High surface area
 - Alkaline pH
 - Stable C structures
 - High cation exchange capacity

Russell Ranch Biochar Experimental Plots

- Established in Spring 2012 and planted in tomatoes
- Used walnut biochar from Dixon Ridge Farms, Winters, CA
- 4 treatments replicated 4 times

The experimental treatments were:

Biochar + Mineral Fertilizer	Compost	Mineral Fertilizer	Biochar + Compost
4.5 tons/acre biochar; 145:45:45 lb/acre N:P:K	4.5 tons/acre	145:45:45 lb/acre N:P:K	4.5 tons/acre of each



Mean ± SD. Different letters indicate statistically significant differences.