Effects of Cover Crops on a Northern California Vineyard Ecosystem

Chuck Ingels, UCCE Sacramento County
 Kate Scow, LAWR Dept., UCD
 Desley Whisson, WFC Biology Dept., UCD
 Laura Tourte, UCCE Santa Cruz County

Funding

UC Sustainable Agriculture
Research & Education Program

Lodi-Woodbridge Winegrape Commission

Experimental Methods Vineyard

- Merlot on 5BB, eastern Sac. County
- 2.1 x 3.3 m (7 x 11 ft.)
- Planted 1993
- Drip irrigated, fertigated

Experimental Methods Cover Crop Mixes

Fall 1996:

- Calif. native perennial grass (NT)
 Fall 1997:
- Annual clover (NT)Fall 1997-99
- Bell bean / vetch / pea / oat (D)
- Barley / oat (D)
- Disked control (D)

Experimental Methods Cover Crop Fertilization

45 kg/ha N/year (urea):

- Calif. native perennial grass
- Barley / oat
 No additional fertilizer:
- Annual clover
- Bell bean / vetch / pea / oat
- Disked control


















Results (3 Year Study)


Little or no effect on:

- Yield
- Juice quality
- Vine water stress

Cover Crop N Content 1998-2000




Cover Crop Biomass April, 1998




 $(MT/ha \times 0.46 = tons/acre)$

(LSD, p > 0.05)

Cover Crop N Content April, 1998




Weed Biomass






Pocket Gophers 1999




Soil Microbial Biomass



Wine Quality



Cover Crop Costs per Acre (Planted Every Row)




Are Cover Crops Sustainable?


It depends:

- Source of organic N
- Erosion protection
- Water penetration
- Vigor management, wine quality
- Do benefits outweigh costs?
- Resident vegetation may be as good

Veraison Petiole Nitrate 1998



Bloom Petiole Nitrate 2001 (Post Experiment)



Approximate Cover Crop Costs (Yr. 1 and Yr. 2+)

	Disk		Plant		Mow/Chop	Fert.
Nat Gr	2	0	Y	N	2	Y
BB/Vet	6	6	Y	Y	1	N
Clover	2	0	Y	N	3	N
Cereal	6	6	Y	Y	2	Y
Disked	5	5	N	N	1	N
Res Veg	0	0	N	N	3	N