
AIR POLLUTION

Latino Policy & Politics Institute UCLA Climate & Health Dashboard

San Joaquin County

County Statistics

Factors Influencing Exposure to Air Pollution



Age Latino: 28 yrs NL white: 47 yrs **Noncitizen Population**

Latino: 18% NI white: 2%

Limited English Proficiency

Latino: 26% NL white: 1%

Households

Latino: 48% NL white: 31%

Rate

Latino: 14% NL white: 10%



Median Income (Household)

Latino: \$74k NL white: \$91k



Benefits

Latino: 22% NL white: 8%



Insecurity Latino: 19% NL white: 9%



Latino: 10% NL white: 4%



Health Status Latino: 21%

NL white: 14%




Expectancy

Latino: 79 yrs NL white: 76 yrs

Neighborhood Statistics

Air Pollutants

Latino Neighborhoods and Exposure to Particulate Matter 2.5 (PM2.5), 2015-2017



Note: $\mu g/m^3$ = one-millionth of a gram per cubic meter of air. Note: California's state standard for PM2.5 is an annual average of 12 μ g/m³, while the federal standard is 9 μ g/m³. There is no state or federal or state standard for Diesel PM.

PM2.5

PM2.5 is produced from sources like vehicle exhaust, wildfires, and industrial activity. These fine air particles enter the lungs and bloodstream and worsen conditions like asthma and heart disease.

Latino neighborhoods had similar exposure to PM2.5 as NL white neighborhoods.

> 11 μg/m³ Latino neighborhoods

10 μg/m³

NL white neighborhoods

Annual mean concentration

Diesel PM

Diesel emissions from vehicles and heavy-duty equipment release harmful particulate matter. Exposure to diesel exhaust can raise blood pressure, trigger heart attacks, and worsen lung conditions.

Latino neighborhoods had higher exposure to diesel PM than NL white neighborhoods.

> 0.31 tons/year Latino neighborhoods

0.12 tons/year NL white neighborhoods

Emissions

Latino neighborhoods = Census tracts with 70%+ Latino residents NL white neighborhoods = Census tracts with 70%+ NL white residents

^{*}NL white = Non-Latino white

Latino Policy & Politics Institute UCLA Climate & Health Dashboard

Neighborhood Statistics (cont.)

Proximity to Major Sources of Air Pollution

Note: Exposure and proximity scores take into account the number of sites/facilities and their proximity to neighborhoods. Higher scores = more exposure to pollutants for residents.

Cleanup sites, such as Superfunds, are polluted with materials like lead and asbestos. Examples include old and abandoned processing plants and manufacturing facilities.

Exposure Score

Latino neighborhoods **NL** white neighborhoods treatment, storage, and disposal sites. They can release toxic substances such as carcinogens, mercury, and asbestos into the air, water, and soil.

Hazardous waste facilities are

Exposure Score

0.3 0.3

Latino neighborhoods NL white neighborhoods

RMP facilities are sites where hazardous chemicals—like propane, pesticides, ammonia, and explosives—are present, posing risks to the environment and communities if released.

Proximity Score

2.4 0.9

Latino neighborhoods **NL white** neighborhoods

Vehicle Types and Traffic

Lower-emission vehicles (LEVs)

use battery electric, plug-in hybrid, or hybrid technology to reduce greenhouse gas emissions.

% of LEVs owned

2% 4%

Latino neighborhoods **NL** white neighborhoods of pollutants because they lack advanced emission-control equipment.

Clunker vehicles (vehicles 20

years or older) emit high levels

% of clunker vehicles owned

16% 12%

Latino neighborhoods **NL** white neighborhoods vehicles on roads within an area. Neighborhoods near major roadways face greater exposure to harmful emissions released from vehicles.

Traffic density measures the concentration of

Vehicle kilometers per hour

856 km/hr 401 km/hr

Latino neighborhoods NL white neighborhoods

Vulnerable Groups

Age

Children and older adults are more vulnerable to air pollution and have a higher risk of developing respiratory and cardiovascular diseases.

8% ages 0-5

Latino neighborhoods

9% ages 65+ 5%

20%

ages 0-5 ages 65+

NL white neighborhoods

Health

Air pollution worsens pre-existing health conditions like asthma and coronary heart disease, increasing emergency visits and health complications. Long-term exposure to air pollution can cause chronic illness and premature death.

% of Adults (18+) with Pre-Existing Conditions

6% Latino

NL white neighborhoods neighborhoods

Coronary Heart Disease

11% 10% NL white Latino

neighborhoods neighborhoods

Asthma

Low Birth Weight (LBW) Babies

LBW babies are born under 5 lbs. LBW increases the risk of infant mortality, developmental delays, and chronic health conditions. Exposure to air pollution, such as PM2.5, contributes to higher rates of LBW

babies.

% of Infants

6%

Latino neighborhoods

4%

NL white neighborhoods

Emergency Department Visits (per 10,000 people)

21

15

Latino **NL** white neighborhoods neighborhoods

Heart Attacks

86 Latino

43 **NL** white neighborhoods neighborhoods

Asthma Attacks

Disadvantaged Communities

The CA Environmental Protection Agency defines disadvantaged communities based on their environmental pollution burden and population characteristics. Under Senate Bill 535, revenue from CA's Cap-and-Trade Program is partly directed toward these communities through the CA Climate Investments program to reduce pollution, enhance climate resilience, and improve health and economic well-being.

% of Disadvantaged Communities

100%

Latino neighborhoods

NL white neighborhoods

14%