
# AIR POLLUTION

# UCLA Latino Policy & Politics Institute Climate & Health Dashboard

# **Madera County**

# **County Statistics**

## **Factors Influencing Exposure to Air Pollution**



Median
Age
Latino: 28 yrs
NL white: 50 yrs

Noncitizen Population

Latino: 21% NL white: 1% AZ Limited English

Proficiency Latino: 30%

Latino: 30% NL white: 1% A

Households Latino: 46% NL white: 21% H

Rate

Latino: 24% NL white: 12%



Median Income (Household)

Latino: \$65k NL white: \$81k



SNAP Benefits

Latino: 27% NL white: 11%



Insecurity
Latino: 23%
NL white: 11%



Latino: 10% NL white: 4%



Latino: 24% NL white: 15%




Expectancy
Latino: 78 yrs
NL white: 77 yrs

# **Neighborhood Statistics**

**Air Pollutants** 

# Latino Neighborhoods and Exposure to Particulate Matter 2.5 (PM2.5), 2015-2017



Note:  $\mu g/m^3$  = one-millionth of a gram per cubic meter of air Note: California's state standard for PM2.5 is an annual average of 12  $\mu g/m^3$ , while the federal standard is 9  $\mu g/m^3$ . There is no state or federal or state standard for Diesel PM.

### PM2.5

**PM2.5** is produced from sources like vehicle exhaust, wildfires, and industrial activity. These fine air particles enter the lungs and bloodstream and worsen conditions like asthma and heart disease.

Latino neighborhoods had <u>similar exposure</u> to PM2.5 as NL white neighborhoods.

13 μg/m³ Latino neighborhoods **9** μg/m³

NL white neighborhoods

Annual mean concentration

### **Diesel PM**

**Diesel emissions** from vehicles and heavy-duty equipment release harmful particulate matter. Exposure to diesel exhaust can raise blood pressure, trigger heart attacks, and worsen lung conditions.

Latino neighborhoods had <u>higher exposure</u> to diesel PM, a higher exposure than NL white neighborhoods.

**0.21** tons/year Latino neighborhoods

**0.01** tons/year NL white neighborhoods

Emissions

Latino neighborhoods = Census tracts with 70%+ Latino residents NL white neighborhoods = Census tracts with 70%+ NL white residents

<sup>\*</sup>NL white = Non-Latino white

#### Latino Policy & Politics Institute UCLA Climate & Health Dashboard

# **Neighborhood Statistics (cont.)**

## **Proximity to Major Sources of Air Pollution**

Note: Exposure and proximity scores take into account the number of sites/facilities and their proximity to neighborhoods. Higher scores = more exposure to pollutants for residents.

Cleanup sites, , such as Superfunds, are polluted with materials like lead and asbestos. Examples include old and abandoned processing plants and manufacturing facilities.

#### **Exposure Score**

Latino neighborhoods **NL** white neighborhoods

#### Hazardous waste facilities are

treatment, storage, and disposal sites. They can release toxic substances such as carcinogens, mercury, and asbestos into the air, water, and soil.

#### **Exposure Score**

0.3 0.2

Latino neighborhoods NL white neighborhoods

RMP facilities are sites where hazardous chemicals—like propane, pesticides, ammonia, and explosives—are present, posing risks to the environment and communities if released.

#### **Proximity Score**

0.7 0.0

Latino neighborhoods **NL white** neighborhoods

## **Vehicle Types and Traffic**

#### Lower-emission vehicles (LEVs)

use battery electric, plug-in hybrid, or hybrid technology to reduce greenhouse gas emissions.

% of LEVs owned

2% 5%

Latino neighborhoods

**NL** white neighborhoods

**Clunker vehicles** (vehicles 20 years or older) emit high levels of pollutants because they lack advanced emission-control equipment.

% of clunker vehicles owned

13% 16%

Latino neighborhoods **NL** white neighborhoods vehicles on roads within an area. Neighborhoods near major roadways face greater exposure to harmful emissions released from vehicles.

Traffic density measures the concentration of

#### Vehicle kilometers per hour

3%

821 km/hr 397 km/hr

Latino neighborhoods NL white neighborhoods

## **Vulnerable Groups**

Age

Children and older adults are more vulnerable to air pollution and have a higher risk of developing respiratory and cardiovascular diseases.

8% ages 0-5

Latino neighborhoods

9%

ages 65+

30% ages 65+

ages 0-5 NL white neighborhoods

Health

Air pollution worsens pre-existing health conditions like asthma and coronary heart disease, increasing emergency visits and health complications. Long-term exposure to air pollution can cause chronic illness and premature death.

#### % of Adults (18+) with Pre-Existing Conditions

5% Latino

**NL** white neighborhoods neighborhoods

Coronary Heart Disease

11% Latino

NL white neighborhoods neighborhoods

**Asthma** 

10%

#### Low Birth Weight (LBW) Babies

LBW babies are born under 5 lbs. LBW increases the risk of infant mortality, developmental delays, and chronic health conditions. Exposure to air pollution, such as PM2.5, contributes to higher rates of LBW

babies.

% of Infants

5%

Latino neighborhoods

4%

**NL** white neighborhoods

#### Emergency Department Visits (per 10,000 people)

**31** 

Latino **NL** white neighborhoods neighborhoods

**Heart Attacks** 

94 30 Latino **NL** white neighborhoods neighborhoods

Asthma Attacks

## Disadvantaged Communities

The CA Environmental Protection Agency defines disadvantaged communities based on their environmental pollution burden and population characteristics. Under Senate Bill 535, revenue from CA's Cap-and-Trade Program is partly directed toward these communities through the CA Climate Investments program to reduce pollution, enhance climate resilience, and improve health and economic well-being.

% of Disadvantaged Communities

100%

Latino neighborhoods

0% **NL** white neighborhoods