
# AIR POLLUTION

# UCLA Latino Policy & Politics Institute Climate & Health Dashboard

# San Bernardino County

## **County Statistics**

## **Factors Influencing Exposure to Air Pollution**



Median Age Latino: 29 NL white: 45 Noncitizen

Population
Latino: 14%
NI. white: 2%

AZ mited End

Limited English Proficiency

Latino: 21% NL white: 2%

Households Latino: 42% NL white: 31% H

Rate
Latino: 15%

NL white: 10%

Median Income
(Household)

Latino: \$75k NL white: \$83k



SNAP Benefits

Latino: 17% NL white: 10%



Insecurity
Latino: 18%
NL white: 10%

Uninsured Rate

Latino: 11% NL white: 5% Fair/Poor

Health Status Latino: 17%

NL white: 13%




Expectancy
Latino: 78 yrs
NL white: 75 yrs

# **Neighborhood Statistics**

#### **Air Pollutants**

# Latino Neighborhoods and Exposure to Particulate Matter 2.5 (PM2.5), 2015–2017



Note: µg/m³ = one-millionth of a gram per cubic meter of air

Note: California's state standard for PM2.5 is an annual average of 12 µg/m³, while the federal standard is 9 µg/m³. There is no state or federal or state standard for Diesel PM.

#### PM2.5

**PM2.5** is produced from sources like vehicle exhaust, wildfires, and industrial activity. These fine air particles enter the lungs and bloodstream and worsen conditions like asthma and heart disease.

Latino neighborhoods had <u>higher exposure</u> to PM2.5 than NL white neighborhoods.

12 μg/m³ Latino neighborhoods 8 μg/m³

ods **NL white** neighborhoods

Annual mean concentration

#### **Diesel PM**

**Diesel emissions** from vehicles and heavy-duty equipment release harmful particulate matter. Exposure to diesel exhaust can raise blood pressure, trigger heart attacks, and worsen lung conditions.

Latino neighborhoods had <u>higher exposure</u> to diesel PM than NL white neighborhoods.

**0.27** tons/year Latino neighborhoods

0.03 tons/year
NL white neighborhoods

**Emissions** 

Latino neighborhoods = Census tracts with 70%+ Latino residents NL white neighborhoods = Census tracts with 70%+ NL white residents

<sup>\*</sup>NL white = Non-Latino white

#### Latino Policy & Politics Institute UCLA Climate & Health Dashboard

# **Neighborhood Statistics (cont.)**

## **Proximity to Major Sources of Air Pollution**

Note: Exposure and proximity scores take into account the number of sites/facilities and their proximity to neighborhoods. Higher scores = more exposure to pollutants for residents.

Cleanup sites, such as Superfunds, are polluted with materials like lead and asbestos. Examples include old and abandoned processing plants and manufacturing facilities.

#### **Exposure Score**

10

Latino neighborhoods

**NL** white neighborhoods

#### Hazardous waste facilities are

treatment, storage, and disposal sites. They can release toxic substances such as carcinogens, mercury, and asbestos into the air, water, and soil.

#### **Exposure Score**

0.5 0.2

Latino neighborhoods **NL** white neighborhoods

RMP facilities are sites where hazardous chemicals—like propane, pesticides, ammonia, and explosives—are present, posing risks to the environment and communities if released.

#### **Proximity Score**

1.2 0.1

Latino neighborhoods **NL white** neighborhoods

## **Vehicle Types and Traffic**

#### Lower-emission vehicles (LEVs)

use battery electric, plug-in hybrid, or hybrid technology to reduce greenhouse gas emissions.

% of LEVs owned

2%

Latino neighborhoods **NL** white neighborhoods

**Clunker vehicles** (vehicles 20 years or older) emit high levels of pollutants because they lack advanced emission-control equipment.

% of clunker vehicles owned

11% 13%

Latino neighborhoods **NL** white neighborhoods vehicles on roads within an area. Neighborhoods near major roadways face greater exposure to harmful emissions released from vehicles.

5%

ages 0-5

#### Vehicle kilometers per hour

**Traffic density** measures the concentration of

994 km/hr 375 km/hr

Latino neighborhoods **NL** white neighborhoods

### **Vulnerable Groups**

#### Age

Children and older adults are more vulnerable to air pollution and have a higher risk of developing respiratory and cardiovascular diseases.

7% ages 0-5 9%

ages 65+ Latino neighborhoods

26% ages 65+

NL white neighborhoods

#### Health

Air pollution worsens pre-existing health conditions like asthma and coronary heart disease, increasing emergency visits and health complications. Long-term exposure to air pollution can cause chronic illness and premature death.

#### % of Adults (18+) with Pre-Existing Conditions

5% Latino

**NL** white neighborhoods neighborhoods

**Coronary Heart Disease** 

10% Latino

10% NL white

neighborhoods neighborhoods

**Asthma** 

#### Emergency Department Visits (per 10,000 people)

19

19

Latino **NL** white neighborhoods neighborhoods

**Heart Attacks** 

68

51

Latino **NL** white neighborhoods neighborhoods

Asthma Attacks

#### Low Birth Weight (LBW) Babies

LBW babies are born under 5 lbs. LBW increases the risk of infant mortality, developmental delays, and chronic health conditions. Exposure to air pollution, such as PM2.5, contributes to

higher rates of LBW

babies.

% of Infants

6%

Latino neighborhoods

\*\*\*\*

5%

**NL** white neighborhoods

### Disadvantaged Communities

The CA Environmental Protection Agency defines disadvantaged communities based on their environmental pollution burden and population characteristics. Under Senate Bill 535, revenue from CA's Cap-and-Trade Program is partly directed toward these communities through the CA Climate Investments program to reduce pollution, enhance climate resilience, and improve health and economic well-being.

## % of Disadvantaged Communities

87%

Latino

neighborhoods

4% **NL** white neighborhoods