

CSE 150A-250A AI: Probabilistic Methods

Lecture 5

Fall 2026

Trevor Bonjour
Department of Computer Science and Engineering
University of California, San Diego

Slides adapted from previous versions of the course (Prof. Lawrence, Prof. Alvarado, Prof Berg-Kirkpatrick)

Agenda

Review

Inference

Exact Inference: Variable Elimination

Review


d-separation and conditional independence

- Theorem

$P(X, Y|E) = P(X|E)P(Y|E)$ if and only if every *path* from a node in X to a node in Y is *blocked* by E .

- Definition

A path π is **blocked** if there exists a node $Z \in \pi$ for which one of three conditions holds:



$\text{descendants}(Z) \cap E = \emptyset$

A **descendant** of Z is *any* node (e.g., child, grandchild) that lies on a directed path from Z .

D-Separation Algorithm

1. Shade all observed nodes $\{Z_1, \dots, Z_k\}$ in the graph.
2. Enumerate all undirected paths from X to Y .
3. For each path:
 - 3.1 Decompose the path into triples (segments of 3 nodes).
 - 3.2 If none of the d-separation blocking conditions apply to any of the triples on the path, then the path is **active** and **d-connects** X and Y . Return $X \not\perp\!\!\!\perp Y \mid \{Z_1, \dots, Z_k\}$
4. If all paths are blocked , then

$$X \perp\!\!\!\perp Y \mid \{Z_1, \dots, Z_k\}.$$

Loopy example

A. TRUE or B. FALSE?

5. $P(B|D, E) \stackrel{?}{=} P(B|D)$



The evidence is $\{D\}$.

There are two paths from B to E .

Path $B \rightarrow D \rightarrow E$
is blocked by node D ,
satisfying condition (1).


Path $B \rightarrow D \leftarrow A \rightarrow C \rightarrow E$
is not blocked by any node.

The statement is .



Markov Blanket

A **Markov Blanket** B_X of node X consists of **parents** of X , **children** of X and **"spouses"** (other parents of children of X , but not X) of X .



Every variable is conditionally independent of any other variable given it's **Markov Blanket**.

Inference

- **Problem**

Given a set E of evidence nodes, and a set Q of query nodes, how to compute the posterior distribution $P(Q|E)$?

- **More precisely**

How to express $P(Q|E)$ in terms of the CPTs $P(X_i|pa(X_i))$ of the BN, which are assumed to be given?

- **Tools at our disposal**

Bayes rule

marginalization

product rule


marginal independence

conditional independence

Strategy to compute $P(Q|E)$

Bayes rule	Use to express $P(Q E)$ in terms of conditional probabilities that respect the order of the DAG.
marginalization	Use to introduce nodes on the left side of the conditioning bar when they need to appear as parents.
product rule	Use to express joint predictions (over multiple variables) in terms of simpler individual predictions.
marginal and conditional independence	Use to remove non-informative variables from the right side of the conditioning bar.

Inference Example




Q. What are the CPTs associated with the DAG shown above?


A. $P(B|J,M)$ B. $P(B)$ C. $P(A|B,E)$

D. A, B and C E. B and C

Inference example



Inference Example



$$P(B = 1 | J = 1, M = 1) = ??$$

Inference Example: Enumeration

$$P(B = 1|J = 1, M = 1)$$

$$= \frac{P(B = 1, J = 1, M = 1)}{P(J = 1, M = 1)}$$

$$= \alpha P(B = 1, J = 1, M = 1)$$

$$= \alpha \sum_e \sum_a P(B = 1, J = 1, M = 1, E = e, A = a)$$

$$= \alpha \sum_e \sum_a P(B = 1)P(E = e)P(A = a|B = 1, E = e)P(J = 1|A = a)P(M = 1|A = a)$$

Q. How many operations does it take to calculate $P(B = 1, J = 1, M = 1)$?

Inference Example: Enumeration

$$P(B = 1, J = 1, M = 1)$$

$$\begin{aligned} &= P(B = 1) \times P(E = 0) \times P(A = 0|B = 1, E = 0) \times P(J = 1|A = 0) \times P(M = 1|A = 0) \\ &+ P(B = 1) \times P(E = 0) \times P(A = 1|B = 1, E = 0) \times P(J = 1|A = 1) \times P(M = 1|A = 1) \\ &+ P(B = 1) \times P(E = 1) \times P(A = 0|B = 1, E = 1) \times P(J = 1|A = 0) \times P(M = 1|A = 0) \\ &+ P(B = 1) \times P(E = 1) \times P(A = 1|B = 1, E = 1) \times P(J = 1|A = 1) \times P(M = 1|A = 1) \end{aligned}$$

Inference Example: Enumeration

$$P(B = 1 \mid J = 1, M = 1) = \frac{P(B = 1, J = 1, M = 1)}{P(B = 0, J = 1, M = 1) + P(B = 1, J = 1, M = 1)}$$

Q. How many multiplications and additions does it take to compute $P(B = 1 \mid J = 1, M = 1)$?

- A. 16 multiplications, 3 additions
- B. 48 multiplications, 7 additions
- C. 32 multiplications, 6 additions
- D. 32 multiplications, 7 additions
- E. I have no idea

Inference Example: Enumeration

$$P(B = 1|J = 1, M = 1)$$

$$= \frac{P(B = 1, J = 1, M = 1)}{P(J = 1, M = 1)}$$

$$= \alpha P(B = 1, J = 1, M = 1)$$

$$= \alpha \sum_e \sum_a P(B = 1, J = 1, M = 1, E = e, A = a)$$

$$= \alpha \sum_e \sum_a P(B = 1)P(E = e)P(A = a|B = 1, E = e)P(J = 1|A = a)P(M = 1|A = a)$$

$$= \alpha P(B = 1) \sum_e P(E = e) \sum_a P(A = a|B = 1, E = e)P(J = 1|A = a)P(M = 1|A = a)$$

Q. How many operations does it take now to calculate $P(B = 1, J = 1, M = 1)$?

Inference Example: Enumeration

$$P(B = 1, J = 1, M = 1)$$

$$\begin{aligned} = & P(B = 1) \times \left[P(E = 0) \times (P(A = 0|B = 1, E = 0) \times P(J = 1|A = 0) \times P(M = 1|A = 0)) \right. \\ & + P(A = 1|B = 1, E = 0) \times P(J = 1|A = 1) \times P(M = 1|A = 1)) \\ & + P(E = 1) \times (P(A = 0|B = 1, E = 1) \times P(J = 1|A = 0) \times P(M = 1|A = 0)) \\ & \left. + P(A = 1|B = 1, E = 1) \times P(J = 1|A = 1) \times P(M = 1|A = 1)) \right] \end{aligned}$$

Inference Example: Enumeration

$$P(b|j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A|b, E) P(j|A) P(m|A)$$




Image Source: *Artificial Intelligence: A Modern Approach* (Russell & Norvig, 2020)

Inference Example: Enumeration

$$P(B = 1 \mid J = 1, M = 1) = \frac{P(B = 1, J = 1, M = 1)}{P(B = 0, J = 1, M = 1) + P(B = 1, J = 1, M = 1)}$$


Q. How many multiplications and additions does it take to compute $P(B = 1 \mid J = 1, M = 1)$ using this rearrangement?

- A. 11 multiplications, 3 additions
- B. 22 multiplications, 6 additions
- C. 22 multiplications, 7 additions
- D. 33 multiplications, 7 additions
- E. I have no idea

Exact Inference: Variable Elimination

- **Idea:** Eliminate **repeated calculations** by storing intermediate results in “factors”.
- A **factor** is a function of a set of variables that evaluates to a number.
- Variable Elimination (VE) successively eliminates all non-query variables, one at a time, until only factors involving the query variables remain.
- To eliminate a variable:
 - *join* all factors containing that variable.
 - *sum out* that variable from the resulting factor.

VE Example



$$P(B = 1 | J = 1, M = 1) = ??$$

$$\begin{aligned} P(B \mid J = 1, M = 1) \\ &= \alpha P(B, J = 1, M = 1) \\ &= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) P(M = 1 \mid A) \end{aligned}$$

$$\begin{aligned}P(B \mid J = 1, M = 1) \\&= \alpha P(B, J = 1, M = 1) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) P(M = 1 \mid A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) f_1(A)\end{aligned}$$

$$\begin{aligned}P(B \mid J = 1, M = 1) \\&= \alpha P(B, J = 1, M = 1) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) P(M = 1 \mid A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) f_1(A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) f_2(A)\end{aligned}$$

VE Example

$$\begin{aligned}P(B \mid J = 1, M = 1) &= \alpha P(B, J = 1, M = 1) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) P(M = 1 \mid A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) f_1(A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) f_2(A) \\&= \alpha P(B) \sum_E P(E) f_3(B, E)\end{aligned}$$

$$\begin{aligned}P(B \mid J = 1, M = 1) &= \alpha P(B, J = 1, M = 1) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) P(M = 1 \mid A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) f_1(A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) f_2(A) \\&= \alpha P(B) \sum_E P(E) f_3(B, E) \\&= \alpha P(B) f_4(B)\end{aligned}$$

VE Example

$$\begin{aligned}P(B \mid J = 1, M = 1) \\&= \alpha P(B, J = 1, M = 1) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) P(M = 1 \mid A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) f_1(A) \\&= \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) f_2(A) \\&= \alpha P(B) \sum_E P(E) f_3(B, E) \\&= \alpha P(B) f_4(B) \\&= \alpha f_5(B)\end{aligned}$$

$$P(B \mid J = 1, M = 1) = \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) P(M = 1 \mid A)$$

A	$f_1(A) = P(M = 1 \mid A)$
0	0.01
1	0.7

VE Example

$$P(B \mid J = 1, M = 1) = \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) f_1(A)$$

A	$P(J = 1 \mid A)$
0	0.05
1	0.9

A	$f_1(A)$
0	0.01
1	0.7

A	$f_2(A) = P(J = 1 \mid A) f_1(A)$
0	
1	

VE Example

$$P(B \mid J = 1, M = 1) = \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(J = 1 \mid A) f_1(A)$$

A	$P(J = 1 \mid A)$
0	0.05
1	0.9

A	$f_1(A)$
0	0.01
1	0.7

A	$f_2(A) = P(J = 1 \mid A) f_1(A)$
0	0.05×0.01
1	0.9×0.7

VE Example

$$P(B \mid J = 1, M = 1) = \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) f_2(A)$$

A	$f_2(A)$
0	0.0005
1	0.63

B	E	$P(A = 1 \mid B, E)$
0	0	0.001
1	0	0.94
0	1	0.29
1	1	0.95

B	E	$f_3(B, E) = \sum_A P(A \mid B, E) f_2(A)$
0	0	
1	0	
0	1	
1	1	

VE Example

$$P(B \mid J = 1, M = 1) = \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) f_2(A)$$

A	$f_2(A)$
0	0.0005
1	0.63

B	E	$P(A = 1 \mid B, E)$
0	0	0.001
1	0	0.94
0	1	0.29
1	1	0.95

B	E	$f_3(B, E) = \sum_A P(A \mid B, E) f_2(A)$
0	0	$0.001 \times 0.63 + 0.999 \times 0.0005$
1	0	$0.94 \times 0.63 + 0.06 \times 0.0005$
0	1	$0.29 \times 0.63 + 0.71 \times 0.0005$
1	1	$0.95 \times 0.63 + 0.05 \times 0.0005$

VE Example

$$P(B \mid J = 1, M = 1) = \alpha P(B) \sum_E P(E) f_3(B, E)$$

E	$P(E)$
0	0.998
1	0.002

B	E	$f_3(B, E)$
0	0	0.0011295
1	0	0.59223
0	1	0.183055
1	1	0.598525

B	$f_4(B) = \sum_E P(E) f_3(B, E)$
0	$0.998 \times 0.0011295 + 0.002 \times 0.183055$
1	$0.998 \times 0.59223 + 0.002 \times 0.598525$

VE Example

$$P(B \mid J = 1, M = 1) = \alpha P(B) f_4(B)$$

B	$P(B)$
0	0.999
1	0.001

B	$f_4(B)$
0	0.001493351
1	0.59224259

B	$f_5(B) = P(B)f_4(B)$
0	0.999×0.001493351
1	0.001×0.59224259

$$P(B \mid J = 1, M = 1) = \alpha f_5(B)$$

B	$f_5(B)$
0	0.001491858
1	0.000592243

VE Example

$$P(B \mid J = 1, M = 1) = \alpha f_5(B)$$

B	$f_5(B)$
0	0.001491858
1	0.000592243

$$N = 0.001491858 + 0.000592243, \quad \alpha = \frac{1}{N}$$

B	$P(B \mid J = 1, M = 1)$
0	0.716
1	0.284

Q. How many multiplications and additions does it take to compute $P(B = 1 \mid J = 1, M = 1)$ using VE?

$$f_1(A) = P(M = 1 \mid A)$$

$$f_2(A) = P(J = 1 \mid A)f_1(A)$$

$$f_3(B, E) = \sum_A P(A \mid B, E)f_2(A)$$

$$f_4(B) = \sum_E P(E)f_3(B, E)$$

$$f_5(B) = P(B)f_4(B)$$

$$P(B \mid J = 1, M = 1) = \alpha f_5(B)$$

VE Example

Factor	Multiplications	Additions
$f_1(A)$	0	0
$f_2(A)$	2	0
$f_3(B, E)$	8	4
$f_4(B)$	4	2
$f_5(B)$	2	0
Total	16	6

Normalization: + 1 addition

16 multiplications, 7 additions

- Factors are usually represented as a table (therefore an arbitrary function)
- **Caution:** Factors can look like CPTs, and CPTs can be represented as factors, but factors are **not** necessarily probabilities!
- The values in factors only represent intermediate values in the calculations of some probability - with no real meaning in themselves.

Does order of elimination matter?

- In general, yes (but not in the trivial graphs we've been considering)
- Time and space of VE is dominated by the **largest** factor created
- **Heuristic:** Eliminate the variable that will lead to the smallest next factor being created
 - In a **polytree** this leads to **linear** time inference (in size of largest CPT).

Polytree: A belief network with at most one undirected path between any two nodes.

That's All Folks!