

# Problem Set 3 Solutions

Calvin Walker

1.

(a) Consider the M-projection with the factored approximation  $Q(X, Y) = Q(X)Q(Y)$ ,

$$\begin{aligned} D(P||Q) &= \sum_{X,Y} P(X, Y) \log \frac{P(X, Y)}{Q(X)Q(Y)} \\ &= \mathbb{E}_P[\log P(X, Y) - \log Q(X)Q(Y)] \\ &= \mathbb{E}_P[\log P(X, Y)] - \left( \mathbb{E}_p[\log Q(X)] + \mathbb{E}_p[\log Q(Y)] \right) \pm \log(P(X)P(Y)) \\ &= \mathbb{E}_p \left[ \log \frac{P(X, Y)}{P(X)P(Y)} \right] + \mathbb{E}_p \left[ \log \frac{P(X)}{Q(X)} \right] + \mathbb{E}_p \left[ \log \frac{P(Y)}{Q(Y)} \right] \end{aligned}$$

Letting  $Q_M^* = P(X)P(Y)$ ,

$$D(P||Q) = D(P||Q_M^*) + \mathbb{E}_p \left[ \log \frac{P(X)}{Q(X)} \right] + \mathbb{E}_p \left[ \log \frac{P(Y)}{Q(Y)} \right]$$

Thus,  $D(P||Q) \geq D(P||Q_M^*)$ , and  $Q_M^* = P(X)P(Y)$  minimizes the M-projection.

(b)

$$\begin{aligned} \theta^* &= \arg \max_{\theta} \prod_{i=1}^M Q(X^{(i)}; \theta) \\ &= \arg \min_{\theta} \sum_{i=1}^M -\log Q(X^{(i)}; \theta) \\ &= \arg \min_{\theta} \sum_{i=1}^M -\log Q(X^{(i)}; \theta) + P(X^{(i)}) \end{aligned}$$

So if the sample size  $M$  is significantly large,

$$\begin{aligned} \theta^* &= \arg \min_{\theta} \mathbb{E}_P \left[ \log \frac{P(X)}{Q(X; \theta)} \right] \\ &= \arg \min_{\theta} D(P||Q(X; \theta)) \end{aligned}$$

Therefore, the MLE solution  $\theta^*$  minimizes the KL-Divergence  $D(P||Q(X; \theta))$ , and is equivalent to solving for the M-projection  $D(P||Q)$ .

2.

(a)

(b)




Figure 1: Example Cluster Graph

Consider the above Cluster Graph for a pairwise MRF over three binary variables  $A$ ,  $B$ , and  $C$ . We can satisfy the local consistency constraints with the following clique beliefs:

$$\beta_1(A, B) = \begin{bmatrix} 0.3 & 0.2 \\ 0.2 & 0.2 \end{bmatrix} \quad \beta_2(B, C) = \begin{bmatrix} 0.3 & 0.2 \\ 0.2 & 0.3 \end{bmatrix} \quad \beta_3(A, C) = \begin{bmatrix} 0.2 & 0.3 \\ 0.3 & 0.2 \end{bmatrix}$$

Such that the sepset marginals are given by:  $\mu_{1,3}(A) = \mu_{1,2}(B) = \mu_{2,3}(C) = (0.5, 0.5)$ . However, if we sum over the possible outcomes:

$$\sum_{A,B,C} P(A, B, C) = 6 \left( \frac{18}{125} \right) + 2 \left( \frac{8}{125} \right) = \frac{124}{125} \neq 1$$

So the parameterization does not correspond to a valid probability distribution, and we can see for a cluster graph that is not a tree, the marginal polytope is strictly contained by the local consistency polytope.

**3.**

(a)