

Problem Set 3 Solutions

Calvin Walker

1.

- (a) Consider the M-projection with the factored approximation $Q(X, Y) = Q(X)Q(Y)$,

$$\begin{aligned} D(P||Q) &= \sum_{X,Y} P(X, Y) \log \frac{P(X, Y)}{Q(X)Q(Y)} \\ &= \mathbb{E}_P[\log P(X, Y) - \log Q(X)Q(Y)] \\ &= \mathbb{E}_P[\log P(X, Y)] - \left(\mathbb{E}_p[\log Q(X)] + \mathbb{E}_p[\log Q(Y)] \right) \pm \log(P(X)P(Y)) \\ &= \mathbb{E}_p \left[\log \frac{P(X, Y)}{P(X)P(Y)} \right] + \mathbb{E}_p \left[\log \frac{P(X)}{Q(X)} \right] + \mathbb{E}_p \left[\log \frac{P(Y)}{Q(Y)} \right] \end{aligned}$$

Letting $Q_M^* = P(X)P(Y)$,

$$D(P||Q) = D(P||Q_M^*) + \mathbb{E}_p \left[\log \frac{P(X)}{Q(X)} \right] + \mathbb{E}_p \left[\log \frac{P(Y)}{Q(Y)} \right]$$

Thus, $D(P||Q) \geq D(P||Q_M^*)$, and $Q_M^* = P(X)P(Y)$ minimizes the M-projection.

(b)

$$\begin{aligned} \theta^* &= \arg \max_{\theta} \prod_{i=1}^M Q(X^{(i)}; \theta) \\ &= \arg \min_{\theta} \sum_{i=1}^M -\log Q(X^{(i)}; \theta) \\ &= \arg \min_{\theta} \sum_{i=1}^M -\log Q(X^{(i)}; \theta) + P(X^{(i)}) \end{aligned}$$

So if the sample size M is significantly large,

$$\begin{aligned} \theta^* &= \arg \min_{\theta} \mathbb{E}_P \left[\log \frac{P(X)}{Q(X; \theta)} \right] \\ &= \arg \min_{\theta} D(P||Q(X; \theta)) \end{aligned}$$

Therefore, the MLE solution θ^* minimizes the KL-Divergence $D(P||Q(X; \theta))$, and is equivalent to solving for the M-projection $D(P||Q)$.

(c)

2.

- (a) $\mathcal{M} \subseteq \text{Local}[\mathcal{U}]$, since any valid distribution P over \mathcal{X} will satisfy the constraints of the local-consistency polytope. For a clique tree \mathcal{T} , under the constraints of the local-consistency polytope, the pseudo marginals must be locally consistent,

$$\mu_{i,j}(S_{i,j}) = \sum_{C_i - S_{i,j}} \beta_i(C_i) = \sum_{C_j - S_{i,j}} \beta_j(C_j)$$

which implies that the clique tree is calibrated. So we have the clique tree invariant

$$\tilde{P}_{\Phi}(\mathcal{X}) = \frac{\prod_i \beta_i(C_i)}{\prod_{i,j} \mu_{i,j}(S_{i,j})}$$

where $\beta_i(C_i) \propto \tilde{P}_{\Phi}(C_i)$ and $\mu_{i,j}(S_{i,j}) \propto \tilde{P}_{\Phi}(S_{i,j})$. Therefore, the clique and sepset marginals define a valid distribution over \mathcal{X} , and $\text{Local}[\mathcal{U}] \subseteq \mathcal{M}$. So we have that $\text{Local}[\mathcal{U}]$ is equivalent to \mathcal{M} for a clique tree \mathcal{T} .




Figure 1: Example Cluster Graph

(b)

Consider the above Cluster Graph for a pairwise MRF over three binary variables A , B , and C . We can satisfy the local consistency constraints with the following clique beliefs:

$$\beta_1(A, B) = \begin{bmatrix} 0.3 & 0.2 \\ 0.2 & 0.2 \end{bmatrix} \quad \beta_2(B, C) = \begin{bmatrix} 0.3 & 0.2 \\ 0.2 & 0.3 \end{bmatrix} \quad \beta_3(A, C) = \begin{bmatrix} 0.2 & 0.3 \\ 0.3 & 0.2 \end{bmatrix}$$

Such that the sepset marginals are given by: $\mu_{1,3}(A) = \mu_{1,2}(B) = \mu_{2,3}(C) = (0.5, 0.5)$. However, if we sum over the possible outcomes:

$$\sum_{A,B,C} P(A, B, C) = 6 \left(\frac{18}{125} \right) + 2 \left(\frac{8}{125} \right) = \frac{124}{125} \neq 1$$

So the parameterization does not correspond to a valid probability distribution, and we can see for a cluster graph that is not a tree, the marginal polytope is strictly contained by the local consistency polytope.

3.

(a)