

CS 754 : Advanced Image Processing Assignment 2

Meet Udeshi - 14D070007

Arka Sadhu - 140070011

February 10, 2017

Q2

A2.3

We have the relation :

$$E_u = \sum_{t=1}^T C_t \cdot F_t$$

Consider

$$E_1 = C_1 \cdot F_1$$

and suppose that we want to construct it as a matrix product, then we can write it as

$$E_1 = \phi_1 f_1$$

where $\phi_1 = \text{diag}(C_1)$ and $f_1 = \text{vec}(F_1)$ Hence

$$E_u = [\phi_1 | \phi_2 | \dots | \phi_T] \begin{bmatrix} f_1 \\ f_2 \\ \dots \\ f_T \end{bmatrix}$$

Therefore

$$x = \begin{bmatrix} f_1 \\ f_2 \\ \dots \\ f_T \end{bmatrix}$$
$$y = \text{vec}(Eu)$$
$$A = [\phi_1 | \phi_2 | \dots | \phi_T]$$

Q3

A3

Coherence is defined as :

$$\mu(\phi, \psi) = \sqrt{n} * \max_{i,j \in 0,1,\dots,n-1} |\phi^{i^t} \psi_j|$$

- (a) Upper Bound: Consider a row of ϕ matrix to be exactly same as one of the columns of ψ the inner product would be 1, because both matrices are unit normalized, and result in $\mu_{max}(\phi, \psi) = \sqrt{n}$.
- (b) Lower Bound: Let g be a row of the ϕ matrix, which is unit normalized. g can be written with basis vectors as columns of ψ . That is

$$g = \sum_{k=1}^n \alpha_k \psi_k$$

Also since g is unit norm,

$$\sum_{k=1}^n \alpha_k^2 = 1$$

When we take the inner product of g with ψ_k , only one α will remain. Now consider the coherence of g and ψ . Clearly

$$\mu(g, \psi) = \sqrt{n} * \max(\alpha_j)_{j=1}^n$$

where α_j is corresponds to the j th coefficient. Now we also know that

$$\max(\alpha_j)_{j=1}^n \geq \text{avg}(\alpha_j)_{j=1}^n$$

and equality occurs when all α_j are equal. Hence to get the minimum coherence we need all α_j to be equal and using the previous constraint on unit norm we get

$$\alpha_j = \frac{1}{\sqrt{n}}$$

. This gives

$$\mu_{\min}(g, \psi) = 1$$

This is the bound for all rows, and hence we have obtained

$$\mu_{\min}(\phi, \psi) = 1$$

Q4

A4