Ans to the ques no'- 1

Here,
$$A(0,0,4)$$
 is translated to $A_1^*(2,3,6)$

$$\begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} T_{\chi} \\ T_{y} \\ 4 + T_{z} \\ 1 \end{bmatrix}$$

So,
$$T_x = 2$$

 $T_y = 3$
 $T_z = 6-4=2$

i. Translation matrix =
$$\begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Activity of

and the

-

The same of the sa

1

III_

Following the translation matrix,

coordinate of
$$O_1: \begin{bmatrix} x' \\ y' \\ \hline z' \\ 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 2 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 2 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{cases} x' \\ y' \\ = \begin{cases} 2 \\ 3 \\ 2 \\ 1 \end{cases}$$

.. The new point is at O1 (2,3,2).

20

En

C. T.

二九

=-1

न त

Application of the last of the

_ त

E-a

The same of the sa

-4

Ans to the ques no :- 2

Considering the Center of Rotation to be at $O_1(2,3,2)$, we need to firstly shift the corner points in such way that the Center of Rotation lies at the Origin. Then, we need to rotate. Finally we will retranslate to original position.

Here, if O₁ is translated to origin, the whole Figure-2 will shift to be Figure-1. Now, we will notate Figure-1. The notated points 2 by 30°:

$$A_{2} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos 30^{\circ} & -\sin 30^{\circ} & 0 \\ 0 & \sin 630^{\circ} & \cos 30^{\circ} & 0 \end{bmatrix} = \begin{bmatrix} 0 \\ -2 \\ 2\sqrt{3} \\ 1 \end{bmatrix}$$

$$C_{2}^{1} = \begin{bmatrix} 1 & 0 & 0 & 0 & 2 \\ 0 & \cos 30^{\circ} & -\sin 30^{\circ} & 0 \\ 0 & \sin 30^{\circ} & \cos 30^{\circ} & 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 0 \\ 1 \end{bmatrix}$$

$$D_{2}^{1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos 30^{\circ} & -\sin 30^{\circ} & 0 \\ 0 & \sin 30^{\circ} & \cos 30^{\circ} & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \\ 4 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -2+\sqrt{3} \\ 1+2\sqrt{3} \\ 1 \end{bmatrix}$$

$$O_{2}^{\prime} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos 30^{\circ} & -\sin 30^{\circ} & 0 \\ 0 & \sin 30^{\circ} & \cos 30^{\circ} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

$$F_{2} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos 30^{\circ} & -\sin 30^{\circ} & 0 \\ 0 & \sin 30^{\circ} & \cos 30^{\circ} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$G_{12} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos 30^{\circ} & -\sin 30^{\circ} & 0 \\ 0 & \sin 30^{\circ} & \cos 30^{\circ} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 0 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ \sqrt{3} \\ 1 \end{bmatrix}$$

$$E_{2}^{1} = \begin{cases} 0 & 0 & 0 & 0 \\ 0 & \cos 30^{\circ} & -\sin 30^{\circ} & 0 \\ 0 & \sin 30^{\circ} & \cos 30^{\circ} & 0 \end{cases} \quad 0 \quad 0$$

Finally, applying the transition matrices to all the interim points:

$$A2 = \begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 6 & 3 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -2 \\ 2\sqrt{3} \end{bmatrix} = \begin{bmatrix} 2\sqrt{3}+2 \\ 1 \\ 1 \end{bmatrix}$$


Similarly,

$$B_{2} = \begin{bmatrix} 2 \\ 5.46 \\ 5.73 \end{bmatrix}$$
, $C_{2} = \begin{bmatrix} 4 \\ 6.46 \\ 4 \\ 1 \end{bmatrix}$


$$F_2 = \begin{bmatrix} 2 \\ 2 \\ 3.73 \\ 1 \end{bmatrix}$$

$$G_{2} = \begin{bmatrix} 4 \\ 2 \\ 3^{1}73 \end{bmatrix}$$

$$E_{\lambda} = \begin{bmatrix} 4 \\ 3 \\ 2 \\ 1 \end{bmatrix}$$



The Figure will be,



Norry

Midzo[®]

Cipran[®]

Seronex