



1. Download the data set "Revenue.csv" into your working directory.

a) Draw a scatter plot. X-axis is the "Day" and Y-axis is the "Revenue".

```
> df <- read.table("C:\\Users\\Udaya Vijay Anand\\Downloads\\Revenue.csv", header=T)
> head(df)
  Period Day Revenue Sales_quantity Average_cost The_average_annual_payroll_of_the_region
1 01.01.2015   1 16010072      12729      1257.764                    30024676
2 01.02.2015   2 15807587      11636      1358.507                    30024676
3 01.03.2015   3 22047146      15922      1384.697                    30024676
4 01.04.2015   4 18814583      15227      1235.607                    30024676
5 01.05.2015   5 14021480       8620      1626.622                    30024676
6 01.06.2015   6 16783929      13160      1275.375                    30024676
> |
```



b) Draw a line plot.



c) Looking at the plots in a) and b) do you observe any trends or patterns?

Observing the scatter plot and line plot in a) and b), it can be observed that the revenue exhibits a positive trend as time passes, although there are noticeable fluctuations in the data.

## Question 2

The following data set shows the heights and the arm spans of 10 persons.

Height (cm) 178.5 | 176 | 185.6 | 179 | 172 | 167.7 | 174.3 | 136.5 | 164.5 | 165.5

Arm Span (cm) 183.5 | 170 | 194.1 | 181 | 181 | 171.7 | 171 | 135.5 | 164 | 160.5


Create a data frame named mydata using “mydata <- data.frame(height, armspan)”. Here note that you have to create two vectors height, armspan manually.

a) Add a new variable, “apeindex”, as the ratio between armspan and height, to your data. Show your new data set.

```
> height <- c(178.5, 176, 185.6, 179, 172, 167.7, 174.3, 136.5, 164.5, 165.5)
> armspan <- c(183.5, 170, 194.1, 181, 181, 171.7, 171, 135.5, 164, 160.5)
> mydata <- data.frame(height, armspan)
> mydata$apeindex <- mydata$armspan / mydata$height
> mydata$apeindex <- mydata$armspan / mydata$height
> mydata
  height armspan apeindex
1  178.5   183.5  1.0280112
2  176.0   170.0  0.9659091
3  185.6   194.1  1.0457974
4  179.0   181.0  1.0111732
5  172.0   181.0  1.0523256
6  167.7   171.7  1.0238521
7  174.3   171.0  0.9810671
8  136.5   135.5  0.9926740
9  164.5   164.0  0.9969605
10 165.5   160.5  0.9697885
> |
```

b) Attach a boxplot of apeindex. Are there any outliers?

```
> boxplot(mydata$apeindex)
> |
```



c) Find the min, Q1, Q2, Q3, and max of apeindex.

```
> summary(mydata$apeindex)
   Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
0.9659  0.9840  1.0041  1.0068  1.0270  1.0523
> |
```

d) Find IQR,  $Q1 - 1.5 \cdot IQR$ , and  $Q3 + 1.5 \cdot IQR$ .

```
> IQR(mydata$apeindex)
[1] 0.04300259
> Q1 <- quantile(mydata$apeindex, 0.25)
> Q3 <- quantile(mydata$apeindex, 0.75)
> lower <- Q1 - 1.5*IQR(mydata$apeindex)
> upper <- Q3 + 1.5*IQR(mydata$apeindex)
> |
```

```
> lower
   25%
0.919465
> upper
   75%
1.091475
> |
```


e) What is the lower adjacent value (LAV) and upper adjacent value (UAV)? (Note that they're not necessarily equal to  $Q1 - 1.5 \cdot IQR$  or  $Q3 + 1.5 \cdot IQR$ !!!)

```
> LAV <- min(mydata$apeindex[mydata$apeindex > lower])
> UAV <- max(mydata$apeindex[mydata$apeindex < upper])
>
> LAV
[1] 0.9659091
> UAV
[1] 1.052326
> |
```

3. Consider an unfair coin with a chance of  $1/5$  to get head and  $4/5$  to get tail in each toss

a). For 3 independent tosses, what are the outcomes in terms of heads (H) and tails (T) and their relative frequencies? Draw a line plot.

```
> # Simulate 3 coin tosses
> A <- sample(c(0, 1), 3, replace = TRUE)
> A
[1] 0 0 0
>
> # Calculate relative frequency of heads in A
> Rpre <- sum(A == 1) / 3
> Rpre
[1] 0
>
> # Simulate 10000 coin tosses
> X <- sample(c(0, 1), 10000, replace = TRUE)
> Rpreq <- sum(X == 1) / 10000
> Rpreq
[1] 0.4989
>
> # Calculate relative frequency of heads as tosses increase
> Headnums <- cumsum(X == 1)
> Proportions <- Headnums / (1:10000)
>
> # Plot the relative frequencies
> plot(1:10000, Proportions, type = "l", xlab = "# tosses", ylab = "Frequency of heads")
>
> # Add a horizontal line at y=0.5
> lines(1:10000, rep(0.5, 10000), col = "blue")
> |
```



b). For 1000 independent tosses, draw a plot to see how the relative frequency converges

```
> # Define the probabilities of getting heads and tails
> p_H <- 1/5
> p_T <- 4/5
>
> # Simulate 1000 coin tosses with the given probabilities
> outcomes <- sample(c("H", "T"), size = 1000, replace = TRUE, prob = c(p_H, p_T))
>
> # Calculate the relative frequency of heads as the number of tosses increases
> r_freq <- cumsum(outcomes == "H") / (1:1000)
>
> # Plot the relative frequency as a function of the number of tosses
> plot(r_freq, type = "l", xlab = "Toss Number", ylab = "Relative Frequency of H", main = "Relative Frequency Convergence")
>
```

