

# Hypothesis tests with continuous variables

Alex Sanchez, Miriam Mota, Ricardo Gonzalo and  
Santiago Perez-Hoyos

Statistics and Bioinformatics Unit. Vall d'Hebron Institut de Recerca

Versión 2021-11-29


# Outline

- 1 INTRODUCTION
- 2 TYPE OF TESTS
- 3 NORMALITY TESTS
- 4 ONE GROUP COMPARISON
- 5 TWO GROUPS COMPARISON IN INDEPENDENT SAMPLES
- 6 TWO GROUPS COMPARISON IN DEPENDENT SAMPLES
- 7 *K GROUPS COMPARISON IN INDEPENDENT SAMPLES*
- 8 *Multiple comparisons and multiple testing*

# Introduction

- Once the concept of hypothesis testing is established,
- Researchers face the problem of *which test should be applied at every possible situation*.
- Best solution:
  - understand the problem and the questions addressed
  - know available tests for each problem
  - be aware of applicability assumptions of each test and how to check them.
- Easier to say than to do.
  - Sometimes cheatsheets may be helpful, but be warned against a blind use, that is understand and be critic with the steps.

# Which test is appropriate for which problem



## Example situation

- A study was designed to compare two distinct hypertension control programs.
- 60 individuals with HTA were randomly assigned to either one or the other group (30 per group)
- Blood pressure was measured each month during a year

```


display.par(mfrow=c(1,1)) # Guarda los parámetros para el dibujo
par(mfrow=c(2,2)) # Dibuja cuatro gráficos por grafico
with(hta, boxplot(tas1, main="Box-plot") )

with(hta, hist(tas1) )

with(hta, qqnorm(tas1, main="Normal QQplot") );with(hta, qqline(tas1) )

par(oldpar) # Vuelve a los parámetros de dibujo originales

```



# Normality Test

```
with(hta, shapiro.test(tad1) ) # Shapiro Wilk test
```

```
##
```

```
## Shapiro-Wilk normality test
```

```
##
```

```
## data: tad1
```

```
## W = 0.96622, p-value = 0.09512
```

# One sample Test

```
with(hta,t.test(tad1,mu=90) ) # One sample T.test
```

```
##  
## One Sample t-test  
##  
## data: tad1  
## t = -1.2137, df = 59, p-value = 0.2297  
## alternative hypothesis: true mean is not equal to 90  
## 95 percent confidence interval:  
## 85.80626 91.02707  
## sample estimates:  
## mean of x  
## 88.41667
```



# Homogeneity variance Test

```
library(car)
hta%>%
  group_by(sexo) %>%
  summarise(var = sd(tas1))
```

```
## # A tibble: 2 x 2
##   sexo    var
##   <chr> <dbl>
## 1 MUJER  17.6
## 2 VARON  22.1
```

```
with(hta, leveneTest(tad1~factor(sexo), center="median"))
```

```
## Levene's Test for Homogeneity of Variance (center = "median")
##           Df F value Pr(>F)
## group    1  1.3506 0.2499
##           58
```

- p value is over 0.05
- We can assume homogeneity of variances

# T test when variances are equal

```
with(hta,t.test(tas1~factor(sexo),var.equal=TRUE ))
```

```
##
```

```
## Two Sample t-test
```

```
##
```

```
## data: tas1 by factor(sexo)
```

```
## t = -0.2471, df = 58, p-value = 0.8057
```

```
## alternative hypothesis: true difference in means between groups
```

```
## 95 percent confidence interval:
```

```
## -11.603461 9.053519
```

```
## sample estimates:
```

```
## mean in group MUJER mean in group VARON
```

```
## 149.5946 150.8696
```

- Type I Error is over than 0.05
- We cannot reject mean equality

# T test when variances are unequal

```
with(hta,t.test(tas1~factor(sexo),var.equal=FALSE ))  
  
##  
##  Welch Two Sample t-test  
##  
## data:  tas1 by factor(sexo)  
## t = -0.23436, df = 39.098, p-value = 0.8159  
## alternative hypothesis: true difference in means between groups is not equal to 0  
## 95 percent confidence interval:  
##  -12.277927    9.727986  
## sample estimates:  
## mean in group MUJER mean in group VARON  
##           149.5946           150.8696
```

- Same conclusions as before
- Test is also known as Welch test

## U Mann-Whitney or Sum Rank non parametric test

```
with(hta,wilcox.test(tad1~factor(sexo)  
  ,alternative='two.sided',exact=TRUE, correct=FALSE))
```

```
##
```

```
## Wilcoxon rank sum test
```

```
##
```

```
## data: tad1 by factor(sexo)
```

```
## W = 434, p-value = 0.8955
```

```
## alternative hypothesis: true location shift is not equal to 0
```

```
hta%>%
```

```
  group_by(sexo) %>%
```

```
  summarise(median = median(tad1))
```

```
## # A tibble: 2 x 2
```

```
##   sexo median
```

```
##   <chr>   <dbl>
```

```
## 1 MUJER    90
```

```
## 2 VARON    90
```

- Null Hypothesis cannot be rejected

## Paired T-test

```
with(hta,t.test(tas1,tas12,paired=TRUE))
```

```
##
```

```
## Paired t-test
```

```
##
```

```
## data: tas1 and tas12
```

```
## t = 6.0672, df = 51, p-value = 1.609e-07
```

```
## alternative hypothesis: true difference in means is not equal
```

```
## 95 percent confidence interval:
```

```
## 8.518285 16.943253
```

```
## sample estimates:
```

```
## mean of the differences
```

```
## 12.73077
```

```
summary(hta$tas1)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
```

```
##    100.0   140.0   145.0   150.1   160.0   210.0
```

```
summary(hta$tas12)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
```

# Paired Sign-Rank Wilcoxon Test

```
with(hta,wilcox.test(tad1,tad12,  
    exact=TRUE, paired=TRUE))
```

```
##
```

```
## Wilcoxon signed rank test with continuity correction
```

```
##
```

```
## data: tad1 and tad12
```

```
## V = 478.5, p-value = 0.05333
```

```
## alternative hypothesis: true location shift is not equal to 0
```

# Read diabetes data

```
library(readxl)
library(dplyr)
library(magrittr)
diabetes <- read_excel("datasets/diabetes.xls")
sapply(diabetes, class)
```

```
##      numpacie      mort      tempsviu      edat      bmi      edatdiag
## "numeric" "character" "numeric" "numeric" "numeric" "numeric"
##      tabac      sbp      dbp      ecg      chd
## "character" "numeric" "numeric" "character" "character"
```

```
diabetes_factor <- diabetes %>%
  mutate_if(sapply(diabetes, is.character), as.factor) %>%
  select (-numpacie)
```

```
diabetes_factor %>%
  group_by(ecg) %>%
  summarise( n=n(),
             mean = mean(edat),
             sd=sd(edat))
```

```
## # A tibble: 3 x 4
##   ecg      n mean    sd
##   <chr> <int> <dbl> <dbl>
## 1 Anormal    11  64.9  6.76
## 2 Frontera   27  53.8 11.4
## 3 Normal   111  50.5 11.5
```

# ANOVA

```
anova<-aov(edat~ecg,data=diabetes_factor)
summary(anova)
```

```
##              Df Sum Sq Mean Sq F value    Pr(>F)
## ecg           2    2166   1083.0     8.619 0.00029 ***
## Residuals    146   18347    125.7
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1
```



# Multicomparison

```
library(multcomp)
tuk <- glht(anova, linfct = mcp(ecg = "Tukey"))

print(summary(tuk)) # pairwise tests

##
## Simultaneous Tests for General Linear Hypotheses
##
## Multiple Comparisons of Means: Tukey Contrasts
##
##
## Fit: aov(formula = edat ~ ecg, data = diabetes_factor)
##
## Linear Hypotheses:
##
##           Estimate Std. Error t value Pr(>|t|)
## Frontera - Anormal == 0 -11.094      4.010  -2.767 0.016508 *
## Normal - Anormal == 0  -14.405      3.543  -4.065 0.000221 ***
## Normal - Frontera == 0   -3.310      2.405  -1.376 0.345701
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## (Adjusted p values reported -- single-step method)
```

```
print(confint(tuk, level=0.95)) # confidence intervals
```

```
##
## Simultaneous Confidence Intervals
##
## Multiple Comparisons of Means: Tukey Contrasts
##
##
## Fit: aov(formula = edat ~ ecg, data = diabetes_factor)
##
## Quantile = 2.3458
## 95% family-wise confidence level
##
##
## Linear Hypotheses:
##           Estimate lwr      upr
## Frontera - Anormal == 0 -11.0943 -20.5003 -1.6882
## Normal - Anormal == 0  -14.4046 -22.7168 -6.0924
## Normal - Frontera == 0   -3.3103  -8.9530  2.3324
```

# Multicomparison plot

```
plot(confint(tuk))
```

# Kruskal-Wallis Test

```
diabetes_factor%>%  
  group_by(ecg) %>%  
  summarise(median = median(edat))
```

```
## # A tibble: 3 x 2  
##   ecg      median  
##   <fct>    <dbl>  
## 1 Anormal     64  
## 2 Frontera    53  
## 3 Normal     49
```

```
kruskal.test(edat~ecg, data=diabetes_factor)
```

```
##
```

```
## Kruskal-Wallis rank sum test
```

```
##
```

```
## data: edat by ecg
```

```
## Kruskal-Wallis chi-squared = 17.483, df = 2, p-value = 0.0001
```

# Dunn Test for multiple comparison

```
library(dunn.test)
with(diabetes_factor,dunn.test(edat,ecg,method="bonferroni"))
```

```
##    Kruskal-Wallis rank sum test
##
## data: edat and ecg
## Kruskal-Wallis chi-squared = 17.4826, df = 2, p-value = 0
##
##
##                                     Comparison of edat by ecg
##                                     (Bonferroni)
## Col Mean-|
## Row Mean |      Anormal    Frontera
## -----+-----
## Frontera |      2.721182
##           |      0.0098*
##           |
## Normal   |      4.075469    1.467464
##           |      0.0001*      0.2134
##
```