

The following is a description of a project aimed at developing a more direct way of measuring subtle energy. Various forms of indirect measures are available, all of which have various large measures of error or uncertainty. As a starting place, dowsing was chosen as an indirect measure for comparison and in general, to work with a subtle energy source.

Dowsing is indirect in that the operator is part of the instrument with the process subject to high levels of 'experimenter effect'. A direct measure then is one where there is no operator only an observer, one that can be removed in space and time from the experiment. The 'experimenter effect' can then be greatly reduced, to something like 'observer effect'. This will be discussed later because it is important in calibrating the instrument.

Another aspect of 'indirectness' is the interfaceability of the active measuring element to subtle energy. Electro-magnetic interactions seem removed from subtle energy. For example, in Kirlian photography it seems as though the ionization process is not as important as film emulsion reaction. Hence it is difficult to determine exactly where and when the subtle energy interfacing is occurring.

To consider what does constitute a more direct interface, the source of subtle energy to be measured must be considered. A principle of synthetic geometry based accumulators is primarily that a precise form in space and time is in a state of harmonic relationship with the space time manifold and in some manner subtle energy is 'accumulated'. No kinetic energy is transferred to the accumulator yet it undergoes a 'change of state' which challenges conservation of energy, especially if the accumulator puts out kinetic energy and performs work. This suggests another dimensional avenue of exchange for kinetic energy and would seem to satisfy some theories of multi-dimensional universes and is important in considering the manifold.

The space time manifold can be thought of as the sum total of the potential energy of the universe. In that sense it represents



the archetype form of the sum total vectors of the potential, in other words the 4th dimension is generated by the 3 dimensions. Holisms that evolve in the manifold have a potential energy relationship with the 4th dimension or manifold.

This 'potential relativity' between micro-cosmic holism and the universe archetype can be thought of as a hologram like seed pattern in the 4th dimension along which holisms evolve. Subtle energy can then be seen as changes in state of potential energy relationships or the invisible exchange of energy through or between an alternative dimensional space. In dowsing, the operator is the holism that is sensitive to the potential pattern.

The potential relationship of a single atom is a 'form' of probabilities or tendencies in and around the atom in 4 dimensions, giving rise to its structure.

A simple way of approaching this is to look at the atom in a 'Newtonian' sense, in that the particles are orbiting and can be found in many possible configurations. Some of the possible configurations are geometrically resonant with the manifold. When the atom reaches one of these configurations it is open to explicate effects. More specifically, when the particles transit the resonant configuration shear waves are generated opening other dimensional paths to the atom. This process can be seen as the trigger in spontaneous events in the atom like radioactive decay or spontaneous transitions. Alternately, an interstitial path is formed when an electron emits light or a nucleus emits a decay product.

Since it is not possible to predict an event it is said to be random and a half life of a particular population of atoms is a constant. However, an atom can be externally triggered if the potential probabilities are changed i.e., if it is exposed to external subtle energy or if it is involved geometrically with a larger holism, it will change its half life by some degree proportionally to the change in subtle energy. This principle also opens the possibility of some periodicity in an isotope's output, an amount that would loosely reflect the geometry of the manifold.



Measuring the change or variance of an isotope's output is somewhat more direct in that it responds to the manifold through a mediator to be measured.

An isotope standing alone responds to the manifold directly and yields what we observe as random with a hint at some periodicity. Receiving meaningful data about a modifier can be accomplished with 2 isotopes, one as a control (interfaced to background) the other involved with a modifier, in essence yielding data about the differences in the relationship of an isotope to the manifold and an isotope through a modifier to the manifold. The variance detector is just this, a differential geiger counter. This principle was first used by Helmut Schmidt as a test for PK. He and others have found a variation proportional to PK.

In order to develop a better understanding I chose to work with the Genesa crystals as the source of subtle energy. Cognitive information about the manifold is inherent in the Genesa system and it is possible to explore a modifier's (Genesa crystal) relationship to the manifold in a much more detailed way than a complex PK source because the crystal can be accessed in several specific ways (i.e., along different rays), yielding more information about the manifold.

During the development of the instrument it became apparent that before any serious Genesa crystal studies could be done an extensive calibration and background study had to be performed. Also this needs to be done separately for alpha, beta, and gamma based detectors.

My work to date with beta and gamma have shown some peculiar background data. For one, the 'observer effect' is readily demonstrable. If I am anywhere near the detectors during a prolonged zero differential reading, the average zero reading is higher than if I am not present. Also periodicities come and go and seem affected by my presence.

Other considerations that seem to affect zero reading are: sun and moon positions, weather (other than just humidity a factor in alpha attenuation) and the relative positions of the two

detectors (including distance from each other).

Preliminary results of the Genesa Crystal testing are favorable. The first experiments were simply to try and verify a dowsing reading of a ray of the crystal. The variance detector yields as much as a 2% differential with zero being established at no greater than .8%. This is in keeping with the results of Schmidt's PK testing.

Before proceeding with Genesa crystal testing a thorough accounting of the background and more refining of the detectors is required. Since the hardware that supports the isotope and GM tube is itself a modifier, an accounting of the interfaceability of its hardware needs to be considered. Selection of isotopes and baffles is a subject in the refinement of detectors. The purity of the type of radiation to be observed is important in the elimination of 'noise' and in accurate zero calibration. Calibration to offset the average isotope output differences is accomplished by adjusting the source distance to the GM tube. This can be a source of noise if different types of particles are mixed, in that air attenuation is different for each type. Other sources of noise to be reckoned with include cosmic ray intrusion, GM tube characteristics and possible subtle energy effects in the tube.

It is hoped that the refinement of this technique will lead to some interesting experimental possibilities. It is significant alone that a radio isotope can be affected without some specific kinetic interaction like particle collisions as in a fission reactor. Further there is hope of developing an accurate tool for the study and development of psychotronic technology.