



BLM3620 Digital Signal Processing*

Erkan Uslu

euslu@yildiz.edu.tr

Yıldız Technical University – Computer Engineering

*Based on lecture notes from Ali Can Karaca & Ahmet Elbir

Lecture #13 – z - Transform

- Introduce the z-Transform
- Examples
- Transform Example
- MATLAB Applications


- The FFT is simply an algorithm for efficiently calculating the DFT
- Computational efficiency of an N-Point FFT:
 - DFT: N^2 Complex Multiplications
 - FFT: $(N/2) \log_2(N)$ Complex Multiplications

| N | DFT Multiplications | FFT Multiplications | FFT Efficiency |
|-------|---------------------|---------------------|----------------|
| 256 | 65,536 | 1,024 | 64 : 1 |
| 512 | 262,144 | 2,304 | 114 : 1 |
| 1,024 | 1,048,576 | 5,120 | 205 : 1 |
| 2,048 | 4,194,304 | 11,264 | 372 : 1 |
| 4,096 | 16,777,216 | 24,576 | 683 : 1 |

Discrete Fourier Transform (DFT)

$$X[k] = \sum_{n=0}^{N-1} x[n] e^{-j(2\pi/N)kn}$$

Recap: Frequency Response $H(e^{j\hat{\omega}})$



$$H(e^{j\hat{\omega}}) = \sum_{n=-\infty}^{\infty} h[n]e^{-j\hat{\omega}n}$$


$$\text{Periodic : } H(e^{j(\hat{\omega}+2\pi)}) = H(e^{j\hat{\omega}})$$

$$y[n] = A \cdot |H(e^{j\hat{\omega}_0})| \cos(\hat{\omega}_0 n + \varphi + \angle H(e^{j\hat{\omega}_0}))$$

Recap: Transfer Function/System Function $H(z)$

Given impulse response of an LTI system $h[n]$, the transfer function can be found as:

$$H[z] = \sum_{k=-\infty}^{\infty} h[k] z^{-k}$$



$$= e^{j\hat{w}}$$

Z transform can be computed if FT exists or not.


Easy to compute the system output.

Mostly used with block diagrams to illustrate the system.

Recap: Transfer Function/System Function $H(z)$

Given impulse response of an LTI system $h[n]$, the transfer function can be found as:

$$H[z] = \sum_{k=-\infty}^{\infty} h[k] z^{-k}$$



Z-transform is a Generalized version of Discrete Time Fourier
 $= e^{j\hat{w}}$)

Z transform can be computed if FT exists or not.

Easy to compute the system output.


Mostly used with block diagrams to illustrate the system.

Recap: Transfer Function/System Function $H(z)$

e j w e e e j w j j w w w w e j w)

Given impulse response of an LTI system $h[n]$, the transfer function can be found as:

$$H[z] = \sum_{k=-\infty}^{\infty} h[k] z^{-k}$$



Z-transform is a Generalized version of Discrete Time Fourier Transform ($z = e^{j\hat{\omega}}$)

Z transform can be computed if FT exists or not.

Easy to compute the system output.


Mostly used with block diagrams to illustrate the system.

Recap: Transfer Function/System Function $H(z)$

e j w e e e j w j j w w w w e j w)

Given impulse response of an LTI system $h[n]$, the transfer function can be found as:

$$H[z] = \sum_{k=-\infty}^{\infty} h[k] z^{-k}$$



Z-transform is a Generalized version of Discrete Time Fourier Transform ($z = e^{j\hat{\omega}}$)

Z transform can be computed if FT exists or not.

Easy to compute the system output.

Mostly used with block diagrams to illustrate the system.

Given impulse response of an LTI system $h[n]$, the transfer function can be found as:

The diagram consists of two parts, each showing a block labeled "System".

In the top part, an input signal z^n enters the system from the left, and the output signal is $H(z)z^n$ exiting to the right.

In the bottom part, an input signal $e^{j\hat{w}n}$ enters the system from the left, and the output signal is $H(e^{j\hat{w}})e^{j\hat{w}n}$ exiting to the right.

Z transform can be computed if FT exists or not.

Easy to compute the system output.

Mostly used with block diagrams to illustrate the system

Given impulse response of an LTI system $h[n]$, the transfer function can be found as:

The diagram consists of two parts, one above the other, each showing a block labeled "System".

In the top part, an input signal z^n enters the system from the left, and the output signal is $H(z)z^n$ exiting to the right.

In the bottom part, the input signal is $e^{j\hat{w}n}$ entering the system from the left, and the output signal is $H(e^{j\hat{w}})e^{j\hat{w}n}$ exiting to the right.

Easy to compute the system output.


Mostly used with block diagrams to illustrate the system.

Recap: Transfer Function/System Function $H(z)$

e j w e e e j w j j w w w w e j w)

Given impulse response of an LTI system $h[n]$, the transfer function can be found as:

$$H[z] = \sum_{k=-\infty}^{\infty} h[k] z^{-k}$$



Z-transform is a Generalized version of Discrete Time Fourier Transform ($z = e^{j\hat{\omega}}$)

Z transform can be computed if FT exists or not.


Z transform can be computed if FT exists or not.

Easy to compute the system output.


Easy to compute the system output.

Mostly used with block diagrams to illustrate the system. Mostly used with block diagrams to illustrate


Three Domains



Three Domains



Three Domains



Z-Transform DEFINITION

- POLYNOMIAL Representation of LTI SYSTEM:

- EXAMPLE:

$$H(z) = \sum_n h[n]z^{-n}$$

*APPLIES to
any SIGNAL*

Z-Transform DEFINITION

- POLYNOMIAL Representation of LTI SYSTEM:

- EXAMPLE:

$$H(z) = \sum_n h[n]z^{-n}$$

$$\{h[n]\} = \{2, 0, -3, 0, 2\}$$

*APPLIES to
any SIGNAL*

Z-Transform DEFINITION



- POLYNOMIAL Representation of LTI SYSTEM:

- EXAMPLE:

$$H(z) = \sum_n h[n]z^{-n}$$

*APPLIES to
any SIGNAL*

$$\{h[n]\} = \{2, 0, -3, 0, 2\}$$

$$H(z) = 2z^{-0} + 0z^{-1} - 3z^{-2} + 0z^{-3} + 2z^{-4}$$

Z-Transform DEFINITION



- POLYNOMIAL Representation of LTI SYSTEM:

- EXAMPLE:

$$H(z) = \sum_n h[n]z^{-n}$$

*APPLIES to
any SIGNAL*

$$\{h[n]\} = \{2, 0, -3, 0, 2\}$$

$$\begin{aligned} H(z) &= 2z^{-0} + 0z^{-1} - 3z^{-2} + 0z^{-3} + 2z^{-4} \\ &= 2 - 3z^{-2} + 2z^{-4} \end{aligned}$$

Z-Transform DEFINITION



- POLYNOMIAL Representation of LTI SYSTEM:

- EXAMPLE:

$$H(z) = \sum_n h[n]z^{-n}$$

*APPLIES to
any SIGNAL*

$$\{h[n]\} = \{2, 0, -3, 0, 2\}$$

$$H(z) = 2z^{-0} + 0z^{-1} - 3z^{-2} + 0z^{-3} + 2z^{-4}$$

$$= 2 - 3z^{-2} + 2z^{-4}$$

$$= 2 - 3(z^{-1})^2 + 2(z^{-1})^4$$

POLYNOMIAL in z^{-1}

$y[n] - y[n-1] = x[n] + 2x[n-1]$ sisteminin transfer fonksiyonunu bulunuz.

Fark denkleminde her iki tarafın z-dönüşümü alındığında

$$Y(z) - z^{-1}Y(z) = X(z) + 2z^{-1}X(z)$$

elde edilmektedir. Sistemin transfer fonksiyonu

$$H(z) = \frac{Y(z)}{X(z)} = \frac{1 + 2z^{-1}}{1 - z^{-1}}$$

olarak bulunmaktadır.

Blok Diyagram Gösterimi

$y[n] - 4.5y[n-1] + 2y[n-2] = x[n]$ şeklinde tanımlanmış sistemin blok diyagramını çizelim.

Blok Diyagram Gösterimi


$y[n] - 4.5y[n-1] + 2y[n-2] = x[n]$ şeklinde tanımlanmış sistemin blok diyagramını çizelim.

$$y[n] = x[n] + 4.5y[n-1] - 2y[n-2]$$

Blok Diyagram Gösterimi

$y[n] - 4.5y[n-1] + 2y[n-2] = x[n]$ şeklinde tanımlanmış sistemin blok diyagramını çizelim.

$$y[n] = x[n] + 4.5y[n-1] - 2y[n-2]$$




Credit by Sarp Ertürk

Blok Diyagram Gösterimi

$y[n] - 4.5y[n-1] + 2y[n-2] = x[n]$ şeklinde tanımlanmış sistemin blok diyagramını çizelim.

$$y[n] = x[n] + 4.5y[n-1] - 2y[n-2]$$

$$H(z) = \frac{1}{1 - 4.5z^{-1} + 2z^{-2}}$$



Credit by Sarp Ertürk

Z-Transform EXAMPLE

- ANY SIGNAL has a z-Transform:

$$X(z) = \sum_n x[n]z^{-n}$$

$$H(z) = \sum_n h[n]z^{-n}$$

Z-Transform EXAMPLE

- ANY SIGNAL has a z-Transform:

$$X(z) = \sum_n x[n]z^{-n}$$

$$H(z) = \sum_n h[n]z^{-n}$$

Example 7.1

| n | $n < -1$ | -1 | 0 | 1 | 2 | 3 | 4 | 5 | $n > 5$ |
|--------|----------|------|-----|-----|-----|-----|-----|-----|---------|
| $x[n]$ | 0 | 0 | 2 | 4 | 6 | 4 | 2 | 0 | 0 |

Z-Transform EXAMPLE

- ANY SIGNAL has a z-Transform:

$$X(z) = \sum_n x[n]z^{-n}$$

$$H(z) = \sum_n h[n]z^{-n}$$

Example 7.1

| n | $n < -1$ | -1 | 0 | 1 | 2 | 3 | 4 | 5 | $n > 5$ |
|--------|----------|------|-----|-----|-----|-----|-----|-----|---------|
| $x[n]$ | 0 | 0 | 2 | 4 | 6 | 4 | 2 | 0 | 0 |

$$X(z) = ?$$

Z-Transform EXAMPLE

- ANY SIGNAL has a z-Transform:

$$X(z) = \sum_n x[n]z^{-n}$$

$$H(z) = \sum_n h[n]z^{-n}$$

Example 7.1

| n | $n < -1$ | -1 | 0 | 1 | 2 | 3 | 4 | 5 | $n > 5$ |
|--------|----------|------|-----|-----|-----|-----|-----|-----|---------|
| $x[n]$ | 0 | 0 | 2 | 4 | 6 | 4 | 2 | 0 | 0 |

$$X(z) = ?$$

$$X(z) = 2 + 4z^{-1} + 6z^{-2} + 4z^{-3} + 2z^{-4}$$

Z-Transform EXAMPLE

- ANY SIGNAL has a z-Transform:

$$X(z) = \sum_n x[n]z^{-n}$$

$$H(z) = \sum_n h[n]z^{-n}$$

Example 7.1

| n | $n < -1$ | -1 | 0 | 1 | 2 | 3 | 4 | 5 | $n > 5$ |
|--------|----------|------|-----|-----|-----|-----|-----|-----|---------|
| $x[n]$ | 0 | 0 | 2 | 4 | 6 | 4 | 2 | 0 | 0 |

$$X(z) = ?$$

$$X(z) = 2 + 4z^{-1} + 6z^{-2} + 4z^{-3} + 2z^{-4}$$

Example 9.2



$$X(z) = 1 - 2z^{-1} + 3z^{-3} - z^{-5}$$

**EXPONENT GIVES
TIME LOCATION**

$$x[n] = ?$$

Example 9.2



$$X(z) = 1 - 2z^{-1} + 3z^{-3} - z^{-5}$$

EXPONENT GIVES
TIME LOCATION

$$x[n] = ?$$

$$x[n] = \begin{cases} 0 & n < 0 \\ 1 & n = 0 \\ -2 & n = 1 \\ 0 & n = 2 \\ 3 & n = 3 \\ 0 & n = 4 \\ -1 & n = 5 \\ 0 & n > 5 \end{cases}$$

$$x[n] = \delta[n] - 2\delta[n - 1] + 3\delta[n - 3] - \delta[n - 5]$$

Example 9.2



$$X(z) = 1 - 2z^{-1} + 3z^{-3} - z^{-5}$$

EXPONENT GIVES
TIME LOCATION

$$x[n] = ?$$

$$x[n] = \begin{cases} 0 & n < 0 \\ 1 & n = 0 \\ -2 & n = 1 \\ 0 & n = 2 \\ 3 & n = 3 \\ 0 & n = 4 \\ -1 & n = 5 \\ 0 & n > 5 \end{cases}$$

$$x[n] = \delta[n] - 2\delta[n - 1] + 3\delta[n - 3] - \delta[n - 5]$$

Z-Transform Property: Delay Property

A delay of one sample multiplies the z -transform by z^{-1} .

$$x[n - 1] \iff z^{-1} X(z)$$

Time delay of n_0 samples multiplies the z -transform by z^{-n_0}

$$x[n - n_0] \iff z^{-n_0} X(z)$$

Example:



Find z-transform of shifted impulse $x[n] = 3\delta[n - 5]$.

$$X(z) = \sum_{n=-\infty}^{\infty} x[n] z^{-n}$$

$$X(z) = \sum_{n=-\infty}^{\infty} 3\delta[n - 5] z^{-n} = 3\delta[0]z^{-5} = 3z^{-5}$$

Example:



Find z-transform of shifted impulse $x[n] = 3\delta[n - 5]$.

$$X(z) = \sum_{n=-\infty}^{\infty} x[n] z^{-n}$$

$$X(z) = \sum_{n=-\infty}^{\infty} 3\delta[n - 5] z^{-n} = 3\delta[0]z^{-5} = 3z^{-5}$$

Fourier transform?

Example:



Find z-transform of shifted impulse $x[n] = 3\delta[n - 5]$.

$$X(z) = \sum_{n=-\infty}^{\infty} x[n] z^{-n}$$

$$X(z) = \sum_{n=-\infty}^{\infty} 3\delta[n - 5] z^{-n} = 3\delta[0]z^{-5} = 3z^{-5}$$

Fourier transform?

$$X(e^{j\Omega}) = X[z] \Big|_{z=e^{j\omega}} = 3e^{-j5\omega}$$

Z-Transform of FIR Filter

- CALLED the **SYSTEM FUNCTION**
 - because $h[n]$ is same as $\{b_k\}$

SYSTEM
FUNCTION

$$H(z) = \sum_{k=0}^M b_k z^{-k} = \sum_{k=0}^M h[k] z^{-k}$$

$$y[n] = \sum_{k=0}^M b_k x[n-k] = \sum_{k=0}^M h[k] x[n-k]$$

FIR DIFFERENCE EQUATION

CONVOLUTION

Z-Transform of FIR Filter

- Get $H(z)$ DIRECTLY from the $\{b_k\}$
- Example 7.3 in the book:

$$y[n] = 6x[n] - 5x[n-1] + x[n-2]$$

$$\{b_k\} = \{6, -5, 1\}$$

$$H(z) = \sum b_k z^{-k} = 6 - 5z^{-1} + z^{-2}$$

- Input is $x[n]$, find $y[n]$ (for FIR, $h[n]$)
- How to combine $X(z)$ and $H(z)$?

Example 7.5

$$x[n] = \delta[n - 1] - \delta[n - 2] + \delta[n - 3] - \delta[n - 4]$$

$$\text{and } h[n] = \delta[n] + 2\delta[n - 1] + 3\delta[n - 2] + 4\delta[n - 3]$$

$$X(z) = 0 + 1z^{-1} - 1z^{-2} + 1z^{-3} - 1z^{-4}$$

$$\text{and } H(z) = 1 + 2z^{-1} + 3z^{-2} + 4z^{-3}$$

FIR Filter = CONVOLUTION

$$y[n] = \sum_{k=0}^M b_k x[n-k] = \sum_{k=0}^M h[k] x[n-k]$$

CONVOLUTION

FIR Filter = CONVOLUTION

| | | | | | | | | |
|--------------|---|----|----|----|----|----|----|----|
| $x[n], X(z)$ | 0 | +1 | -1 | +1 | -1 | | | |
| $h[n], H(z)$ | 1 | 2 | 3 | 4 | | | | |
| <hr/> | | | | | | | | |
| | 0 | +1 | -1 | +1 | -1 | | | |
| | | 0 | +2 | -2 | +2 | -2 | | |
| | | | 0 | +3 | -3 | +3 | -3 | |
| | | | | 0 | +4 | -4 | +4 | -4 |
| <hr/> | | | | | | | | |
| $y[n], Y(z)$ | 0 | +1 | +1 | +2 | +2 | -3 | +1 | -4 |

$$y[n] = \sum_{k=0}^M b_k x[n-k] = \sum_{k=0}^M h[k] x[n-k]$$

CONVOLUTION

CONVOLUTION PROPERTY



- PROOF:

$$y[n] = x[n] * h[n] = \sum_{k=0}^M h[k]x[n - k]$$

$$Y(z) = \sum_{k=0}^M h[k] (z^{-k} X(z))$$

MULTIPLY
z-TRANSFORMS

$$= \left(\sum_{k=0}^M h[k]z^{-k} \right) X(z) = H(z)X(z).$$



- **MULTIPLY** the z-TRANSFORMS:

Example 7.5

$$x[n] = \delta[n - 1] - \delta[n - 2] + \delta[n - 3] - \delta[n - 4]$$

$$\text{and } h[n] = \delta[n] + 2\delta[n - 1] + 3\delta[n - 2] + 4\delta[n - 3]$$

$$X(z) = 0 + 1z^{-1} - 1z^{-2} + 1z^{-3} - 1z^{-4}$$

$$\text{and } H(z) = 1 + 2z^{-1} + 3z^{-2} + 4z^{-3}$$

$$y[n] = h[n] * x[n] \quad \leftrightarrow \quad Y(z) = H(z)X(z)$$

CONVOLUTION EXAMPLE



- Finite-Length input $x[n]$
- FIR Filter ($L=4$)


**MULTIPLY
Z-TRANSFORMS**

$$\begin{aligned} Y(z) &= H(z)X(z) \\ &= (1 + 2z^{-1} + 3z^{-2} + 4z^{-3})(z^{-1} - z^{-2} + z^{-3} - z^{-4}) \\ &= z^{-1} + (-1 + 2)z^{-2} + (1 - 2 + 3)z^{-3} + (-1 + 2 - 3 + 4)z^{-4} \\ &\quad + (-2 + 3 - 4)z^{-5} + (-3 + 4)z^{-6} + (-4)z^{-7} \\ &= z^{-1} + z^{-2} + 2z^{-3} + 2z^{-4} - 3z^{-5} + z^{-6} - 4z^{-7} \end{aligned}$$

$y[n] = ?$


Z-Transform Property: CASCADE EQUIVALENT

- Multiply the System Functions





Z-Transform Property: CASCADE EQUIVALENT



- Multiply the System Functions




CASCADE EXAMPLE




CASCADE EXAMPLE




CASCADE EXAMPLE



CASCADE EXAMPLE



CASCADE EXAMPLE



Region of Convergence (ROC)



- Give a sequence, **the set of values of z** for which the z -transform **converges**, i.e., $|X(z)| < \infty$, is called the **region of convergence**.

$$ROC = \{z = re^{j\omega} \mid R_{x-} < r < R_{x+}\}$$


Region of Convergence (ROC)

- Give a sequence, **the set of values of z** for which the z -transform **converges**, i.e., **$|X(z)| < \infty$** , is called the **region of convergence**.

$$|X(z)| = \left| \sum_{n=-\infty}^{\infty} x(n) z^{-n} \right| = \sum_{n=-\infty}^{\infty} |x(n)| |z|^{-n} < \infty$$

ROC is centered on origin and consists of a set of rings.


$$ROC = \{z = re^{j\omega} \mid R_{x-} < r < R_{x+}\}$$



Stable Systems




- A stable system requires that its **Fourier transform** is uniformly convergent.



Stable Systems



- A stable system requires that its **Fourier transform** is uniformly convergent.




- Fact: Fourier transform is to evaluate z -transform on a unit circle.

Stable Systems



- A stable system requires that its **Fourier transform** is uniformly convergent.



- Fact: Fourier transform is to evaluate z -transform on a unit circle.
- A stable system requires the ROC of z -transform to include the unit circle.

Example: A right sided Sequence



$$x(n) = a^n u(n)$$

$$X(z) = \sum_{n=-\infty}^{\infty} a^n u(n) z^{-n}$$

Example: A right sided Sequence



$$x(n) = a^n u(n)$$

$$\begin{aligned} X(z) &= \sum_{n=-\infty}^{\infty} a^n u(n) z^{-n} \\ &= \sum_{n=0}^{\infty} a^n z^{-n} \end{aligned}$$

Example: A right sided Sequence



$$x(n) = a^n u(n)$$

$$\begin{aligned} X(z) &= \sum_{n=-\infty}^{\infty} a^n u(n) z^{-n} \\ &= \sum_{n=0}^{\infty} a^n z^{-n} \\ &= \sum_{n=0}^{\infty} (az^{-1})^n \end{aligned}$$

For convergence of $X(z)$, we require that

Example: A right sided Sequence



$$x(n) = a^n u(n)$$

$$\begin{aligned} X(z) &= \sum_{n=-\infty}^{\infty} a^n u(n) z^{-n} \\ &= \sum_{n=0}^{\infty} a^n z^{-n} \\ &= \sum_{n=0}^{\infty} (az^{-1})^n \end{aligned}$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |az^{-1}| < \infty$$

Example: A right sided Sequence



$$x(n) = a^n u(n)$$

$$\begin{aligned} X(z) &= \sum_{n=-\infty}^{\infty} a^n u(n) z^{-n} \\ &= \sum_{n=0}^{\infty} a^n z^{-n} \\ &= \sum_{n=0}^{\infty} (az^{-1})^n \end{aligned}$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |az^{-1}| < \infty \quad \longrightarrow \quad |az^{-1}| < 1$$

Example: A right sided Sequence



$$x(n) = a^n u(n)$$

$$\begin{aligned} X(z) &= \sum_{n=-\infty}^{\infty} a^n u(n) z^{-n} \\ &= \sum_{n=0}^{\infty} a^n z^{-n} \\ &= \sum_{n=0}^{\infty} (az^{-1})^n \end{aligned}$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |az^{-1}| < \infty \quad \longrightarrow \quad |az^{-1}| < 1$$

$$\longrightarrow \quad |z| > |a|$$

Example: A right sided Sequence

$$x(n) = a^n u(n)$$

$$\begin{aligned} X(z) &= \sum_{n=-\infty}^{\infty} a^n u(n) z^{-n} \\ &= \sum_{n=0}^{\infty} a^n z^{-n} \\ &= \sum_{n=0}^{\infty} (az^{-1})^n \end{aligned}$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |az^{-1}| < \infty \quad \longrightarrow \quad |az^{-1}| < 1$$

$$\longrightarrow \quad |z| > |a|$$

$$X(z) = \sum_{n=0}^{\infty} (az^{-1})^n = \frac{1}{1 - az^{-1}} = \frac{z}{z - a}$$


$$|z| > |a|$$

Example: A right sided Sequence ROC for $x(n)=a^n u(n)$

$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$


Example: A right sided Sequence ROC for $x(n)=a^n u(n)$

$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$




Example: A right sided Sequence ROC for $x(n)=a^n u(n)$

$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$





Example: A right sided Sequence ROC for $x(n)=a^n u(n)$

$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$





Example: A right sided Sequence ROC for $x(n)=a^n u(n)$

$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$





Example: A right sided Sequence ROC for $x(n)=a^n u(n)$

$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$



Example: A right sided Sequence ROC for $x(n)=a^n u(n)$



$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$



Example: A right sided Sequence ROC for $x(n)=a^n u(n)$

$$X(z) = \frac{z}{z-a}, \quad |z| > |a|$$


Which one is stable?



Another Example



$$x(n) = -a^n u(-n-1)$$



Example: A left sided Sequence



$$x(n) = -a^n u(-n-1)$$

$$X(z) = -\sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n}$$

Example: A left sided Sequence



$$x(n) = -a^n u(-n-1)$$

$$\begin{aligned} X(z) &= - \sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n} \\ &= - \sum_{n=-\infty}^{-1} a^n z^{-n} \end{aligned}$$

Example: A left sided Sequence



$$x(n) = -a^n u(-n-1)$$

$$\begin{aligned} X(z) &= - \sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n} \\ &= - \sum_{n=-\infty}^{-1} a^n z^{-n} \\ &= - \sum_{n=1}^{\infty} a^{-n} z^n \end{aligned}$$

Example: A left sided Sequence



$$x(n) = -a^n u(-n-1)$$

$$\begin{aligned} X(z) &= -\sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n} \\ &= -\sum_{n=-\infty}^{-1} a^n z^{-n} \\ &= -\sum_{n=1}^{\infty} a^{-n} z^n \\ &= 1 - \sum_{n=0}^{\infty} a^{-n} z^n \end{aligned}$$

For convergence of $X(z)$, we require that

Example: A left sided Sequence



$$x(n) = -a^n u(-n-1)$$

$$\begin{aligned} X(z) &= -\sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n} \\ &= -\sum_{n=-\infty}^{-1} a^n z^{-n} \\ &= -\sum_{n=1}^{\infty} a^{-n} z^n \\ &= 1 - \sum_{n=0}^{\infty} a^{-n} z^n \end{aligned}$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |a^{-1} z| < \infty$$

Example: A left sided Sequence

$$x(n) = -a^n u(-n-1)$$

$$\begin{aligned} X(z) &= -\sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n} \\ &= -\sum_{n=-\infty}^{-1} a^n z^{-n} \\ &= -\sum_{n=1}^{\infty} a^{-n} z^n \\ &= 1 - \sum_{n=0}^{\infty} a^{-n} z^n \end{aligned}$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |a^{-1} z| < \infty \quad \longrightarrow \quad |a^{-1} z| < 1$$

Example: A left sided Sequence

$$x(n) = -a^n u(-n-1)$$

$$\begin{aligned} X(z) &= -\sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n} \\ &= -\sum_{n=-\infty}^{-1} a^n z^{-n} \\ &= -\sum_{n=1}^{\infty} a^{-n} z^n \\ &= 1 - \sum_{n=0}^{\infty} a^{-n} z^n \end{aligned}$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |a^{-1} z| < \infty \quad \longrightarrow \quad |a^{-1} z| < 1$$

$$\longrightarrow \quad |z| < |a|$$

Example: A left sided Sequence

$$x(n) = -a^n u(-n-1)$$

$$X(z) = -\sum_{n=-\infty}^{\infty} a^n u(-n-1) z^{-n}$$

$$= -\sum_{n=-\infty}^{-1} a^n z^{-n}$$

$$= -\sum_{n=1}^{\infty} a^{-n} z^n$$

$$= 1 - \sum_{n=0}^{\infty} a^{-n} z^n$$

For convergence of $X(z)$, we require that

$$\sum_{n=0}^{\infty} |a^{-1} z| < \infty \quad \longrightarrow \quad |a^{-1} z| < 1$$

$$\longrightarrow \quad |z| < |a|$$

$$X(z) = 1 - \sum_{n=0}^{\infty} (a^{-1} z)^n = 1 - \frac{1}{1 - a^{-1} z} = \frac{z}{z - a}$$

$$|z| < |a|$$


Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$



$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$


Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$

$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$




Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$

$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$





Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$

$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$





Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$

$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$





Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$

$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$



Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$



$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$



Example: A left sided Sequence ROC for $x(n)=-a^n u(-n-1)$

$$X(z) = \frac{z}{z-a}, \quad |z| < |a|$$

Which one is stable?



Example: A right sided Sequence



$$x(n) = a^n u(n)$$

Example: A right sided Sequence




$$x(n) = a^n u(n) \quad \longrightarrow \quad X(z) = \frac{z}{z - a}, \quad |z| > |a|$$

Example: A right sided Sequence




$$x(n) = a^n u(n) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| > |a|$$



Example: A right sided Sequence




$$x(n) = a^n u(n) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| > |a|$$



Example: A right sided Sequence




$$x(n) = a^n u(n) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| > |a|$$



Example: A right sided Sequence




$$x(n) = a^n u(n) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| > |a|$$



Example: A right sided Sequence



$$x(n) = a^n u(n) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| > |a|$$



ROC is **bounded by the pole** and is the **exterior of a circle**.

Example: A left sided Sequence



$$x(n) = -a^n u(-n-1)$$

Example: A left sided Sequence




$$x(n) = -a^n u(-n-1) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| < |a|$$

Example: A left sided Sequence




$$x(n) = -a^n u(-n-1) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| < |a|$$



Example: A left sided Sequence




$$x(n) = -a^n u(-n-1) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| < |a|$$



Example: A left sided Sequence




$$x(n) = -a^n u(-n-1) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| < |a|$$



Example: A left sided Sequence




$$x(n) = -a^n u(-n-1) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| < |a|$$



Example: A left sided Sequence



$$x(n) = -a^n u(-n-1) \quad \longrightarrow \quad X(z) = \frac{z}{z-a}, \quad |z| < |a|$$



ROC is **bounded by the pole** and is the **interior of a circle**.

Example: Sum of Two Right Sided Sequences




Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$

Example: Sum of Two Right Sided Sequences


$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$



$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}}$$

Example: Sum of Two Right Sided Sequences


$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$


$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$

Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$


➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$



Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$


➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$



Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$


➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$



Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$


➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$



Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$


➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$



Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$


➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$



Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$

➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$




ROC is **bounded by poles**
and is the **exterior of a circle**.

Example: Sum of Two Right Sided Sequences

$$x(n) = \left(\frac{1}{2}\right)^n u(n) + \left(-\frac{1}{3}\right)^n u(n)$$

➔
$$X(z) = \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}} = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$



ROC is **bounded by poles**
and is the **exterior of a circle**.

ROC does not include any pole.

Example: A Two Sided Sequence




$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$

Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$



$$X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}}$$

Example: A Two Sided Sequence




$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$

$$\rightarrow X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$$


Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$




$$X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$$




Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$


 $X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$




Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$




$$X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$$




Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$




$$X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$$




Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$


 $X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$




Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$


 $X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$



Example: A Two Sided Sequence

$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$

$$\rightarrow X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$$



ROC is **bounded by poles**
and is a **ring**.


Example: A Two Sided Sequence



$$x(n) = \left(-\frac{1}{3}\right)^n u(n) - \left(\frac{1}{2}\right)^n u(-n-1)$$

➔

$$X(z) = \frac{z}{z + \frac{1}{3}} + \frac{z}{z - \frac{1}{2}} = \frac{2z(z - \frac{1}{12})}{(z + \frac{1}{3})(z - \frac{1}{2})}$$



ROC is **bounded by poles**
and is a **ring**.

ROC does not include any pole.

Represent z-transform as a Rational Function



$$X(z) = \frac{P(z)}{Q(z)}$$

where $P(z)$ and $Q(z)$ are polynomials in z .

Represent z-transform as a Rational Function

$$X(z) = \frac{P(z)}{Q(z)} \quad \text{where } P(z) \text{ and } Q(z) \text{ are polynomials in } z.$$

Zeros: The values of z 's such that $X(z) = 0$

Represent z-transform as a Rational Function

$$X(z) = \frac{P(z)}{Q(z)} \quad \text{where } P(z) \text{ and } Q(z) \text{ are polynomials in } z.$$

Zeros: The values of z 's such that $X(z) = 0$

Poles: The values of z 's such that $X(z) = \infty$

Sıfır / Kutup Gösterimi



$$X(z) = \frac{1 - 0.64z^{-2}}{1 - 0.2z^{-1} - 0.08z^{-2}} \text{ için kutup-sıfır grafiğini çizelim.}$$

Credit by Sarp Ertürk

Sıfır / Kutup Gösterimi


$$X(z) = \frac{1 - 0.64z^{-2}}{1 - 0.2z^{-1} - 0.08z^{-2}} \text{ için kutup-sıfır grafiğini çizelim.}$$

$$X(z) = \frac{(1 - 0.8z^{-1})(1 + 0.8z^{-1})}{(1 - 0.4z^{-1})(1 + 0.2z^{-1})}$$

Sıfır / Kutup Gösterimi

$X(z) = \frac{1 - 0.64z^{-2}}{1 - 0.2z^{-1} - 0.08z^{-2}}$ için kutup-sıfır grafiğini çizelim.

$$X(z) = \frac{(1 - 0.8z^{-1})(1 + 0.8z^{-1})}{(1 - 0.4z^{-1})(1 + 0.2z^{-1})}$$



Credit by Sarp Ertürk

POLES of $H(z)$



- Find z , where
 - FIR only has poles at $z=0$

$$H(z) \rightarrow \infty$$

POLES of $H(z)$

- Find z , where

- FIR only has poles at $z=0$ $H(z) \rightarrow \infty$

$$H(z) = \frac{z^3 - 2z^2 + 2z - 1}{z^3}$$

Three Poles at : $z = 0$

POLES of $H(z)$

- Find z , where

- FIR only has poles at $z=0$ $H(z) \rightarrow \infty$

$$H(z) = \frac{z^3 - 2z^2 + 2z - 1}{z^3}$$

Three Poles at : $z = 0$

$$\text{Roots : } z = 1, \frac{1}{2} \pm j \frac{\sqrt{3}}{2}$$

POLES of $H(z)$

- Find z , where

- FIR only has poles at $z=0$

$$H(z) \rightarrow \infty$$


$$H(z) = \frac{z^3 - 2z^2 + 2z - 1}{z^3}$$

Three Poles at : $z = 0$


Roots : $z = 1, \frac{1}{2} \pm j \frac{\sqrt{3}}{2}$

$$e^{\pm j\pi/3}$$


PLOT ZEROS in z-DOMAIN




PLOT ZEROS in z-DOMAIN




PLOT ZEROS in z-DOMAIN



PLOT ZEROS in z-DOMAIN



$$H(z) = \frac{z^2}{z^2 - 0.97z + 0.94}$$



```
%% Z transform
```

```
sym z;
```

```
Hz = 1/( 1 - 0.97*z^(-1) + 0.94*z^(-2));
```

```
[N,D] = numden(Hz);
```


```
pay = roots([0 0 100]);
```

```
payda = roots([94 -97 100]);
```

```
%% Z-plane
```

```
zplane(pay, payda);
```

$$H(z) = \frac{z^2}{z^2 - 0.97z + 0.94}$$



```
%% Z transform
```

```
sym z;
```

```
Hz = 1/( 1 - 0.97*z^(-1) + 0.94*z^(-2));
```

```
[N,D] = numden(Hz);
```

```
pay = roots([0 0 100]);
```

```
payda = roots([94 -97 100]);
```

```
%% Z-plane
```

```
zplane(pay, payda);
```

```
%% Freq. Response
```

```
[h,w] = freqz([0 0 100],[94 -97  
100], 'whole', 2001);
```

```
plot(w/pi, 20*log10(abs(h)))
```

```
ax = gca;
```

```
ax.XTick = 0:.5:2;
```

```
xlabel('Normalized Frequency')
```

```
ylabel('Magnitude (dB)')
```

$$H(z) = \frac{z^2}{z^2 - 0.97z + 0.94}$$

```
%% Z transform
```

```
syms a b;
```

```
Hz = 1/( 1 - 0.97*(a+1j*b)^(-1) + 0.94*(a+1j*b)^(-2));
```

```
fsurf(abs(Hz));
```

```
xlim([-1 1]);
```


```
ylim([-1 1]);
```

```
hold on;
```

```
omega = 0:pi/100:2*pi;
```

```
s = exp(-1j*omega);
```

```
plot3(real(s),imag(s),2*ones(201,1),'LineWidth',3);
```



EXERCISE 9.2: Determine the system function $H(z)$ of an FIR filter whose impulse response is

$$h[n] = \delta[n] - 7\delta[n - 2] - 3\delta[n - 3]$$

McClellan, Schafer, and Yoder, *DSP First, 2e*, ISBN 0-13-065562-7.
Prentice Hall, Upper Saddle River, NJ 07458. ©2016 Pearson Education, Inc.



Exercise-1



EXERCISE 9.2: Determine the system function $H(z)$ of an FIR filter whose impulse response is

$$h[n] = \delta[n] - 7\delta[n - 2] - 3\delta[n - 3]$$

SOLUTION to EXERCISE 9.2:

DSP First 2e

$$\begin{aligned} H(z) &= \sum_{n=0}^3 h[n] z^{-n} \\ &= \sum_{n=0}^3 (\delta[n] - 7\delta[n-2] - 3\delta[n-3]) z^{-n} \\ &= z^{-0} - 7z^{-2} - 3z^{-3} \\ H(z) &= 1 - 7z^{-2} - 3z^{-3} \end{aligned}$$

EQUALS ONE AT $n=0$

Exercise-2

EXERCISE 9.5: Use z -transforms to combine the following cascaded systems

$$w[n] = x[n] + x[n - 1]$$

$$y[n] = w[n] - w[n - 1] + w[n - 2]$$

into a single difference equation for $y[n]$ in terms of $x[n]$.

McClellan, Schafer, and Yoder, *DSP First, 2e*, ISBN 0-13-065562-7.
Prentice Hall, Upper Saddle River, NJ 07458. ©2016 Pearson Education, Inc.



Exercise-2

EXERCISE 9.5: Use z -transforms to combine the following cascaded systems

$$w[n] = x[n] + x[n-1]$$

$$y[n] = w[n] - w[n-1] + w[n-2]$$

into a single difference equation for $y[n]$ in terms of $x[n]$.

SOLUTION to EXERCISE 9.5:

DSP First 2e



$$\begin{aligned} w[n] &= x[n] + x[n-1] \\ \downarrow & \quad \downarrow \quad \downarrow \\ W(z) &= X(z) + z^{-1}X(z) \quad \leftarrow \text{DELAY PROPERTY} \\ y[n] &= w[n] - w[n-1] + w[n-2] \\ \downarrow \\ Y(z) &= W(z) - z^{-1}W(z) + z^{-2}W(z) \\ Y(z) &= (1 - z^{-1} + z^{-2})W(z) \\ &= (1 - z^{-1} + z^{-2})(1 + z^{-1})X(z) \\ &= (1 + z^{-3})X(z) \quad \leftarrow \text{MULTIPLY POLYNOMIALS} \\ Y(z) &= X(z) + z^{-3}X(z) \\ \downarrow \quad \downarrow \quad \downarrow \\ y[n] &= x[n] + x[n-3] \end{aligned}$$



Exercise-3



Example 9-9: Consider a DTFT function expressed as

$$H(e^{j\hat{\omega}}) = (1 + \cos 2\hat{\omega})e^{-j3\hat{\omega}}$$

Using the inverse Euler formula for the cosine term gives

$$H(e^{j\hat{\omega}}) = \left(1 + \frac{e^{j2\hat{\omega}} + e^{-j2\hat{\omega}}}{2}\right) e^{-j3\hat{\omega}} = (e^{j\hat{\omega}})^{-3} + \frac{1}{2}(e^{j\hat{\omega}})^{-1} + \frac{1}{2}(e^{j\hat{\omega}})^{-5}$$

Making the substitution $e^{j\hat{\omega}} = z$ gives

$$H(z) = z^{-3} + \frac{1}{2}z^{-1} + \frac{1}{2}z^{-5} = \frac{1}{2}z^{-1} + z^{-3} + \frac{1}{2}z^{-5}$$

If the impulse response of the system is needed, the inverse z -transform gives

$$h[n] = \frac{1}{2}\delta[n-1] + \delta[n-3] + \frac{1}{2}\delta[n-5]$$

McClellan, Schafer, and Yoder, *DSP First, 2e*, ISBN 0-13-065562-7.
Prentice Hall, Upper Saddle River, NJ 07458. ©2016 Pearson Education, Inc.

