

Introduction to Statistical Modeling

Categorical variables

Joris Vankerschaver

Example: BIRNH study

- Epidemiological follow-up study in the mid 80s where nutritional and health data in Belgium were measured ($n = 5,815$)
- **Goal:** effect of smoking on cholesterol
- Since people from different provinces might have a different smoking and dietary behaviour, we want to correct for province
- Possible values of this variable:
 - 1: West Flanders
 - 2: East Flanders
 - 3: Flemish Brabant
 - 7: Antwerp
 - 8: Limburg

A first analysis ...

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	420.48851244	7.684314e+01	5.472037	4.646481e-08
SMOKING	1.60405509	1.384697e+00	1.158416	2.467450e-01
AGE	-14.85347652	4.442769e+00	-3.343292	8.334607e-04
I(AGE^2)	0.34173838	8.359639e-02	4.087956	4.414182e-05
I(AGE^3)	-0.00234162	5.126759e-04	-4.567447	5.045194e-06
SEX	9.77881436	1.290010e+00	7.580415	4.021198e-14
PROVINCE	-1.68519064	2.307958e-01	-7.301653	3.249025e-13

Model implicitly assumes that mean difference in cholesterol

- between Limburg and West-Flanders is 7 times as large as
- the one between East- and West-Flanders

Dummy variables

- Create 4 **dummy variables**

$$P_2 = \begin{cases} 1 & \text{East Flanders} \\ 0 & \text{other} \end{cases}$$

$$P_3 = \begin{cases} 1 & \text{Flemish Brabant} \\ 0 & \text{other} \end{cases}$$

$$P_7 = \begin{cases} 1 & \text{Antwerp} \\ 0 & \text{other} \end{cases}$$

$$P_8 = \begin{cases} 1 & \text{Limburg} \\ 0 & \text{other} \end{cases}$$

Dummy variables

- These 4 dummy variables carry same information as variable PROVINCE:
 - In West Flanders: $(P_2, P_3, P_7, P_8) = (0, 0, 0, 0)$
 - In East Flanders: $(P_2, P_3, P_7, P_8) = (1, 0, 0, 0)$
 - In Flemish Brabant: $(P_2, P_3, P_7, P_8) = (0, 1, 0, 0)$
 - In Antwerp: $(P_2, P_3, P_7, P_8) = (0, 0, 1, 0)$
 - In Limburg: $(P_2, P_3, P_7, P_8) = (0, 0, 0, 1)$
- Each categorical variable with k levels can be transformed into $k - 1$ dummy variables by choosing 1 level as **reference**:
- In R:

```
m <- lm(TCHOL ~ SMOKING + AGE + I(AGE^2) + I(AGE^3)
        + SEX + factor(PROVINCE))
```

Analysis with dummy variables

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	427.509211438	7.681468e+01	5.565462	2.738817e-08
SMOKING	1.691720424	1.383183e+00	1.223063	2.213583e-01
AGE	-15.186319875	4.440163e+00	-3.420217	6.302833e-04
I(AGE^2)	0.347689341	8.354742e-02	4.161581	3.208966e-05
I(AGE^3)	-0.002376407	5.123749e-04	-4.638023	3.599409e-06
SEX	9.777916700	1.288500e+00	7.588602	3.777949e-14
factor(PROVINCE)2	-8.004574314	2.043346e+00	-3.917385	9.060444e-05
factor(PROVINCE)3	-8.332945248	1.487402e+00	-5.602350	2.217790e-08
factor(PROVINCE)7	-11.567195456	1.858041e+00	-6.225478	5.157986e-10
factor(PROVINCE)8	-13.559736351	1.999453e+00	-6.781721	1.312645e-11

How to test for effect of province?

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	427.509211438	7.681468e+01	5.565462	2.738817e-08
SMOKING	1.691720424	1.383183e+00	1.223063	2.213583e-01
AGE	-15.186319875	4.440163e+00	-3.420217	6.302833e-04
I(AGE^2)	0.347689341	8.354742e-02	4.161581	3.208966e-05
I(AGE^3)	-0.002376407	5.123749e-04	-4.638023	3.599409e-06
SEX	9.777916700	1.288500e+00	7.588602	3.777949e-14
factor(PROVINCE)2	-8.004574314	2.043346e+00	-3.917385	9.060444e-05
factor(PROVINCE)3	-8.332945248	1.487402e+00	-5.602350	2.217790e-08
factor(PROVINCE)7	-11.567195456	1.858041e+00	-6.225478	5.157986e-10
factor(PROVINCE)8	-13.559736351	1.999453e+00	-6.781721	1.312645e-11

Necessary to test if multiple coefficients are zero

Partial F-test

Assume we want to compare 2 **nested models**:

- **Complete model (C)** with p_C parameters; e.g., $p_C = 10$ and

$$E(Y|X, P) = \beta_0 + \beta_1 X + \beta_2 P$$

- **Reduced model (R)** with p_R parameters; e.g., $p_R = 6$ and

$$E(Y|X, P) = \beta_0^* + \beta_1^* X$$

- Testing $H_0 : \beta_2 = 0$ is equivalent to testing if complete and reduced model are equal.

Partial F-test

- Under null hypothesis, residual sums of squares of both models will be approximately same.
- **Test statistic:**

$$SSE(R) - SSE(C).$$

- What is distribution under null hypothesis?

$$\frac{SSE(R) - SSE(C)}{p_C - p_R} \div \frac{SSE(C)}{n - p_C} \sim F_{p_C - p_R, n - p_C}.$$

under null hypothesis

Partial F-test in R

Analysis of Variance Table

Model 1: TCHOL ~ SMOKING + AGE + I(AGE^2) + I(AGE^3) + SEX

Model 2: TCHOL ~ SMOKING + AGE + I(AGE^2) + I(AGE^3) + SEX + factor(PRO

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	5480	10460957				
2	5476	10329117	4	131840	17.474	2.943e-14 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1