

# Leveraging 4f-SAMMY-seq protocol in Prokaryotes as a high-resolution technology to study Nucleoid accessibility

Ugo Maria Iannacchero<sup>1</sup>, Elena Chiti<sup>2</sup>, Valentina Rosti<sup>1,3</sup>, Andrea Vannini<sup>2</sup>, Chiara Lanzuolo<sup>1,3</sup>, Davide Roncarati<sup>2</sup>, Eva Maria Pinatel<sup>1</sup>

<sup>1</sup>Institute of Biomedical Technologies, National Research Council, Segrate, Italy ; <sup>2</sup>Department of Pharmacy and Biotechnology (FaBiT), University of Bologna, Bologna, Italy ;

<sup>3</sup>INGM, National Institute of Molecular Genetics "Romeo ed Enrica Invernizzi", Milan, Italy

## • Background

The bacterial chromosome self-organizes into the nucleoid, where Nucleoid Associated Proteins (NAPs) and RNAs regulate DNA accessibility. While bacterial Hi-C protocols have recently provided insights into Nucleoid organization, their use remains technically demanding and limited to a few bacterial species, due to high-sequencing depth requirements, crosslinking biases and species-specific restriction enzymes. Moreover, the limited achievable resolution (>5 kb) fails to capture gene-level interactions, leaving key aspects of nucleoid organization unresolved.

## • A novel epigenomic technology

**4f-SAMMY-seq** (4-fractions Sequential Analysis of MacroMolecules accessibility) is a high-throughput sequencing technology originally developed in mammalian systems [1]. It relies on the **biochemical separation of chromatin into four solubility-based fractions**, each reflecting a different level of DNA accessibility. This enables genome-wide profiling of **both open and closed regions**, and has been demonstrated to reconstruct **chromatin compartmentalization with Hi-C like resolution and accuracy at lower sequencing depth** [2].

## • Why 4f-SAMMY-seq in Prokaryotes?

- Broadly **applicable across different bacterial species**.
- **Does not involve cross-linking or restriction enzymes**.
- **Does not rely on the use of antibodies**.
- **Cost-effective and fast experimental workflow**.
- **Works efficiently with only ~0.5–2M reads per fraction**, given the small size of bacterial genomes.


## • Discussion and next steps

We adapted the **nf-core/sammyseq** [3,4] pipeline to **prokaryotes** by integrating **read processing tools better suited for bacterial genomes**, accommodating non-standard chromosome names **ensuring compatibility with Hi-C tools** and **refining genome binning strategy to support finer-scale resolution** through smaller window-sizes.

We applied it to ***Helicobacter pylori* in exponential growth**, where we observed **distinct signal differences between more and less accessible DNA fractions**, and we obtained **DNA accessibility domains at sub-kilobase resolution** down to the scale of gene-promoters, with as few as **500,000 reads per fraction**.

Ongoing work will focus on **finalizing the integration of the Differential Solubility module** into the pipeline, with further adaptation for prokaryotic genomes. Additionally, we aim to enhance data interpretation by **incorporating COG pathway enrichment module** to better understand the functional impact of nucleoid accessibility changes.


## Methods of analysis



## Analysis output


### 1 STAGE Pre-processing, alignment and filtering

Raw sequencing reads from each solubility fraction are aligned to the reference genome, with quality control metrics including **GC content** and **read alignment distribution** assessed. To evaluate sample similarity and variability, **Principal Component Analysis (PCA)** is performed on the processed coverage profiles.




### 2 STAGE Compartments analysis (Hi-C like)

**Genome-wide coverage profiles** are generated for each fraction, and cross-correlation values among the four fractions are computed using Hi-C analysis tools to generate an **NxN correlation matrix** computed over user-defined genomic bin size. Eigenvector decomposition of this matrix enables identification of genome-wide compartmental organizations leading to the **generation of a consensus profile across replicates** that, when compared across conditions, can be used to individuate compartment transitions.



### 3 STAGE Differential solubility analysis (ChIP-seq like)

**Pairwise relative ratios** between the most representative fractions are calculated to detect continuous changes in DNA accessibility, enabling identification of gene-level alterations.



### References

1. Sebestyén, E., Marullo, F., Lucini, F., Petri, C., Bianchi, A., Valsoni, S., Olivieri, L., Antonelli, L., Gregorini, F., Oliva, G., Ferrari, F., & Lanzuolo, C. (2020). SAMMY-seq reveals early alteration of heterochromatin and deregulation of bivalent genes in Hutchinson-Gilford Progeria Syndrome. *Nature Communications*, 11(1), 6274. <https://doi.org/10.1038/s41467-020-20048-9>
2. Lucini, F., Petri, C., Salvato, E., Pal, K., Rosti, V., Gorini, F., Santarelli, P., Quadri, R., Lembo, G., Graziano, G., Di Patrizio Soldateschi, E., Tagliaferri, L., Pinatel, E., Sebestyén, E., Rotta, L., Gentile, F., Vaira, V., Lanzuolo, C., & Ferrari, F. (2024). Biochemical properties of chromatin domains define genome compartmentalization. *Nucleic Acids Research*, 52(12), e54-e54. <https://doi.org/10.1093/nar/gkac454>
3. Ewels, P. A., Peltzer, A., Fillinger, S., Patel, H., Alnberg, J., Wilms, A., Garcia, M. U., Di Tommaso, P., & Nahnsen, S. (2020). The nf-core framework for community-curated bioinformatics pipelines. *Nature Biotechnology*, 38(3), 276–278. <https://doi.org/10.1038/s41587-020-0439-x>
4. <https://nf-co.re/sammyseq/dev/> (<https://github.com/nf-core/sammyseq/tree/dev>)

### Contacts

✉ ugo.iannacchero@itb.cnr.it  
📧 @ugoiann.bsky.social

