

# Positive Semigroups on $C^\star$ - and $W^\star$ - Algebras

by

Ulrich Groh<sup>1</sup>

## D-I: Basic Results on Semigroups and Operator Algebras

This is not a systematic introduction into the theory of strongly continuous semigroups on  $C^\star$  and  $W^\star$ -algebras. For that we refer to *Bratteli-Robinson (1979)*, *Davies (1976)* and the survey article of *Oseledets (1984)*. We only prepare for the subsequent chapters on spectral theory and asymptotics by fixing the notations and introducing some standard constructions.

### 1.1 Notations

1. By  $M$  we shall denote a  $C^\star$ -algebra with unit  $\mathbb{1}$ .

$$M^{\text{sa}} := \{x \in M : x^* = x\}$$

is the selfadjoint part of  $M$  and

$$M_+ := \{x^*x : x \in M\}$$

the positive cone in  $M$ .

If  $M'$  is the dual of  $M$ , then

$$M'_+ = \{\phi \in M' : \phi(x) \geq 0 \text{ for alle } x \in M_+\}$$

is a weak\*-closed generating cone in  $M'$  and we call

$$S(M) := \{\phi \in M'_+ : \phi(\mathbb{1}) = 1\}$$

---

<sup>1</sup> This is the main part of my "Habilitationsschrift" (accepted by Mathematische Fakultät, Universität Tübingen, 1984). I wish to thank Professor T. Ando for his warm hospitality during a one year stay at the Institute of Applied Electricity, Hokkaido University, Sapporo. My thanks go also to the Alexander von Humboldt Stiftung and to NIHON GAKUJITSU SHENEOKAE, whose support made this stay possible.

the state space of  $M$ . For the theory of  $C^*$ -algebras and related notations we refer to [Pedersen (1979)].

$M$  is called a  $W^*$ -algebra, if there exists a Banach space  $M_*$ , such that its dual space  $(M_*)'$  is (isomorphic to)  $M$ . We call  $M_*$  the predual of  $M$  and  $\phi \in M_*$  a normal linear functional. It is known that  $M_*$  is unique [Sakai (1971), 1.13.3.]. For further properties of  $M_*$  and  $M$  we refer to [Takesaki (1979), Chapter III].

2. A map  $T \in L(M)$  is called positive (in symbols  $T \geq 0$ ) if  $T(M_+) \subseteq M_+$  and  $T$  is called  $n$ -positive ( $n \in \mathbb{N}$ ) if  $T \otimes id_n$  is positive from  $M \otimes M_n$  in  $M \otimes M_n$ , where  $id_n$  is the identity map on the  $C^*$ -algebra  $M_n$  of all  $n \times n$ -matrices. Obviously, every  $n$ -positive map is positive.

We call contraction  $T \in L(M)$  a Schwarz map if  $T$  satisfies the inequality

$$T(x) T(x)^* \leq T(xx^*) \quad (x \in M).$$

It is well known that every  $n$ -positive contraction,  $n \geq 2$  and that every positive contraction on a commutative  $C^*$ -algebra is a Schwarz map [Takesaki (1979), Corollary IV. 3.8.]. As we shall see, the Schwarz inequality is crucial for our investigations.

3. If  $M$  is a  $C^*$ -algebra we assume  $T = (T(t))_{t \geq 0}$  to be a strongly continuous semigroup (abbreviated semigroup) while on  $W^*$ -algebras we consider weak\*-semigroups, i.e. the mapping

$$t \rightarrow T(t)_* x : \mathbb{R}_+ \rightarrow M_*$$

is continuous,  $M_*$  the predual of  $M$ , and every  $T(t)$   $w^*$ -continuous. Note that the preadjoint semigroup

$$T_* = \{T(t)_* : T(t) \in t \geq 0\}$$

is weakly, hence strongly continuous on  $M_*$  (see e.g., Davies (1980), Prop.1.23). We call  $T$  identity preserving if  $T(t) \mathbb{1} = \mathbb{1}$  and of Schwarz type if every  $T(t)$  is a Schwarz map.

For the notations concerning one-parameter semigroups we refer to Part-A. In addition we recommend to compare the results of this section of the book with the corresponding results for commutative  $C^*$ -algebras, i.e. for  $C_0(X)$ ,  $C(K)$  and  $L^\infty(\mu)$  (see Part-B).

## 1.2 A Fundamental Inequality for the Resolvent

If  $(T(t))_t$  is a strongly continuous semigroup of Schwarz maps on a  $C^*$ -algebra  $M$  (resp. a weak\*-semigroup of schwarz type on a  $W^*$ -algebra  $M$ ) with generator  $A$ ,

then the spectral bound  $s(A) \leq 0$ . Then for  $\lambda \in \mathbb{C}$ ,  $\operatorname{Re}(\lambda) > 0$ , there exists a representation for the resolvent  $R(\lambda, A)$  given by the formula

$$R(\lambda, A)x = \int_0^\infty e^{-\lambda t} T(t)x \, dt \quad (x \in M)$$

where the integral exists in the norm topology.

**Theorem 1** *Let  $T = (T(t))_{t \geq 0}$  be a semigroup of schwarz type and  $S = (S(t))_{t \geq 0}$  a semigroup on a  $C^*$ -algebra  $M$  with generators  $A$  and  $B$ , respectively. If*

$$(S(t)x)(S(t)x)^* \leq T(t)xx^* \quad (*)$$

*for all  $x \in M$  and  $t \in \mathbb{R}_+$ , then*

$$(\mu R(\mu, B)x)(\mu R(\mu, B)x)^* \leq \mu R(\mu, A)xx^*$$

*for all  $x \in M$  and  $\mu \in \mathbb{R}_+$ . The same result holds if  $T$  is a weak\*-semigroup of schwarz type and  $S$  is a weak\*-semigroup on a  $W^*$ -algebra  $M$  such that  $(*)$  is fulfilled.*