

# Characterization of Positive Semigroups on $W^*$ -Algebras

Ulrich Groh

**Abstract** Since the positive cone of a  $C^*$ -algebra has non-empty interior many results of Chapter B-II can be applied verbatim to the characterization of the generator of positive semigroups on  $C^*$ -algebras. On the other hand a concrete and detailed representation of such generators has been found only in the uniformly continuous case (see Lindblad (1976)).

A third area of active research has been the following: Which maps on  $C^*$ -algebras (in particular, which derivations) commuting with certain automorphism groups are automatically generators of strongly continuous positive semigroups.

For more information we refer to the survey article of Evans [1].



# Chapter 1

## Characterization of Positive Semigroups on $W^*$ -Algebras

Since the positive cone of a  $C^*$ -algebra has non-empty interior many results of Chapter B-II can be applied verbatim to the characterization of the generator of positive semigroups on  $C^*$ -algebras. On the other hand a concrete and detailed representation of such generators has been found only in the uniformly continuous case (see Lindblad (1976)). A third area of active research has been the following: Which maps on  $C^*$ -algebras (in particular, which derivations) commuting with certain automorphism groups are automatically generators of strongly continuous positive semigroups. For more information we refer to the survey article of Evans [1].

### 1 Semigroups on Properly Infinite $W^*$ -Algebras

The aim of this section is to show that strongly continuous semigroups of Schwarz maps on properly infinite  $W^*$ -algebras are already uniformly continuous. In particular, our theorem is applicable to such semigroups on  $B(H)$ .

It is worthwhile to remark, that the result of Lotz [2] on the uniform continuity of every strongly continuous semigroup on  $L^\infty$  (see A-II, Sec.3) does not extend to arbitrary  $W^*$ -algebras.

*Example 1* Take  $M = B(H)$ ,  $H$  infinite dimensional, and choose a projection  $p \in M$  such that  $Mp$  is topologically isomorphic to  $H$ . Therefore  $M = H \oplus M_0$ , where  $M_0 = \ker(x \mapsto xp)$ . Next take a strongly, but not uniformly continuous, semigroup  $S$  on  $H$  and consider the strongly continuous semigroup  $S \oplus \text{Id}$  on  $M$ .

For results from the classification theory of  $W^*$ -algebras needed in our approach we refer to Sakai [3, 2.2] and Takesaki [4, V.1].

**Theorem 1** *Every strongly continuous one-parameter semigroup of Schwarz type on a properly infinite  $W^*$ -algebra  $M$  is uniformly continuous.*

*Proof.* Let  $T = (T(t)_{t \geq 0})$  be strongly continuous on  $M$  and suppose  $T$  not to be uniformly continuous. Then there exists a sequence  $(T_n) \subset T$  and  $\varepsilon > 0$  such that

$\|T_n - \text{Id}\| \geq \varepsilon$  but  $T_n \rightarrow \text{Id}$  in the strong operator topology. We claim that for every sequence  $(P_k)$  of mutually orthogonal projections and all bounded sequences  $(x_k)$  in  $M$

$$\lim_n \|(T_n - \text{Id})(P_k x_k P_k)\| = 0$$

uniformly in  $k \in \mathbb{N}$ . This follows from an application of the *Lemma of Phillips* and the fact that the sequence  $(P_k x_k P_k)$  is summable in the  $s^*(M, M_*)$ -topology (compare Elliot (1972)).

Let  $(P_k)$  be a sequence of mutually orthogonal projections in  $M$  such that every  $P_k$  is equivalent to 1 via some  $u_k \in M$  [3, 2.2]. Without loss of generality we may assume  $\|(T_n - \text{Id})(u_n)\| \leq n^{-1}$  since the semigroup  $T$  is strongly continuous. Thus we obtained the following:

- (i)  $\lim_n \|(T_n - \text{Id})(P_k x_k P_k)\| = 0$  uniformly in  $k \in \mathbb{N}$  for every bounded sequence  $(x_k)$  in  $M$ .
- (ii) Every projection  $P_k$  is equivalent to 1 via some  $u_k \in M$ .
- (iii)  $\|(T_n - \text{Id})u_n\| \leq n^{-1}$  for all  $n \in \mathbb{N}$ .

For the following construction see A-I,3.6 and D-II,Sec.2. Let

- (i)  $\widehat{M}$  be an ultrapower of  $M$ ,
- (ii) let  $p := \widehat{(P_k)} \in \widehat{M}$ ,
- (iii)  $T := \widehat{(T_n)} \in L(\widehat{M})$
- (iv) and  $u := \widehat{(u_k)} \in \widehat{M}$ .

Then  $T$  is identity preserving and of Schwarz type on  $\widehat{M}$ . Since  $u^*u = p$  and  $uu^* = 1$  it follows  $pu^* = u^*$  and  $(uu^*)x(uu^*) = x$  for all  $x \in \widehat{M}$ . Finally,  $T(pxp) = pxp$  for all  $x \in \widehat{M}$ , which follows from (i), and  $T(u^*) = T(pu^*) = pu^* = u^*$  and  $T(u) = u$ , which follows from (iii). Using the Schwarz inequality we obtain

$$T(uu^*) = T(1) \leq 1 = uu^* = T(u)T(u)^*.$$

Using D-III, Lemma 1.1. we conclude  $T(ux) = uT(x)$  and  $T(xu^*) = T(x)u^*$  for all  $x \in \widehat{M}$ . Hence

$$\begin{aligned} T(x) &= T(uu^*xu^*) = uT(u^*xu)u^* = uT(pu^*xup)u^* \\ &= upu^*xupu^* = uu^*xu^* = x \end{aligned}$$

for all  $x \in \widehat{M}$ . From this we obtain that for every bounded sequence  $(x_k)$  in  $M$

$$\lim_k \|T_k x_k - x_k\| = 0$$

for some subsequence of the  $T_n$ 's and of the  $x_k$ 's. This conflicts with our assumption at the beginning, hence the theorem is proved.  $\square$

## References

1. Evans, D.: Quantum dynamical semigroups, symmetry groups, and locality. *Acta Applicandae Mathematicae* **2**, 333–352 (1984)
2. Lotz, H.P.: Uniform convergence of operators on  $L^\infty$  and similar spaces. *Math. Z.* **190**, 207–220 (1985)
3. Sakai, S.:  $C^*$ -Algebras and  $W^*$ -Algebras. Springer, Berlin-Heidelberg-New York (1971)
4. Takesaki, M.: *Theory of Operator Algebras I*. Springer, New York-Heidelberg-Berlin (1979)