# Fahnenkorrektur: *One-parameter Semigroups of Positive Operators* (2. Aufl.)

Springer LNM 1184, Proof-PDF vom 14.10.2026 (514 PDF-Seiten). Geprüft: Text ab S. 1 bis Index. Titelei, Formeln und Layout wurden nicht geprüft.

**Seitenangaben** = gedruckte Seitenzahl (PDF-Seite = gedruckte Seite + 18).
**Sicherheit:** ● hoch (eindeutiger Fehler) · ◐ mittel (sehr wahrscheinlich, kurz prüfen) · ○ niedrig (Hinweis, Ermessenssache)

Insgesamt **419 Punkte**: 122 ●, 162 ◐, 135 ○.

## A. Übergreifende Punkte

1. **Einleitungsseiten der Teile (S. 2, 136, 264, 408):** Die Bezeichnung „Part A–D“ ist gewollt. Diese Seiten sind aber in einer anderen Schrift gesetzt (Computer Modern statt Times). Falls das nicht beabsichtigt ist, bitte an Springer melden.
2. **Inhaltsverzeichnis (S. xi–xiv):** „Uniformly Ergodic Semig**o**ups“ → „Semig**r**oups“ (Teil IV, 3.2; gleicher Fehler in Überschrift und Kolumnentitel S. 432–436). „Basic Results **of** Spaces $C_0(X)$“ → vermutlich „**on**“ (Teil II, Kap. 1; auch Kolumnentitel).
3. **Kolumnentitel:** Teil I, Kap. 1 heißt im Kolumnentitel „Basic Results on Banach Spaces“ (Kapiteltitel: „… on Semigroups on Banach Spaces“). Teil IV, Kap. 3 schreibt „W∗-algebras“ klein, sonst „W∗-Algebras“. Kurztitel sonst konsistent.
4. **„Part-I“ im Fließtext:** Die Bindestrich-Form ist gewollt. Gelistet sind nur Stellen, an denen sie fehlt (S. 174, zweimal).

   > **Merkposten (noch zu entscheiden):** Im Fließtext steht 357-mal „Part-I/II/III/IV“ mit Bindestrich. Ohne Bindestrich steht es nur auf S. 174 (zweimal: „Part I, Proposition 1.1.9“, „Part I, Corollary 2.1.33“). Kolumnentitel, Teil-Titelseiten, Inhaltsverzeichnis und „Updated Notes Part I–IV“ schreiben dagegen ohne Bindestrich.
   >
   > - **Variante 1:** So lassen, nur S. 174 an „Part-I“ angleichen.
   > - **Variante 2:** Überall „Part I“ ohne Bindestrich (im Englischen üblich, passt zu Titelei und Kolumnentiteln). Das geht nur als Sammelanweisung an Springer: „im gesamten Text ‚Part-I‘ → ‚Part I‘ usw.“
5. **Britische Schreibweise** nur vereinzelt in den Updated Notes (S. 480: characterise, generalisation), sonst amerikanisch. „behaviour“ kommt nur in Literaturtiteln vor, das ist okay.
6. **Einheitlichkeit:** „quasi-compactness“ / „quasicompactness“ (S. 380), „well defined“ / „well-defined“, „Markov-operator“ / „Markov operator“ u. Ä. Siehe die Einzelpunkte unten (Kategorie „Konsistenz“).
7. **Literatur:** Alle Zitate [n] haben einen Eintrag im jeweiligen Literaturverzeichnis, und jeder Eintrag wird zitiert (automatisch geprüft). Unten aufgeführt sind nur inhaltliche Zuordnungsfehler und Formatfragen.
8. **Keine Author Queries** im PDF gefunden. Falls Springer eine separate AQ-Liste geschickt hat, bitte nachreichen.

## B. Einzelkorrekturen

### Teil I, Kap. 1 – Basic Results on Semigroups on Banach Spaces

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 4 | ◐ | is continuous for the strong operator topology on L(E), e.g., lim ‖T(t)f − T(t0)f‖ = 0 | is continuous for the strong operator topology on L(E), i.e., lim … | Grammatik | Definition, kein Beispiel: 'i.e.' statt 'e.g.'. |
| | | the generator is everywhere | | | |

| | | | | | |
|---|---|---|---|---|---|
| 5 | ○ | defined and therefore bounded (use Proposition 1.1.9 (ii)) | (use Proposition 1.1.9 (i)) | Querverweis | Beschränktheit folgt aus Abgeschlossenheit = 1.1.9 (i); bitte prüfen. |
| 10 | ◐ | As summary we obtain | In summary we obtain | Grammatik | |
| 12 | • | Let E to be one of the following function spaces | Let E be one of the following function spaces | Grammatik | |
| 17 | • | generator on a core of its domain (see Proposition 1.1.9 (ii). In particular | (see Proposition 1.1.9 (ii)). In particular | Satz | Schließende Klammer fehlt. |
| 17 | ○ | (Reed & Simon [10, p.28]) | (Reed & Simon [10, p. 28]) | Satz | Ebenso S. 42 'p.252', S. 66 'pp.18-22 ... p.55', S. 67 'p.77'; sonst 'p. 6'. |
| 18 | ○ | There is an easy way how to obtain different (but isomorphic) semigroups | There is an easy way to obtain different (but isomorphic) semigroups | Grammatik | |
| 20 | ◐ | the translation semigroup (T(t))t≥0 where T(t)f(x) := f(x+t) (see Example 1.2.4) | (see Section 1.2.4) | Querverweis | 1.2.4 ist ein Abschnitt, kein Beispiel. |
| 20 | ○ | By Sections 1.2.4 and 1.3.2, the generator A\|F acts as | By Sections 1.2.4 and 1.3.3, the generator A\|F acts as | Querverweis | Restriktion auf Unterraum = Abschnitt 1.3.3 (Subspace Semigroup), nicht 1.3.2 (Rescaled). Bitte prüfen. |
| 21 | ◐ | it is natural to associate (T(t) ')t≥0 its weak*-generator | it is natural to associate with (T(t)')t≥0 its weak*-generator | Grammatik | |
| 23 | • | form a strongly-continuos semigroup with respect to the sup-norm | form a strongly continuous semigroup with respect to the sup-norm | Tippfehler | 'continuos' -> 'continuous'; außerdem kein Bindestrich (sonst überall 'strongly continuous'). |
| 26 | • | In most cases F will be either the Frechét filter or some free ultra filter | In most cases F will be either the Fréchet filter or some free ultrafilter | Tippfehler | Eine Zeile vorher richtig 'Fréchet filter'. |
| 26 | ◐ | (i.e., the filter of sets with finite complement. In most cases F will be ... ultra filter). | (i.e., the filter of sets with finite complement). In most cases F will be ... ultrafilter. | Satz | Klammer schließt an falscher Stelle. |
| 26 | ○ | For the proof we refer to Lemma 1.4. of Derndinger [2]. | For the proof we refer to Lemma 1.4 of Derndinger [2]. | Satz | |

## Teil I, Kap. 2 – Characterization of Semigroups on Banach Spaces

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 32 | • | (which might not be known explicitely) | (which might not be known explicitly) | Tippfehler | |
| 32 | ◐ | 2.1.1 The abstract Cauchy problem / 2.1.2 One-parameter groups / 2.1.4 Holomorphic semigroups … vs. 2.1.3 The Hille-Yosida Theorem, 2.1.8 Perturbation of Generators, 2.1.9 Domains of Uniqueness | Einheitliche Groß-/Kleinschreibung der Unterabschnittsüberschriften (wie in Kap. 1 Title Case, z.B. 'The Abstract Cauchy Problem') | Konsistenz | Betroffen: 2.1.1, 2.1.2, 2.1.4, 2.1.5 (Differentiable semigroups), 2.1.6 (Norm continuous semigroups), 2.1.7 (Compact semigroups). |
| 35 | ◐ | (continuous dependence from the initial value) | (continuous dependence on the initial value) | Grammatik | S. 37 korrekt 'dependence on the initial values'. |
| 36 | • | So by Theorem 1.1.9 (ii) D(A) is a core of B. | So by Proposition 1.1.9 (ii) D(A) is a core of B. | Querverweis | 1.1.9 ist eine Proposition. |
| 36 | ◐ | It is surprising that from condition (b) and (c) in the corollary | It is surprising that from conditions (b) and (c) in the corollary | Grammatik | |
| 38 | • | it follows that T(·)f is measureable. | it follows that T(·)f is measurable. | Tippfehler | |
| 39 | ◐ | (see Davies [11], Goldstein [15] or Pazy [31]) | (see Davies [11], Goldstein [16] or Pazy [31]) | Literatur | [15] ist die Bibliographie, das Lehrbuch ist [16] (so auch S. 32, 53, 68). |
| 40 | ◐ | if and only if it is weakly holomorphic Yosida [40, V.3]. | if and only if it is weakly holomorphic (Yosida [40, V.3]). | Satz | |
| 40 | ○ | There exists r > 0 and β ∈ (−β1, β1) | There exist r > 0 and β ∈ (−β1, β1) | Grammatik | |
| 43 | ○ | We now prove the following extension of Corollary 2.1.12 | We now prove the following extension of Corollary 2.1.12. | Satz | |
| 45 | • | (this follows from Theorem 1.1.6 (ii)) | (this follows from Proposition 1.1.6 (ii)) | Querverweis | 1.1.6 ist eine Proposition. |
| 45 | ○ | Pazy [31, Chapter 3, Theorem 4.7 and 4.8] | Pazy [31, Chapter 3, Theorems 4.7 and 4.8] | Grammatik | |

| | | | | | |
|---|---|---|---|---|---|
| 45 | ○ | the function t ↦ T(t)f is right sided differentiable at t0 | the function t ↦ T(t)f is right-sided differentiable at t0 | Konsistenz | S. 48 und 51: 'right-sided'. |
| 46 | ○ | Proof of Theorem 2.1.19 . It is enough | Proof of Theorem 2.1.19. It is enough | Satz | Leerzeichen vor Punkt, im PDF sichtbar. |
| 47 | • | All together we obtain for $\|r\| \geqslant r0$, All together we obtain for $\|r\| \geqslant r0$, | All together we obtain for $\|r\| \geqslant r0$, | Satz | Satz doppelt gesetzt (im PDF bestätigt). |
| 50 | ○ | (see e.g., Part-II, Corollary 3.2.12) | (see, e.g., Part-II, Corollary 3.2.12) | Satz | Komma nach 'see' fehlt. |
| 51 | • | if and only if the following equivalent assertions holds. | if and only if the following equivalent assertions hold. | Grammatik | |
| 54 | ◐ | Let $\lambda > s(A)$ be so large such that $\|SR(\lambda,A)\| < 1$. | Let $\lambda > s(A)$ be so large that $\|SR(\lambda,A)\| < 1$. | Grammatik | |
| 57 | ○ | Moreover, it follows from Eq. (2.2.1) on the preceding page $p(f) + p(-f) \geqslant p(0) = 0$ | Moreover, it follows from Eq. (2.2.1) on the preceding page that $p(f) + p(-f) \geqslant p(0) = 0$ | Grammatik | |
| 58 | • | Consequently it posseses a σ(E',E)-limit point | Consequently it possesses a σ(E',E)-limit point | Tippfehler | |
| 59 | • | $p(\lambda R(\lambda,A)g) \leqslant p(g)$for all $g \in E$ | $p(\lambda R(\lambda,A)g) \leqslant p(g)$ for all $g \in E$ | Satz | Fehlendes Leerzeichen, im PDF sichtbar. |
| 60 | • | $(2t\, c\|g - Af\| + t^2\, c\|Ag\|)$ (by (2.3)) | (by Eq. (2.2.2)) | Querverweis | Verweis '(2.3)' existiert nicht; gemeint ist offenbar die Abschätzung $\|p(f)\| \leqslant c\|f\|$ (2.2.2). |
| 60 | ○ | A continuous sublinear function $p : E \to R$ is called half-norm if | is called a half-norm if | Grammatik | Analog folgt 'is called a strict half-norm'. |
| 60 | ○ | (by Eq. (2.2.5) on page 58 ) | (by Eq. (2.2.5) on page 58) | Satz | Leerzeichen vor Klammer, im PDF sichtbar. |
| 61 | ◐ | (this is equivalent to the fact that ... such that $\phi = \phi 1 - \phi 2$ (see Batty & Robinson [3] or Schaefer [37, Chapter V]). | (this is equivalent to the fact that ... such that $\phi = \phi 1 - \phi 2$; see Batty & Robinson [3] or Schaefer [37, Chapter V]). | Satz | Klammern nicht ausgeglichen. |

| | | | | | |
|---|---|---|---|---|---|
| 62 | ◐ | Finally, from 2.2.6 it follows that (T(t)) is p-contractive. | Finally, from Theorem 2.2.6 it follows that (T(t)) is p-contractive. | Querverweis | |
| 63 | ○ | We conclude this section indicating the results for the complex case. | We conclude this section by indicating the results for the complex case. | Grammatik | |
| 64 | ◐ | Therefore T is not uniformly continuous however, $(T(t) - Id)^2 = 0$ | Therefore T is not uniformly continuous; however, $(T(t) - Id)^2 = 0$ | Satz | |
| 65 | ○ | Then there exists a bounded sequence (xn) in E and a bounded sequence (x'n) in E' | Then there exist a bounded sequence (xn) in E and a bounded sequence (x'n) in E' | Grammatik | |
| 67 | ◐ | Notes ... Section 1: The abstract Cauchy problem is treated ... | Section 2.1: ... | Konsistenz | In den Notes zu Kap. 2 heißen die Abschnitte 'Section 1/2/3' (S. 67/68), im Buch sonst 2.1, 2.2, 2.3. |
| 67 | ○ | is a Cauchy-sequence for this norm | is a Cauchy sequence for this norm | Tippfehler | |
| 67 | ○ | In Bourgain [7] it is shown, that $H^\infty(D)$ is a Grothendieck space | In Bourgain [7] it is shown that $H^\infty(D)$ is a Grothendieck space | Grammatik | |
| 68 | ◐ | The observation that eventually norm continuity is preserved | The observation that eventual norm continuity is preserved | Grammatik | |
| 68 | ○ | in the book of Pazy [31] who discovered Theorem 2.1.16 and 2.1.17. | who discovered Theorems 2.1.16 and 2.1.17. | Grammatik | |
| 68 | ○ | has the advantage to yield the same permanence | has the advantage of yielding the same permanence | Grammatik | |
| 68 | ○ | A complete information on the general theory can be obtained | Complete information on the general theory can be obtained | Grammatik | |
| 68 | ○ | Bénilan & Picard [5] and Picard [34], where the two last consider non-linear semigroups | where the last two consider non-linear semigroups | Grammatik | |
| 69 | ◐ | [5] P. Bénilan & C. Picard: “Quelques aspects non lineaires du principe du maximum”. Séminaire de Théorie du Potentiel. | ... “Quelques aspects non linéaires du principe du maximum”. ... Springer (1979). | Literatur | Akzent fehlt; Jahr ohne Klammern (sonst '(Jahr)'); ggf. 'Lecture Notes in |

| | | | | | |
|---|---|---|---|---|---|
| | | Vol. 713. Springer 1979. | | | Math.' statt 'Vol.'. |
| 69 | ○ | [7] J. Bourgain: H∞ is a Grothendieck space. Studia Math. 75 (1982) 192–226. | [7] J. Bourgain: H∞ is a Grothendieck space. Studia Math. 75 (1983) 193–216. | Literatur | Jahr/Seiten nach meiner Kenntnis 1983, 193–216; bitte prüfen. |
| 69 | ○ | Proceedings of the Special year in analysis. Springer (1983). | Proceedings of the Special Year in Analysis. Springer (1983). | Literatur | |
| 70 | • | [30] J. J. Moreau: Fonctionelles convexes. Séminaire sur les équations aux derivées partielles. | [30] J. J. Moreau: Fonctionnelles convexes. Séminaire sur les équations aux dérivées partielles. | Literatur | |
| 70 | ◐ | [34] C. Picard: Opérateurs ϕ-accretifs et génération des semi-groupes non linéaires. | [34] C. Picard: Opérateurs ϕ-accrétifs et génération des semi-groupes non linéaires. | Literatur | |
| 70 | ◐ | Ed. by K. D. Bierstedt & E. Fuchssteiner. North Holland 1984. | Ed. by K. D. Bierstedt & B. Fuchssteiner. North-Holland (1984). | Literatur | Herausgeber ist Benno Fuchssteiner; Jahresangabe sonst in Klammern. |
| 70 | ○ | [25] H. P. Lotz: “Uniform convergence of operators on L∞”. 1982. | Unvollständiger Eintrag (z.B. 'Preprint (1982)' ergänzen) | Literatur | |

## Teil I, Kap. 3 – Spectral Theory

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 71 | ○ | called spectral mapping theorem. | called the spectral mapping theorem. | Grammatik | |
| 71 | ○ | These — or similar — relations will be of great use | These—or similar—relations will be of great use | Konsistenz | Sonst Geviertstrich ohne Leerzeichen (S. 9 'are—at least in many aspects—', S. 28, S. 72). |
| 76 | ◐ | = (dist($x$, $\sigma(A)$))^−1 (see Proposition 3.2.5 (ii)) | (see Proposition 3.2.5 (i)) | Querverweis | Die Formel r(R(λ0)) = dist^−1 steht in 3.2.5 (i), nicht (ii). |
| 77 | ◐ | using the notion essential range, see Goldstein [8] | using the notion of essential range | Grammatik | |
| | | we recall the well known | we recall the well-known | | Leerzeichen vor |

| | | | | | |
|---|---|---|---|---|---|
| 79 | • | resolvent equation . | resolvent equation. | Satz | dem Punkt (im PDF sichtbar). |
| 82 | • | the spectral mapping theorem for the resolvent (Theorem 3.2.5) | (Proposition 3.2.5) | Querverweis | 3.2.5 ist eine Proposition; ebenso in den Notes S. 108 'Theorem 3.2.5'. |
| 83 | ◐ | Only by adding appropriate hypotheses we will achieve spectral decompositions | Only by adding appropriate hypotheses will we achieve spectral decompositions | Grammatik | Inversion nach 'Only by ...'. |
| 83 | ○ | follows from the Spectral Inclusion Theorem (see 3.6.2) below. | follows from the Spectral Inclusion Theorem (see Proposition 3.6.2 below). | Querverweis | |
| 86 | ◐ | We review these constructions under the spectral theoretical point of view | from the spectral theoretical point of view | Grammatik | |
| 87 | ◐ | it follows that $(\lambda - A/)\ f\hat{} = g\hat{} = 0$, i.e., $\lambda \in \sigma(A/)$ | ..., i.e., $\lambda \in \sigma(A/)$. | Satz | Schlusspunkt fehlt (im PDF geprüft). |
| 88 | ○ | either well known or has been stated in 3.2.2(ii). | stated in Proposition 3.2.2 (ii). | Querverweis | |
| 92 | ○ | the representations Eq. (3.5.3) on the previous page and Eq. (3.5.4) on the preceding page | the representations (3.5.3) and (3.5.4) on the previous page | Satz | Automatische varioref-Zusätze wirken doppelt/unbeholfen; tritt im Bereich häufig auf (z. B. S. 128 sechsmal 'Eq. (4.2.1) on the previous/preceding page'). |
| 96 | ◐ | Since $R\sigma(A) = P\sigma(A*)$ and $R\sigma(T(t)) = P\sigma(T(t)*)$ (see Proposition 3.4.4) | (see Section 3.4.1) | Querverweis | Proposition 3.4.4 behandelt das F-Produkt; die Aussage über den adjungierten Halbgruppen steht in 3.4.1. |
| 98 | ◐ | on Banach spaces $E1$, $E2$ with generator $A1$, $A2$. | with generators $A1$, $A2$. | Grammatik | |
| 99 | ◐ | (see Theorem 2.1.19 ). By | (see Theorem 2.1.19). By | Satz | Leerzeichen vor ')' bzw. ','; gleiches Muster im PDF bestätigt auf S. 121 ('Corollary 3.6.7 )', 'Theorem 2.1.6 )'), |

| | | | | | |
|---|---|---|---|---|---|
| | | taking | taking | | außerdem S. 123 'Theorem 3.6.3 )', S. 124 'Corollary 3.6.4 ,', S. 125 'Corollary 2.1.12 )'. |
| 100 | • | a "multiplication operator" | a “multiplication operator” | Satz | Gerade Anführungszeichen statt typografischer (sonst überall “...”). |
| 102 | ○ | That $T(\varphi)$ is well-defined follows | well defined (einheitlich) | Konsistenz | Wechsel zwischen 'well defined' (S. 85, 87) und 'well-defined' (S. 102, 128) bzw. 'well known' / 'well-known' (S. 84, 138). |
| 104 | • | (cf. Formula Eq. (1.3.1) on page 19) | (cf. Eq. (1.3.1) on page 19) | Tippfehler | 'Formula' und 'Eq.' doppelt. |
| 104 | ◐ | By the rescaling procedure Section 1.3.2 we may assume | By the rescaling procedure (see Section 1.3.2) we may assume | Grammatik | Ebenso S. 107: 'use the rescaling procedure Section 1.3.2'. |
| 105 | ○ | is Cesàro-summable ... Fourier series is Cesàro summable | einheitlich 'Cesàro summable' (oder durchgehend mit Bindestrich) | Konsistenz | S. 104/105 wechseln 'Cesàro summable' und 'Cesàro-summable'; ebenso 'Fejér’s Theorem' (S. 104) vs. 'Fejér’s theorem' (S. 105) und 'Hahn-Banach theorem' (S. 76, 91) vs. 'Hahn-Banach Theorem' (S. 93). |
| 107 | • | for some $k \in$ Z otherwise if $(\|R(\mu k, A)\|)k\in$Z is unbounded. | for some $k \in$ Z or if $(\|R(\mu k, A)\|)k\in$Z is unbounded. | Grammatik | 'either ... or' statt 'either ... otherwise'. |
| 107 | • | The Example 3.1.3 above in combination with Corollary 3.1.3 shows | Example 3.1.3 above in combination with Corollary 3.1.2 (?) shows | Querverweis | Ein 'Corollary 3.1.3' gibt es nicht (3.1.3 ist das Example). Bitte richtiges Ziel prüfen. |
| 107 | ◐ | (use the rescaling procedure Section 1.3.2) | (use the rescaling procedure, see Section 1.3.2) | Grammatik | |
| 108 | • | The spectral mapping theorem for the resolvent (Theorem 3.2.5) is well | (Proposition 3.2.5) | Querverweis | |

| | | | | | |
|---|---|---|---|---|---|
| | | known | | | |
| 108 | • | can be found in Kato [16], Chap.III,§ 6.4]. | can be found in Kato [16, Chap. III, § 6.4]. | Satz | Klammer falsch gesetzt (öffnende eckige Klammer fehlt), Leerzeichen fehlen. |
| 108 | • | Applications to isolated singularities like Example 3.6 are discussed | like Example 3.3.6 are discussed | Querverweis | Example 3.6 existiert nicht; gemeint ist Example 3.3.6 (Isolated singularities and poles). |
| 108 | ○ | may be violated in various ways [l.c.,Sec.23.16] | [l.c., Sec. 23.16] | Satz | Fehlende Leerzeichen; ähnlich 'Chap.VII' (S. 108), 'Chap.I' (S. 83), 'Chap.8.1' (S. 93), 'Sec.23.16' (S. 101), 'Sec.8' (S. 108) – uneinheitlich zu 'Chap. III, §6.5'. |
| 108 | ○ | Matrix valued multiplication semigroups appear as solution, via Fourier transformation, of systems | appear as solutions, via Fourier transformation, of systems | Grammatik | |
| 109 | ◐ | F. Combes & C. Delaroche: Representations des groupes localement compacts | Représentations des groupes localement compacts | Literatur | Akzent fehlt im französischen Titel. |
| 109 | ◐ | G. Greiner: Some applications of Fejer's theorem to one-parameter semigroups | Some applications of Fejér's theorem | Literatur | Im Text 'Fejér' mit Akzent. |
| 109 | ◐ | G. Greiner, J. Voigt & M. Wolff: On the spectral bound ... J. Operator Theory (1981) 245–256. | J. Operator Theory 5 (1981) 245–256. | Literatur | Bandnummer fehlt; gleicher Eintrag in Kap. 4, Ref. [10] (S. 132). |
| 109 | ○ | Semesterbericht Funktionalanalysis. Universität Tübingen, 1985 pp. 33–49. | Semesterbericht Funktionalanalysis. Universität Tübingen, 1985, pp. 33–49. | Literatur | |
| | | M. P. H. Wolff: "A remark on the spectral bound ... Proceedings of the Conference in | M. P. H. Wolff: A remark on the spectral bound ... In: Functional Analysis and Approximation (Proc. Conf. | | Anführungszeichen umschließen Titel samt Tagungsangabe; Format weicht von |

| | | | | | |
|---|---|---|---|---|---|
| 110 | ◐ | Oberwolfach 1980”. Functional Analysis and Approximation. ... Birkhäuser 1981 pp. 39–50. | Oberwolfach 1980), ed. by P. L. Butzer, B. Sz.-Nagy & E. Görlich. Birkhäuser (1981) 39–50. | Literatur | den übrigen Einträgen ab. Gleicher Eintrag Kap. 4, Ref. [26] (S. 133). |

## Teil I, Kap. 4 – Asymptotics of Semigroups on Banach Spaces

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 114 | • | for all $f \in C0(Rn)$; see also Example 3.1.7. | Verweisziel prüfen (z. B. Example 3.2.3/3.2.4) | Querverweis | In Abschnitt 3.1 gibt es nur 3.1.1–3.1.4; 'Example 3.1.7' existiert nicht. |
| 114 | ○ | beyond the class of eventually norm-continuous semigroups | eventually norm continuous semigroups | Konsistenz | Kap. 3 schreibt durchgehend 'norm continuous' ohne Bindestrich; Kap. 4 mit (auch S. 121). |
| 114 | ○ | (ii) Note that the condition ... (iii) Rescaling this semigroup | (ii) Note that ... / (iii) Rescaling ... (ohne Einzug) | Satz | Unbeabsichtigter Einzug vor den Aufzählungspunkten (im PDF sichtbar); ähnlich S. 79, 87, 121 ('Due to the difficulty'). |
| 117 | ○ | which implies that $g \in \ker(A) = \{0\}$ or $g = 0$. | which implies that $g \in \ker(A) = \{0\}$, i.e., $g = 0$. | Grammatik | 'or' im Sinne von 'bzw./d.h.' (Germanismus); ähnlich S. 126: 'Therefore, $D \subseteq G$ or $G = E$'. |
| 118 | ○ | Proof. (Proof of Theorem 4.1.4) The equality | Proof of Theorem 4.1.4. The equality | Satz | Doppeltes 'Proof'. |
| 119 | • | Example 4.1.2(c) shows that uniform exponential stability is not equivalent to | Example 4.1.2 (ii) shows | Querverweis | Example 4.1.2 hat die Teile (i)–(iii); gemeint ist (ii). Ebenso S. 121: 'Example 4.1.2(c) produces a semigroup'. |
| 121 | • | On the other hand, Example 4.1.2(c) produces a semigroup | Example 4.1.2 (ii) produces a semigroup | Querverweis | |
| 121 | ◐ | holomorphic, or nilpotent ones—the spectral mapping theorem | holomorphic, or nilpotent ones—the spectral mapping theorem | Satz | Leerzeichen nach Geviertstrich, davor keins (im PDF sichtbar). |

| | | | | | |
|---|---|---|---|---|---|
| 122 | ○ | see Widder [24, p.66]; see also the notes in this section. | see also the notes at the end of this chapter. | Querverweis | |
| 124 | ○ | By a classical theorem of Laplace transform theory, (for a proof of the vector-valued version one may follow Widder [25, p.196]), we conclude | By a classical theorem of Laplace transform theory (for a proof ... Widder [25, p. 196]), we conclude | Satz | Überzähliges Komma vor der Klammer. |
| 125 | • | The Laplacian Δ generates a bounded holomorphic semigroups on $Lp$(R$n$) | generates a bounded holomorphic semigroup on | Tippfehler | |
| 125 | ○ | Furthermore the following statements are equivalent, | Furthermore, the following statements are equivalent. | Satz | |
| 126 | ○ | Therefore, $D \subseteq G$ or $G = E$ since $D$ is dense in $E$. | Therefore, $D \subseteq G$, and hence $G = E$ since $D$ is dense in $E$. | Grammatik | |
| 127 | ○ | to be weakly stable if ... or as weakly uniformly stable if the above holds | to be weakly stable if ... or weakly uniformly stable if the above holds | Grammatik | |
| 128 | ○ | (see Pazy [17, Chapter 4,2.4]) | (see Pazy [17, Chapter 4, 2.4]) | Satz | |
| 131 | ◐ | Theorem 4.1.3 (ii) is a semigroup version of Theorem 1.2.7 and 1.2.8 in Doetsch [7]. | of Theorems 1.2.7 and 1.2.8 in Doetsch [7]. | Grammatik | |
| 131 | ○ | was observed by many authors, see for example, Balakrishnan [1, p.178], or Benchimol [2]. | see, for example, Balakrishnan [1, p. 178] or Benchimol [2]. | Satz | |
| 132 | ◐ | A. Balakrishnan: Applied Functional Analysis. Application of Mathematics. Springer (1976). | A. V. Balakrishnan: Applied Functional Analysis. Applications of Mathematics. Springer (1976). | Literatur | Reihenname lautet 'Applications of Mathematics'. |
| 132 | ◐ | G. Doetsch: Handbuch der Laplace Transformation, Band I. | Handbuch der Laplace-Transformation, Band I. | Literatur | |
| 132 | ○ | R. Datko: Extending a theorem of A.M. Liapunov to Hilbert space. | Extending a theorem of A. M. Liapunov to Hilbert space. | Literatur | |
| 132 | ○ | Exponential stability of positive semigroups in a linear topological space I,II. | ... in a linear topological space I, II. | Literatur | |
| | | F. Neubrander: Laplace transform and asymptotic | | | Band 12 der |

| | | | | | |
|---|---|---|---|---|---|
| 132 | ○ | behavior of strongly continuous semigroups. Houston J. Math. 12 (1985) 549–561. | Houston J. Math. 12 (1986) 549–561. | Literatur | Houston J. Math. ist meines Wissens von 1986 – bitte prüfen. |
| 133 | ◐ | M. Slemrod: Asymptotic behavior of $C0$-semigroups ... Indiana Univ. Math. J. 25 (1976) 783–892. | Indiana Univ. Math. J. 25 (1976) 783–792. | Literatur | Seitenumfang 110 Seiten unplausibel; der Artikel umfasst 783–792. |

## Teil II, Kap. 1 – Basic Results on Spaces $C_0(X)$

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 137 | ○ | Basic Results of Spaces $C0(X)$ | Basic Results on Spaces $C0(X)$ | Grammatik | Präposition; betrifft auch den Kolumnentitel. |
| 137 | ○ | 1.1 Algebraic and Order-Structure: Ideals and Quotients | 1.1 Algebraic and Order Structure: Ideals and Quotients | Satz | |
| 139 | • | This identification is a topological isomorphisms with respect to the weak*-topology | This identification is a topological isomorphism with respect to the weak* topology | Tippfehler | |
| 139 | • | (i) positive (notation: $T \geqslant 0$) if $T f$ is a positive whenever $f$ is positive, | if $Tf$ is positive whenever $f$ is positive, | Tippfehler | |
| 139 | ◐ | (iii) Markov-operator if $X$ and $Y$ are compact | (iii) Markov operator if $X$ and $Y$ are compact | Konsistenz | Auf S. 140 'Markov operator' ohne Bindestrich. |

## Teil II, Kap. 2 – Characterization of Positive Semigroups on $C_0(X)$

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 142 | ◐ | similar to a derivation perturbated by a multiplication operator | similar to a derivation perturbed by a multiplication operator | Tippfehler | |
| 143 | • | It will be essential that K is compact for all what follows | It will be essential that K is compact for all that follows | Grammatik | |
| 146 | ◐ | is not positive (see Greiner et al. [16] ). | is not positive (see Greiner et al. [16]). | Satz | Leerzeichen vor schließender Klammer |
| 148 | • | The next result is a strengthened form of Theorem 1.6. | The next result is a strengthened form of Theorem 2.1.6. | Querverweis | Nicht verlinkter Altverweis; gemeint ist Theorem 2.1.6 (S. 144) |
| 148 | ◐ | Thus (P) is not a "metric" condition | Thus (P) is not a “metric” condition | Satz | Gerade statt typografische Anführungszeichen |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | (sonst "…" verwendet) |
| 150 | ◐ | then by Part-I, Section 1.3.2, the operator m · A = ‖m‖∞ B also generates a positive semigroup). | … also generates a positive semigroup. | Satz | Schließende Klammer ohne öffnende (Satz geht auf S. 151 weiter) |
| 152 | • | preceding page succesively to B = m^{1/k} · A | preceding page successively to … | Tippfehler | |
| 154 | • | Then A is the generator of positive semigroup. | Then A is the generator of a positive semigroup. | Grammatik | |
| 155 | ◐ | A semigroup (T(t))t≥0 is called lattice semigroup if | A semigroup (T(t))t≥0 is called a lattice semigroup if | Grammatik | |
| 155 | ◐ | a concrete representation of lattice-semigroups | a concrete representation of lattice semigroups | Konsistenz | Sonst durchgehend ohne Bindestrich |
| 156 | ◐ | right-sided Gateaux-differentiable at every point f ∈ G (vs. 'right-sided Gateaux differentiable' im selben Absatz) | einheitlich 'Gateaux differentiable' (bzw. ggf. 'Gâteaux') | Konsistenz | Auf S. 155/156 gemischt mit und ohne Bindestrich (auch in Prop. 2.2.3) |
| 159 | ○ | For A = Δ (the Laplacian) this is a version of the classical Kato's inequality. | … this is a version of the classical Kato inequality. | Grammatik | |
| 160 | ◐ | Remark Although we assume that (T(t))t≥0 is a real semigroup, i.e., T(t)C(K,R) ⊆ C(K,R) for all t ≥ 0. It is important for the proof | … for all t ≥ 0, it is important for the proof | Grammatik | Nebensatz mit 'Although' als eigener Satz abgetrennt |
| 161 | • | hence T(t) is so as well (cf. Remark 2.1). | hence T(t) is so as well (cf. Remark 2.2.1). | Querverweis | Remark 2.1 existiert nicht; gemeint Remark 2.2.1 (S. 155) |
| 161 | • | we denote by dϕ he linear form on C[−1, 0] | we denote by dϕ the linear form on C[−1, 0] | Tippfehler | |
| 161 | ◐ | The operator Amax on the (real or complex space) C[−1, 0] | The operator Amax on the (real or complex) space C[−1, 0] | Grammatik | |
| 162 | ◐ | Let A on (the real or complex) space C[−1, 0] be given by | Let A on the (real or complex) space C[−1, 0] be given by | Grammatik | |
| 162 | ◐ | A satisfies Kato's Equality Eq. (2.2.9) | A satisfies Kato's equality Eq. (2.2.9) | Konsistenz | Sonst 'Kato's equality' klein |
| 163 | • | this will enable us, to give a detailed description | this will enable us to give a detailed description | Satz | |
| 163 | ◐ | A mapping ϕ : [0, ∞) × K → K is called semiflow if | … is called a semiflow if | Grammatik | |
| 163 | ○ | the space of all real valued continuous functions on K | the space of all real-valued continuous functions on K | Konsistenz | Sonst 'real-valued' |
| 164 | ◐ | An operator δ on C(K) is called derivation if | An operator δ on C(K) is called a derivation if | Grammatik | |
| 166 | ◐ | Theorem 2.3.6 An operator A is generator of a lattice semigroup | An operator A is the generator of a lattice semigroup | Grammatik | |

| | | | | | |
|---|---|---|---|---|---|
| 166 | ○ | In fact, by Eq. (2.1.6) T0(t)f = lim e^{−t/n·h} · T(t/n)f | Verweis prüfen (Eq. (2.1.6) ist ϕx ∈ dpu(f); gemeint vermutlich die Trotter-Produktformel aus Part I) | Querverweis | |
| 167 | • | is closable (by Corollary 2.1.11 ); but | is closable (by Corollary 2.1.11); but | Satz | Im PDF bestätigt |
| 169 | ◐ | Given a compact neighbourhood V of x0 | neighbourhood/neighborhood vereinheitlichen | Konsistenz | S. 169, 179: neighbourhood; S. 180 (3x): neighborhood; außerdem 'behavior' (S. 187) amerikanisch |
| 174 | ◐ | by Part I, Proposition 1.1.9 (or Part I, Corollary 2.1.33) | by Part-I, Proposition 1.1.9 (or Part-I, Corollary 2.1.33) | Konsistenz | Im Fließtext sonst durchgehend „Part-I“ mit Bindestrich, siehe Merkposten A.4. |
| 178 | ◐ | This does no longer hold for functions which are not admissible | This no longer holds for functions which are not admissible | Grammatik | |
| 179 | • | is closed in (a, b) Thus (a, b) ∖ K is the union | is closed in (a, b). Thus (a, b) ∖ K is the union | Satz | Punkt fehlt, im PDF bestätigt |
| 179 | ◐ | Thus αn is strictly monotonous. | Thus αn is strictly monotone. | Tippfehler | 'monotonous' = eintönig; mathematisch 'monotone'/'monotonic' |
| 183 | • | It is easy to verify, that the generator of this group is | It is easy to verify that the generator of this group is | Satz | |
| 183 | ◐ | The operator δ̃m is generator of an automorphism group on C[a, b] | The operator δ̃m is the generator of an automorphism group | Grammatik | |
| 184 | • | we refer to the corresponding notes in Part-III, Chapter 2, Theorem 2.1.6 and 2.1.8 are due to Arendt et al. [4] | ... in Part-III, Chapter 2. Theorems 2.1.6 and 2.1.8 are due to Arendt et al. [4] | Satz | Komma statt Punkt; außerdem Plural 'Theorems' |
| 184 | • | Infinite-dimensional Systems, Retahof 1983. | Infinite-dimensional Systems, Retzhof 1983. | Literatur | Tagungsort Retzhof (LNM 1076) |
| 184 | ◐ | as perturbation of a derivation (Theorem 2.3.5 and 2.3.6) | (Theorems 2.3.5 and 2.3.6) | Grammatik | |
| 184 | ◐ | positive groups on C0(X) (Theorem 2.3.14 ) was obtained | (Theorem 2.3.14) was obtained | Satz | |
| 184 | ◐ | to satisfy the "generalized Osgood condition" | to satisfy the “generalized Osgood condition” | Satz | Gerade Anführungszeichen |
| 184 | ◐ | Uhlig [30] determined all semi-flows on an interval. | Uhlig [30] determined all semiflows on an interval. | Konsistenz | Im Kapitel sonst 'semiflow' |
| 184 | ◐ | has been obtained by Kurose in a series of papers Kurose [21, 22, 23]. | has been obtained by Kurose in a series of papers [21, 22, 23]. | Grammatik | Name doppelt |
| 184 | ○ | (see Part-III, Section 2.5), for general Banach spaces and C0(X) in particular the problem has been considered | (see Part-III, Section 2.5); for general Banach spaces ... | Satz | |

| | | | | | |
|---|---|---|---|---|---|
| 184 | ○ | Our approach is inspired by Laubenfels [24] | Our approach is inspired by de Laubenfels [24] | Konsistenz | Im Literaturverzeichnis 'R. de Laubenfels' |
| 184 | ○ | In the sequel of Batty's work (loc.cit.) | (loc. cit.) | Satz | |
| 184 | ○ | [2] W. Arendt: "Generators of positive semigroups". … Vol. 1076. Springer 1984 pp. 1–15. / [30] H. Uhlig: "Derivationen und Verbandshalbgruppen". | Titel ohne Anführungszeichen, Jahr in Klammern wie bei den übrigen Einträgen (z. B. Lecture Notes in Math. 1076, Springer (1984) 1–15) | Literatur | Formatierung uneinheitlich |
| 185 | • | [11] P. Collatz: Einschließungssatz für die charakteristischen Zahlen von Matrizen. | [11] L. Collatz: Einschließungssatz … | Literatur | Lothar Collatz |
| 185 | ◐ | [16] G. Greiner, J. Voigt & M. Wolff: … J. Operator Theory (1981) 245–256. | J. Operator Theory 5 (1981) 245–256. | Literatur | Bandangabe fehlt |
| 185 | ◐ | [24] R. de Laubenfels: Well behaved derivation on C[0,1]. | R. de Laubenfels: Well-behaved derivations on C[0,1]. | Literatur | |
| 185 | ○ | [19] E. Kamke: Über die partielle Differentialgleichung. Math. Z. 41 (1932) 56–66. | Titel/Angaben prüfen (Titel wirkt unvollständig) | Literatur | |
| 185 | ○ | [20] T. Kato: Schrödinger operators with singular potentials. Israel J. Math. 13 (1973) 135–148. | Israel J. Math. 13 (1972) 135–148 | Literatur | Band 13 erschien 1972 – bitte prüfen |
| 186 | ○ | [27] G. Lumer: Problème de Cauchy pour opérateurs locaux et "changement de temps". Ann. Inst. Fourier (Grenoble) 23 (1974) 409–466. | Ann. Inst. Fourier (Grenoble) 25 (1975) 409–446 | Literatur | Band/Jahr/Seiten wohl aus [26] übernommen – bitte prüfen; [26] ist Bd. 23 (1973) |

## Teil II, Kap. 3 – Spectral Theory of Positive Semigroups on $C_0(X)$

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 187 | ◐ | Krein, Krasnoselski'i, Lotz | Krein, Krasnosel'skii, Lotz | Tippfehler | |
| 187 | ◐ | (see Chapt.V of Schaefer [27]) … (Schaefer [27, V.Proposition 4.1]) | (see Chap. V of Schaefer [27]) … (Schaefer [27, V, Proposition 4.1]) | Satz | Leerzeichen fehlen; Abkürzung uneinheitlich (Chapt./Chap./Chapter, vgl. S. 150 'Chapter I', S. 209 'Chap.I') |
| 188 | ○ | We may assume λ0 > ω0(A) then R(λ0, A) ⩾ 0, and iterating | We may assume λ0 > ω0(A); then R(λ0, A) ⩾ 0, and iterating | Satz | |
| 192 | ○ | equation Eq. (3.1.16) implies that | Eq. (3.1.16) implies that | Satz | Doppelung 'equation Eq.' |

| | | | | | |
|---|---|---|---|---|---|
| 202 | • | A real spectral value λ0 of a generator A is called a dominant provided that | … is called dominant provided that | Grammatik | |
| 204 | • | unless ϕ = r0δ0 for some r0 ∈ R.) | unless ϕ = r0δ0 for some r0 ∈ R. | Satz | Schließende Klammer ohne öffnende, im PDF bestätigt |
| 204 | ◐ | It follows from Eq. (3.2.25) and Eq. (3.2.26) that for λ > ‖ϕ‖, the resolvent R(λ, Aϕ) exists | It follows from Eq. (3.2.27) and Eq. (3.2.28) that … | Querverweis | Die Abschätzung folgt aus (3.2.27)/(3.2.28), nicht aus (3.2.25)/(3.2.26) |
| 205 | ◐ | hence it maps the unit ball in an order interval | hence it maps the unit ball into an order interval | Grammatik | |
| 206 | ◐ | can be extended to systems of functional differential equations even the infinite dimensional case. | … functional differential equations, even to the infinite-dimensional case. | Grammatik | Wort fehlt |
| 206 | ◐ | Then the ideal I generated by {T(t)f : t ⩾ 0} satisfy T ≠ {0} | Then the ideal I generated by {T(t)f : t ⩾ 0} satisfies I ≠ {0} | Grammatik | Verbform; zudem T statt I (auch in der Folgezeile) |
| 207 | ◐ | (e) =⇒ (f): If I is a R(λ, A)-invariant ideal | (e) =⇒ (d): If I is an R(λ, A)-invariant ideal | Querverweis | Beweiskette a⇒b⇒c⇒f⇒e⇒d⇒a; '(e) ⇒ (f)' doppelt bzw. falsch |
| 207 | ○ | The rotation semigroup (see Part-I, Example 1.2.5) | Verweis vereinheitlichen (S. 190, 197, 202: 'Part-I, Example 3.5.6') | Konsistenz | Auch S. 212 'Part-I, Example 1.2.5'; einer der Verweise ist vermutlich falsch |
| 209 | ◐ | (iv) If ker(s(A) − A') contains a positive element (e.g., if X is compact (cf. Theorem 3.1.6), then | (e.g., if X is compact (cf. Theorem 3.1.6)), then | Satz | Klammer nicht geschlossen |
| 209 | ◐ | Protter & Weinberger [20, Chap.I, Theorem 3] | Protter & Weinberger [20, Chap. I, Theorem 3] | Satz | |
| 210 | ○ | a ϕ ⩾ 0 satisfying A'ϕ = 0 always exists (see Theorem 3.1.6) | … always exists (see Theorem 3.1.6). | Satz | Satzpunkt fehlt |
| 210 | ○ | (i) Pσ(A) ∩ iR is a (additive) subgroup of iR. | (i) Pσ(A) ∩ iR is an (additive) subgroup of iR. | Grammatik | |
| 212 | • | such that Pσ(A) = H). Here (H_d)^ denotes | such that Pσ(A) = H. Here (H_d)^ denotes | Satz | Überzählige Klammer, im PDF bestätigt |
| 212 | ◐ | For details see Greiner [15, p.62]. | For details see Greiner [15, p. 62]. | Satz | |

| | | | | | |
|---|---|---|---|---|---|
| 215 | • | A complete proof of this results requires | A complete proof of this result requires | Grammatik | |
| 216 | ◐ | described explicitly by a semi-flow ϕ (S. 216); semi-flows associated with ordinary differential equations (S. 222); dagegen sonst durchgehend 'semiflow' | semiflow / semiflows (einheitlich) | Konsistenz | auch S. 226 'construct semi-flows associated with (T(t))' |
| 218 | ◐ | it is enough to prove assertion (i), (ii) and (iii) respectively | it is enough to prove assertions (i), (ii) and (iii) respectively | Grammatik | |
| 219 | ◐ | (i) By the definition given in Eq. (3.4.1) on page 216 there exists a $\tau > 0$ such that $c_t(h,\phi) \geqslant$ … | (i) By the definition given in Eq. (3.4.4) on page 217 … | Querverweis | c(h,ϕ) ist in (3.4.4) definiert, nicht in (3.4.1) |
| 220 | • | by Part-I, Proposition 3.6.2 Since every boundary point | by Part-I, Proposition 3.6.2. Since every boundary point | Satz | Punkt fehlt |
| 220 | ◐ | there is a $x_f \in K$ such that | there is an $x_f \in K$ such that | Grammatik | |
| 221 | • | by Lemma 3.4.2 (b) there exists $y \in K\infty$ | by Lemma 3.4.2 (ii) there exists $y \in K\infty$ | Querverweis | Lemma 3.4.2 hat Teile (i)-(iii), kein (b) |
| 221 | • | For details see Sect.18.2 of Dieudonné [11]. | For details see Section 18.2 of Dieudonné [11]. | Satz | Leerzeichen fehlt; sonst wird 'Section' ausgeschrieben (im PDF geprüft) |
| 221 | ◐ | ϕ0 and $\tilde{\phi}0$ satisfy condition Theorem 3.4.4 (iii). | ϕ0 and $\tilde{\phi}0$ satisfy condition (iii) of Theorem 3.4.4. | Grammatik | |
| 222 | • | (iii) We consider Ω as in (b) and assume | (iii) We consider Ω as in (ii) and assume | Querverweis | Teile des Beispiels 3.4.6 sind (i)-(iii) nummeriert |
| 222 | • | by remarking that the ideas of (b) and (c) can be combined | by remarking that the ideas of (ii) and (iii) can be combined | Querverweis | |
| 225 | ◐ | for every $x \in K$ there exists a $r = r_x$ with | for every $x \in K$ there exists an $r = r_x$ with | Grammatik | |
| 227 | • | The dichotomy stated in (Eq.(3.2.19) on page 202) is probably | The dichotomy stated in Eq. (3.2.19) on page 202 is probably | Satz | Leerzeichen fehlt, überflüssige Klammern (im PDF geprüft) |
| | | introduced by | Zitat im Literaturformat, z. B. | | Schaefer (1980) steht |

| | | | | | |
|---|---|---|---|---|---|
| 227 | ○ | Derndinger [8] (and Schaefer (1980) respectively) | 'Schaefer [n]', oder Eintrag ergänzen | Literatur | nicht im Literaturverzeichnis des Kapitels |
| 228 | ◐ | [3] J.-M. Bony, P. Courrège & P. Priouret: Semi-groups de Feller sur une variété à bord compact | Semi-groupes de Feller sur une variété à bord compact | Literatur | französischer Originaltitel |
| 229 | • | [22] J. P. Roth: “Les operateurs elliptiques comme générateurs infinitésimaux de semigroups de Feller”. Séminaire de Théorie du Potential | “Les opérateurs elliptiques comme générateurs infinitésimaux de semi-groupes de Feller”. Séminaire de Théorie du Potentiel | Literatur | Akzent fehlt, 'semigroups' englisch, 'Potential' statt 'Potentiel' |
| 229 | ◐ | [28] H. H. Schaefer: “Some recent results on positive groups and semi-groups. Proceedings of the Symposium in Honour of A.C. Zaanen”. From A to Z. ... Vol. 149. Mathematical Centre 1982 | “Some recent results on positive groups and semi-groups”. From A to Z: Proceedings of a Symposium in Honour of A.C. Zaanen. Ed. by ... Mathematical Centre Tracts 149. Mathematisch Centrum, Amsterdam (1982) pp. 69–79. | Literatur | Tagungsbandtitel steckt im Aufsatztitel; Reihe fehlt |
| 229 | ○ | [18] K. Kuhn: ... Semesterbericht Funktionalanalysis. 1984 pp. 125–142. | Semesterbericht Funktionalanalysis. Universität Tübingen, 1984 pp. 125–142. | Literatur | im Literaturverzeichnis von Kap. 4 (Greiner [13]) mit 'Universität Tübingen' |

## Teil II, Kap. 4 – Asymptotics of Positive Semigroups on $C_0(X)$

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 232 | ◐ | Proof. Eq. (4.1.2) follows directly from Section 3.4.1 and the results from Chapter 2 | vermutlich: Theorem 3.4.1 (oder ein anderer gemeinter Verweis) | Querverweis | Einen Abschnitt 3.4.1 gibt es nicht |
| 233 | ○ | Γ{1} is a homoclinic orbit i.e., | Γ{1} is a homoclinic orbit, i.e., | Satz | |
| 234 | ◐ | (2) Assume, that T(t)f = e^{iαt}f | (2) Assume that T(t)f = e^{iαt}f | Grammatik | kein Komma vor 'that' |
| 234 | ◐ | (T(t)f)(z) → 0 μ-a.e.. Lebesgue’s | (T(t)f)(z) → 0 μ-a.e. Lebesgue’s | Satz | doppelter Punkt |
| 235 | ○ | (see Schaefer [29, II.Theorem 5.9]) | (see Schaefer [29, II. Theorem 5.9]) | Satz | vgl. S. 240 'II. Proposition 9.4' |

| | | | | | |
|---|---|---|---|---|---|
| 236 | ○ | (see Asimow & Ellis [1], p.39) | (see Asimow & Ellis [1], p. 39) | Satz | |
| 240 | • | is a countable additive set function | is a countably additive set function | Grammatik | |
| 240 | ○ | (see e.g., Schaefer [29, II. Proposition 9.4 ]) | (see e.g., Schaefer [29, II. Proposition 9.4]) | Satz | Leerzeichen vor schließender Klammer |
| 242 | • | Then ω_ess(T) < 0 by Eq. (3.3.10) on page 85). | Then ω_ess(T) < 0 by Eq. (3.3.10) on page 85. | Satz | überzählige schließende Klammer |
| 245 | ◐ | the asymptotic behavior as t → ∞ of certain semigroups if only its generator is known | … of certain semigroups if only their generator is known | Grammatik | |
| 251 | • | as developed in Part-I, Section 4.1, Section 4.1 | as developed in Part-I, Section 4.1. | Tippfehler | Verweis doppelt, Satzpunkt fehlt |
| 251 | ◐ | Again we use the assertion that dim F < ∞ | Again we use the assumption that dim F < ∞ | Tippfehler | |
| 251 | ◐ | by the Theorem of Arzela-Ascoli | by the Theorem of Arzelà–Ascoli | Tippfehler | |
| 252 | ○ | Proof. By Formula Eq. (4.3.2) on page 248 | Proof. By Eq. (4.3.2) on page 248 | Satz | ebenso S. 258: 'see Formula Eq. (4.3.3)' |
| 253 | • | then σ(A) = ∅ by Proposition 3.4. | then σ(A) = ∅ by Proposition 4.3.4. | Querverweis | |
| 253 | ◐ | by Part-III, Theorem 3.1.1. (i), we obtain | by Part-III, Theorem 3.1.1 (i), we obtain | Satz | |
| 256 | ○ | since, even a scalar valued delay differential equation may be stable | since even a scalar-valued delay differential equation may be stable | Satz | |
| 257 | • | Then (3.11) takes the abstract form | Then Eq. (4.3.10) takes the abstract form | Querverweis | Alter Gleichungsverweis '(3.11)'; gemeint ist die Gleichung aus Example 4.3.11 (bitte Nummer prüfen) |
| 257 | ○ | It is well-known that B̃ generates | It is well known that B̃ generates | Konsistenz | sonst 'well known' (S. 247, 259) |
| 259 | • | showed the translation property (T)for the solution semigroup | showed the translation property (T) for the solution semigroup | Satz | im PDF geprüft |

| | | | | | |
|---|---|---|---|---|---|
| 259 | • | (see Part-III, Section 3.4 can be used to derive | (see Part-III, Section 3.4) can be used to derive | Satz | schließende Klammer fehlt |
| 259 | ◐ | Section 2: ... Theorem 4.1.1 seems to be folklore. | Theorem 4.2.1 seems to be folklore. | Querverweis | In den Notizen zu Abschnitt 2 ist wohl Theorem 4.2.1 (eventuell kompakte Halbgruppen) gemeint; 4.1.1 wird schon in Section 1 genannt |
| 259 | ◐ | This relation more detailed than in Example 4.2.6, can be found | This relation, in more detail than in Example 4.2.6, can be found | Grammatik | |
| 259 | ○ | in Chapter 2 of Casteren [3] | in Chapter 2 of van Casteren [3] | Literatur | Literatureintrag lautet 'J. van Casteren'; ebenso S. 241 |

## Teil III, Kap. 1 – Basic Results on Banach Lattices and Positive Operators

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 265 | • | Rainer Nagel und Ulf Schlotterbeck | Rainer Nagel and Ulf Schlotterbeck | Tippfehler | Deutsches 'und' in der Autorenzeile des Kapitels |
| 267 | • | contains x1, ..., xn (see Section. 1.4). | contains x1, ..., xn (see Section 1.4). | Satz | |
| 268 | ◐ | (more generally, if E is archimedean ordered, then B^dd is the band generated by B | (more generally, if E is archimedean ordered), then B^dd is the band generated by B | Satz | schließende Klammer fehlt |
| 270 | ○ | order-bounded / order bounded; order-continuous / order continuous; order-complete / order complete (gemischt in Kap. III.1, z. B. S. 270 'order-bounded' und 'order bounded functional', S. 272 'order continuous') | einheitliche Schreibung (entweder mit oder ohne Bindestrich) | Konsistenz | |
| 271 | ○ | called an (AM)-space with unit | called an AM-space with unit | Konsistenz | sonst immer 'AM-space' |
| 275 | ◐ | T is called real if T2 is = 0 | T is called real if T2 = 0 | Grammatik | |
| 276 | ◐ | the operator norm and the composition product is Z(E) isomorphic to a Banach lattice algebra | the operator norm and the composition product, Z(E) is isomorphic to a Banach lattice algebra | Grammatik | |

## Teil III, Kap. 2 – Characterization of Positive Semigroups on Banach Lattices

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 280 | ◐ | (this follows from the definition, see Eq. (2.2.3) on page 57. | (this follows from the definition, see Eq. (2.2.3) on page 57). | Satz | schließende Klammer fehlt |
| 284 | • | of all infinitely differentiabele functions on Ω | of all infinitely differentiable functions on Ω | Tippfehler | |
| 284 | ◐ | Consider the function b ∈ C(∂Ω), given by, b(x) = $\bar{f}$(x) | Consider the function b ∈ C(∂Ω) given by b(x) = $\bar{f}$(x) | Satz | Überflüssiges Komma nach „given by“. |
| 285 | ◐ | We know from (iv) that C_c^∞(Ω) ⊆ (Id −A)D(A) | We know from (v) that C_c^∞(Ω) ⊆ (Id −A)D(A) | Querverweis | Die Aussage wird im Beweis von (v) gezeigt, nicht in (iv). |

| | | | | | |
|---|---|---|---|---|---|
| 289 | ○ | Gateaux-derivative … right-sided Gateaux-differentiable | einheitlich ohne Bindestrich wie auf S. 315/317 („Gateaux differentiable“), ggf. „Gâteaux“ | Konsistenz | S. 289 schreibt „Gateaux-differentiable“, S. 315 und 317 „Gateaux differentiable“. |
| 292 | • | It has been proved by Miyajima & Okazawa [23]] that | It has been proved by Miyajima & Okazawa [23] that | Satz | Doppelte schließende Klammer; im PDF bestätigt. |
| 294 | • | In the proof of the "only if" - part of Proposition 2.3.6 | In the proof of the “only if” part of Proposition 2.3.6 | Satz | Gerade Anführungszeichen statt typografischer; dazu ein frei stehender Bindestrich. |
| 301 | ◐ | in a manner analogous to the characterization of positive semigroups in Section 3.1, however, no positive subeigenvectors are needed here. | in a manner analogous to the characterization of positive semigroups in Section 2.3; however, no positive subeigenvectors are needed here. | Querverweis | Die Charakterisierung positiver Halbgruppen steht in Abschnitt 2.3 (Abschnitt 3.1 ist „The Spectral Bound“). Außerdem fehlt ein Semikolon (Komma-Spleiß vor „however“). |
| 301 | ○ | in order to decide whether a semigroup is stable see Part-I, Chapter 4 | in order to decide whether a semigroup is stable, see Part-I, Chapter 4 | Satz | |
| 303 | • | which finally implies that B is closable Now let λ > λ0 | which finally implies that B is closable. | Satz | Am Satzende fehlt der Punkt; im PDF bestätigt. |
| 304 | ◐ | Let M ∈ Z(E) (the center of E (see Section 1.9). | Let M ∈ Z(E) (the center of E, see Section 1.9). | Satz | Die Klammern sind nicht ausgeglichen. |
| 308 | ◐ | how the results of this section look like for C0(X) | what the results of this section look like for C0(X) | Grammatik | |
| 313 | ◐ | a linear operator S on E is called lattice homomorphism if | a linear operator S on E is called a lattice homomorphism if | Grammatik | |
| 314 | ○ | Note, that an operator S is a lattice homomorphism, if and only if S is positive | Note that an operator S is a lattice homomorphism if and only if S is positive | Satz | Kommas nach deutscher Zeichensetzung. |
| 315 | • | is right-sided Gateaux differerentiable and | is right-sided Gateaux differentiable and | Tippfehler | Im PDF bestätigt. |
| 315 | ○ | For that we need, that the modulus function is differentiable. If E = L^p(X,Σ,μ) | For that we need that the modulus function is differentiable. If E = | Grammatik | Komma nach deutscher |

| | | | | | |
|---|---|---|---|---|---|
| | | (1 ⩽ p < ∞) this had been proved in Example 2.2.3. | L^p(X,Σ,μ) (1 ⩽ p < ∞) this has been proved in Example 2.2.3. | | Zeichensetzung; Zeitform. |
| 316 | ◐ | Then by Part-II, Chapter 2, Lemma 2.4 | Then by Part-II, Lemma 2.2.4 | Querverweis | Andere Zitierweise als sonst. Dieselbe Aussage wird auf S. 289 als „Part-II, Lemma 2.2.4“ zitiert. |
| 317 | ○ | (see Section 1.8) ). | (see Section 1.8)). | Satz | Leerzeichen vor der Klammer. |
| 320 | ○ | Here we call a semigroup (T(t))t⩾0 multiplication semigroup if | Here we call a semigroup (T(t))t⩾0 a multiplication semigroup if | Grammatik | |
| 321 | ○ | (see Section 1.9) , it is clear that (a) implies (c). … Hence A ∈ Z(E) The last assertion | (see Section 1.9), it is clear that (a) implies (c). … Hence A ∈ Z(E). | Satz | Leerzeichen vor dem Komma; am Satzende fehlt der Punkt. |
| 322 | ○ | (this follows from Eq. (2.1.6) on page 52) | (this follows from Part-I, Eq. (…) on page 52) | Querverweis | Mehrdeutig: Auch dieses Kapitel hat eine Eq. (2.1.6), nämlich die Gauß-Kern-Formel auf S. 284. Gemeint ist offenbar eine Formel aus einem anderen Teil; bitte den Teil angeben oder den Verweis prüfen. |
| 323 | ◐ | Theorem 2.1.8 has been obtained independently by Arendt (1984a) and van Casteren (1984). | Zitierweise mit Nummern wie sonst (z. B. Arendt [3]); van Casteren fehlt im Literaturverzeichnis | Literatur | Autor-Jahr-Zitate passen nicht zum nummerierten Verzeichnis, und „van Casteren (1984)“ steht nicht in der Liste. |
| 323 | ○ | characterizes the positivity of the semigroup (cf., Nagel & Uhlig [24]) | (cf. Nagel & Uhlig [24]) | Satz | |
| 324 | ◐ | [3] W. Arendt: “Generators of positive semigroups”. Infinite-dimensional Systems, Retahof 1983. Ed. by F. Kappel & W. Schappacher. Vol. 1076. Springer 1984 pp. 1–15. | … Infinite-dimensional Systems, Retzhof 1983. Ed. by F. Kappel & W. Schappacher. Lecture Notes in Math. 1076. Springer 1984, pp. 1–15. | Literatur | Ortsname falsch geschrieben (Retzhof); Reihentitel fehlt vor „Vol. 1076“; vor „pp.“ fehlt ein Komma. |
| 324 | ○ | [10] … Ed. by R. Nagel, U. Schlotterbeck & M. Wolff. | Ed. by R. Nagel, U. Schlotterbeck & M. P. H. Wolff. | Literatur | [17] und [22] nennen die Herausgeber desselben Bandes als „M. P. H. Wolff“. |

| | | | | | |
|---|---|---|---|---|---|
| 325 | • | [27] J. Peetre: Une charactérisation abstraite des opérateurs différentiels. | [27] J. Peetre: Une caractérisation abstraite des opérateurs différentiels. | Literatur | Französische Schreibweise: „caractérisation“. |
| 325 | ○ | [21] W. A. J. Luxemburg: Some Aspects of the Theory of Riesz Spaces. Vol. 4. (1979). | … Vol. 4. Univ. of Arkansas Lecture Notes in Math. (1979). | Literatur | Reihe bzw. Verlag fehlen, der Eintrag ist unvollständig. Bitte die genaue Angabe prüfen. |
| 325 | ○ | [23] S. Miyajima & N. Okazawa: “Generators of positive C0-semigroups on Banach lattices”. 1984. | Bibliografische Angabe ergänzen (Zeitschrift, Band, Seiten bzw. „Preprint“) | Literatur | Unvollständiger Eintrag. |

## Teil III, Kap. 3 – Spectral Theory of Positive Semigroups on Banach Lattices

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 338 | ◐ | Now we apply Proposition 3.2.7(b) and obtain | Now we apply Proposition 3.2.7 and obtain | Querverweis | Proposition 3.2.7 hat keinen Teil (b). |
| 338 | ○ | Then h2 ≔ h1 + I ≠ 0 (by Eq. (3.2.19). Moreover, λR2(λ + iβ)h2 = h2 (by Eq. (3.2.16) on the preceding page and | (by Eq. (3.2.19)). … (by Eq. (3.2.16) on the preceding page) and | Satz | Die Klammern sind nicht ausgeglichen. |
| 339 | ○ | In case that σb(A) is cyclic one has the alternative (cf. Eq. (3.2.19) on page 202) | (cf. Part-II, Eq. (3.2.19) on page 202) | Querverweis | Mehrdeutig: Dieses Kapitel hat auf S. 338 eine eigene Eq. (3.2.19). Gemeint ist wohl die Formel aus Part II. |
| 339 | ○ | then σb(A) is cyclic . | then σb(A) is cyclic. | Satz | |
| 341 | • | then T(t)f is a quasi-interior pointof E+ whenever | then T(t)f is a quasi-interior point of E+ whenever | Satz | Wörter zusammengezogen; im PDF bestätigt. |
| 341 | ◐ | examples of semigroups whose resolvent do not grow slowly | examples of semigroups whose resolvents do not grow slowly | Grammatik | |
| 341 | ○ | (see p.65 in Greiner [13]) | (see p. 65 in Greiner [13]) | Satz | |
| 342 | ○ | Then strong continuity and T(0) = Id ≠ 0 implies that | Then strong continuity and T(0) = Id ≠ 0 imply that | Grammatik | |

| | | | | | |
|---|---|---|---|---|---|
| 345 | ◐ | This extends Theorem 2.3.5 to the setting of Banach lattices. The first assertion of the latter proposition | This extends Part-II, Proposition 3.3.5 to the setting of Banach lattices. The first assertion of the latter proposition | Querverweis | „latter proposition“ und der Beweis („see Part-II, Proposition 3.3.5“) zeigen, dass Part-II, Proposition 3.3.5 gemeint ist. Theorem 2.3.5 gibt es so nicht. |
| 345 | ○ | Suppose A to be the generator of an irreducible, positive semigroup | Suppose A is the generator of an irreducible, positive semigroup | Grammatik | |
| 345 | ○ | By assumption and by assertion (i) there exists $\phi \gg 0$ | By assumption and by assertion (ii) there exists $\phi \gg 0$ | Querverweis | Dass das Funktional strikt positiv ist, folgt aus (ii) (Eigenvektoren von A'), nicht aus (i). |
| 347 | ◐ | (i) $P\sigma(A) \cap iR$ is a (additive) subgroup of iR. | (i) $P\sigma(A) \cap iR$ is an (additive) subgroup of iR. | Grammatik | |
| 347 | ○ | as presented in Sect.V.6 of Schaefer [32] (see Proposition 6.1, Theorem 6.5 Corollary and Theorem 6.5 l.c.). ... The assertions (ii),(iv) and (v) | as presented in Sect. V.6 of Schaefer [32] (see Proposition 6.1, Theorem 6.5, Corollary and Theorem 6.5 l.c.). ... The assertions (ii), (iv) and (v) | Satz | Leerzeichen fehlen; die Aufzählung ist unklar („Theorem 6.5“ steht doppelt), bitte prüfen. |
| 348 | • | which is finer than the Frechet filter | which is finer than the Fréchet filter | Tippfehler | Akzent fehlt. Dasselbe auf S. 350 (PDF 368). |
| 348 | ○ | Hereby, $j \circ i$ is the canonical inclusion of C(Γ) in L1(Γ). | Here, $j \circ i$ is the canonical inclusion of C(Γ) in L1(Γ). | Grammatik | „Hereby“ ist ein Germanismus (von „hierbei“). |
| 348 | ○ | We define $j : C(\Gamma) \to E$ by $j(f) \coloneqq f \circ h$ ... we define $\tilde{j}$ to be the canonical mapping from E | We define $i : C(\Gamma) \to E$ by $i(f) \coloneqq f \circ h$ ... (entsprechend der Bezeichnung i, j in der Aussage) | Konsistenz | In der Aussage heißen die Abbildungen i und j, im Beweis j und $\tilde{j}$. Das betrifft die Bezeichnungen; bitte prüfen. |
| 350 | • | ultrafilter U on N which is finer than the Frechet filter | ultrafilter U on N which is finer than the Fréchet filter | Tippfehler | |
| 352 | • | by calculating the resolvent explicitely | by calculating the resolvent explicitly | Tippfehler | |
| 352 | ◐ | The meaning of r_ess(T(t)) and ω_ess(T) and its properties are explained | The meaning of r_ess(T(t)) and ω_ess(T) and their properties are explained | Grammatik | |
| | | | | | |

| | | | | | |
|---|---|---|---|---|---|
| 352 | ◐ | Proof. There exist ε > 0 such that | Proof. There exists ε > 0 such that | Grammatik | |
| 352 | ○ | an irreducible, eventually norm continuous semigroup | an irreducible, eventually norm-continuous semigroup | Konsistenz | S. 340 (Corollary 3.2.13) schreibt „eventually norm-continuous“. |
| 354 | • | appearing in the the representation Eq. (3.3.21) | appearing in the representation Eq. (3.3.21) | Tippfehler | Doppeltes 'the' |
| 354 | ◐ | For details we refer to Greiner [15] or Voigt [36] where the linear transport equation in higher dimensional spaces is discussed. | … Greiner [15] or Voigt [37] … | Literatur | [36] ist 'Interpolation for positive C0-semigroups'; zur Transportgleichung passt [37] (so auch in den Notes S. 366 zitiert) |
| 355 | ◐ | then one of the following assertions are true | then one of the following assertions is true | Grammatik | |
| 357 | • | the Césaro means | the Cesàro means | Tippfehler | Akzent falsch; S. 382 korrekt 'Cesàro' |
| 359 | ○ | (cf. Part-I, Definition 3.3.1) (One simply performs a spectral decomposition of a single operator $T(t)$). | (cf. Part-I, Definition 3.3.1). (One simply performs a spectral decomposition of a single operator $T(t)$.) | Satz | Satzzeichen vor/in der Klammer |
| 363 | ◐ | are well-defined as bounded linear operator on $Lp(\mu)$ | are well-defined as bounded linear operators on $Lp(\mu)$ | Grammatik | |
| 364 | ○ | then $f \circ \varphi t = g \circ \varphi t$ $\mu$-a.e.. Moreover | then $f \circ \varphi t = g \circ \varphi t$ $\mu$-a.e. Moreover | Satz | Doppelter Punkt; ebenso S. 384 ('$\mu$-a.e..') und S. 385 ('$\mu$ − a.e..') |
| 364 | ○ | Note that for continuous flows condition Eq. (3.4.12) | Note that for continuous flows condition (3.4.12) | Konsistenz | Doppelte Bezeichnung (condition + Eq.); gleiches Muster: 'Formula Eq. (4.3.2)' S. 393, 'formula Eq. (4.3.3)' S. 394, 'equation Eq. (4.3.2)' S. 395, 'Formula Eq. (4.3.7)' S. 396, 'formula Eq. (4.3.9)' S. 398, 'formula Eq. (4.3.3)'? – vermutlich systematisch durch \eqref-Konvertierung |
| 366 | • | or “quasi-strictly positive” in Karlin [21]). Sawashima [30] | or “quasi-strictly positive” in Karlin [21]. Sawashima [30] | Satz | Schließende Klammer ohne öffnende |
| | | [17] G. Greiner, J. Voigt & M. Wolff: On the | | | |

| | | | | | |
|---|---|---|---|---|---|
| 367 | ◐ | spectral bound of the generator of semigroups of positive operators. J. Operator Theory (1981) 245–256. | J. Operator Theory 5 (1981) 245–256. | Literatur | Bandnummer fehlt |
| 368 | • | [34] J. Voigt: On the abszissa of convergence for the Laplace transform | On the abscissa of convergence for the Laplace transform | Literatur | Deutsche Schreibweise im englischen Titel |
| 368 | ○ | [37] J. Voigt: Spectral properties of the neutron transport equations. | Spectral properties of the neutron transport equation. | Literatur | Originaltitel im Singular – bitte prüfen |

## Teil III, Kap. 4 – Asymptotics of Positive Semigroups on Banach Lattices

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 369 | ○ | positive semigroups on AM- , AL-, and Hilbert spaces | positive semigroups on AM-, AL-, and Hilbert spaces | Satz | Leerzeichen vor Komma |
| 370 | • | $s(A) \in \sigma(A)$ (see Corollary 3.1.4 ). | (see Corollary 3.1.4). | Satz | Leerzeichen vor Klammer (im PDF sichtbar); ebenso S. 395 '(see Proposition 2.4.1 )' und '(see Corollary 3.2.12 )', S. 401 '[3, p.257ff] )' |
| 370 | ◐ | (cf. Part-II, Remark 4.1.5.) | (cf. Part-II, Remark 4.1.5) | Satz | Überflüssiger Punkt |
| 372 | ◐ | combining Part-III, Theorems 3.1.2 with Part-I, Theorem 4.1.4 | combining Theorem 3.1.2 with Part-I, Theorem 4.1.4 | Grammatik | Plural 'Theorems' bei nur einem Satz; zudem sonst innerhalb von Part III ohne 'Part-III' zitiert |
| 376 | ○ | In case where $A$ generates a positive semigroup | In case $A$ generates a positive semigroup | Grammatik | |
| 376 | ○ | we discuss the long term behavior of the solutions | we discuss the long-term behavior of the solutions | Konsistenz | S. 369 'long-term behavior'; S. 398 wieder 'long term behavior' |
| 379 | ◐ | Using the general results presented in Section 4.2 and the spectral theoretical result of Chapter 3 | Using the general results presented in Part-II, Section 4.2 and the spectral theoretical results of Chapter 3 | Querverweis | Man befindet sich selbst in Section 4.2; gemeint ist offenbar Part-II (Beweis zitiert Part-II, Theorem 4.2.10 und S. 244) |
| 379 | ○ | (see Part-I, Example 3.1.3 & 3.1.4) | (see Part-I, Examples 3.1.3 and 3.1.4) | Grammatik | |

| | | | | | |
|---|---|---|---|---|---|
| 384 | ◐ | We briefly describe this situation and remind that in this context | We briefly describe this situation and recall that in this context | Grammatik | 'remind' verlangt ein Objekt |
| 385 | • | a Markov transition function satisfying all the assumptions of 4.2.7 induces | a Markov transition function satisfying all the assumptions of Example 4.2.8 induces | Querverweis | 4.2.7 ist ein Korollar; die Annahmen (i)–(v) stehen in Example 4.2.8 |
| 387 | ○ | (see Krengel [25, Chapter 2, Theorem 4.4 and 4.5]) | (see Krengel [25, Chapter 2, Theorems 4.4 and 4.5]) | Grammatik | |
| 390 | ◐ | the semigroup generated by $C$ fulfills the assertions of Theorem 4.2.14 | the semigroup generated by $C$ fulfills the assumptions of Theorem 4.2.14 | Tippfehler | Gemeint sind die Voraussetzungen |
| 393 | ◐ | (compare Part-I, Example 1.2.4. (ii)) | (compare Part-I, Example 1.2.4 (ii)) | Satz | Überflüssiger Punkt |
| 395 | ○ | The semigroup generated by $A0$ is positive (use (RE) on page 392) | The semigroup generated by $A0$ is positive (use Eq. (4.3.1) on page 393) | Querverweis | Positivität von (T0(t)) folgt aus der Darstellung (4.3.1), nicht aus (RE) – bitte prüfen |
| 396 | ○ | According to the proof of 4.3.5 (ii) this is equivalent | According to the proof of Lemma 4.3.5 (ii) this is equivalent | Querverweis | |
| 399 | ◐ | (due to the irreducibility of $K(s)$) , $x'\mu$ is a strictly positive functional | (due to the irreducibility of $K(s)$), $x'\mu$ is a strictly positive functional | Satz | Komma rutscht an den Zeilenanfang (im PDF sichtbar) |
| 399 | ◐ | based on unequal division of cells, (see Arino & Kimmel [1]). | based on unequal division of cells (see Arino & Kimmel [1]). | Satz | Überflüssiges Komma |
| 400 | ◐ | ($\Phi\lambda$)2 is compact [Schaefer [36, Theorem II.9.9]] which | ($\Phi\lambda$)2 is compact (Schaefer [36, Theorem II.9.9]), which | Satz | Verschachtelte eckige Klammern |
| 401 | ○ | The birth-rate is denoted by $\beta$. ... dependence of birth rate on individuals | The birth rate is denoted by $\beta$. ... dependence of birth rate on individuals | Konsistenz | Im selben Absatz 'birth-rate' und 'birth rate'; S. 402 'birth rate' |
| 402 | • | (see Schaefer [36, Section II.5] and thus compactness of $B2$ | (see Schaefer [36, Section II.5]) and thus compactness of $B2$ | Satz | Schließende runde Klammer fehlt |
| 402 | • | As in Example 4.3.13 the first equation in (4.3.14) corresponds to | As in Example 4.3.12 the first equation in (4.3.14) corresponds | Querverweis | Selbstverweis; gemeint ist das vorige Beispiel |

| | | | | | |
|---|---|---|---|---|---|
| | | the aging process. | to the aging process. | | |
| 402 | ◐ | by the Dunford-Pettis-Property of $L1(\Omega)$ (see Schaefer [36, Chapter II, Theorem.9.9]) | by the Dunford–Pettis property of $L1(\Omega)$ (see Schaefer [36, Theorem II.9.9]) | Satz | 'Dunford-Pettis-Property' inkonsistent zu 'Dunford-Pettis property' (S. 398, 400); 'Theorem.9.9' mit Punkt |
| 403 | ◐ | Example 4.2.3 is discussed in more detail in Webb [42] and Greiner [14]. | … in Webb [42] and Greiner [13]. | Literatur | [14] ist 'An irreducibility criterion for the linear transport equation'; zur Altersstruktur passt [13] (so auch S. 381 zitiert). Ebenso 'population equations by Greiner [14]' weiter unten auf S. 403 |
| 403 | ◐ | is given in Chapter 2 of Casteren [4] | is given in Chapter 2 of van Casteren [4] | Literatur | Name laut Literaturliste 'J. van Casteren' |
| 403 | ○ | (see Part-II, Chapter 4, Sec.3) | (see Part-II, Section 4.3) | Konsistenz | Sonst stets 'Section x.y' |
| 404 | ◐ | [15] G. Greiner & R. Nagel: La loi “zero ou deux” et ses conséquences … J. Math. Pures Appl. 9 (1982) 261–273. | La loi « zéro ou deux » et ses conséquences … J. Math. Pures Appl. (9) 61 (1982) 261–273. | Literatur | Akzent 'zéro' fehlt; '9' ist die Serie, Band 61 fehlt |
| 405 | ○ | North Holland 1986. / North Holland 1984. / Riesz Spaces II. North Holland (1983). | North-Holland (1986) / North-Holland (1984) / North-Holland (1983) | Literatur | Verlagsname 'North-Holland'; Jahresangabe teils mit, teils ohne Klammern (auch 'Springer 1984 pp.', 'Marcel Dekker 1985 pp.') |

## Teil IV, Kap. 1 – Basic Results on Semigroups and Operator Algebras

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 409 | ◐ | if $T \otimes \mathrm{Id}n$ is positive from $M \otimes Mn$ in $M \otimes Mn$ | if $T \otimes \mathrm{Id}n$ is positive from $M \otimes Mn$ into $M \otimes Mn$ | Grammatik | |
| 409 | ○ | satisfies the so called Schwarz-inequality | satisfies the so-called Schwarz inequality | Konsistenz | Auf S. 410 'Schwarz inequality' ohne Bindestrich |
| 410 | ◐ | on a commutative C∗-algebra is a Schwarz map. (Takesaki [7, Chapter IV]). | on a commutative C∗-algebra is a Schwarz map (Takesaki [7, Chapter IV]). | Satz | Punkt vor der Quellenklammer |
| 410 | ◐ | If $(S(t)x)(S(t)x)* \leqslant T(t)(xx*)$ for all $x \in M$ and $t \in$ R+. Then | If $(S(t)x)(S(t)x)* \leqslant T(t)(xx*)$ for all $x \in M$ and $t \in$ R+, then | Grammatik | If-Satz durch Punkt abgetrennt (Theorem 1.2.1) |
| 410 | ○ | we recommend to compare the results of this section | we recommend comparing the results of this section | Grammatik | |
| 412 | ◐ | As in Part-III, Section 3.3 the following notion will be an important tool | As in Part-III, Section 3.2 the following notion will be an important tool | Querverweis | Pseudo-Resolventen werden in Part III, Abschnitt 3.2 behandelt (vgl. 'Part-III, Example 3.2.5' wenige Zeilen später) |
| 412 | ○ | T is a semigroup of Schwarz-type, if and only if $\mu R(\mu, A)$ is a Schwarz-operator | T is a semigroup of Schwarz type if and only if $\mu R(\mu, A)$ is a Schwarz operator | Konsistenz | Sonst 'Schwarz type' ohne Bindestrich; Komma vor 'if and only if' überflüssig |
| 413 | • | R$p$ has both the Schwarz type property and is identity preservation. | R$p$ has the Schwarz type property and is identity preserving. | Grammatik | |
| 414 | • | the following equalities holds | the following equalities hold | Grammatik | |
| 414 | ◐ | [1] N. Bourbaki: Éléments des Mathématiques, Intégration, Chapitre 5: Intégration des Mesures. | N. Bourbaki: Éléments de mathématique. Intégration, Chapitre 5: Intégration des mesures. | Literatur | Originaltitel 'Éléments de mathématique' |
| 415 | ○ | Completely positive linear maps, non Hamiltonian evolution | Completely positive linear maps, non-Hamiltonian evolution | Literatur | |

## Teil IV, Kap. 2 – Characterization of Positive Semigroups on W*-Algebras

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 417 | ○ | which derivations) commuting with certain automorphism groups are automatically generators of strongly continuous positive semigroups. | … generators of strongly continuous positive semigroups? | Satz | Formulierung als Frage ('Which maps …'), endet mit Punkt |
| 418 | • | Using the Schwarz, inequality we obtain | Using the Schwarz inequality, we obtain | Satz | Komma an falscher Stelle (im PDF sichtbar) |
| 418 | ◐ | (compare Elliot [1], Lemma 2.). | (compare Elliott [1, Lemma 2]). | Literatur | Name korrekt 'Elliott' (auch in der Literaturliste S. 419 'G. Elliot'); überflüssiger Punkt; Zitierformat sonst '[n, …]' |
| 418 | ○ | Take (i)' $M$ be an ultrapower of $M$, | Take: (i)' let $M$ be an ultrapower of $M$, | Grammatik | (ii)'–(iv)' beginnen mit 'let', (i)' nicht |

## Teil IV, Kap. 3 – Spectral Theory on W*-Algebras and their Preduals

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 422 | • | subinvariant for a positive map $T \in$ L($M$) (resp.. positive semigroup T) | (resp. positive semigroup T) | Satz | Doppelter Punkt |
| 422 | ○ | Kolumnentitel: Part IV 3 Spectral Theory on W$_*$-algebras | Part IV 3 Spectral Theory on W$_*$-Algebras | Konsistenz | Kapiteltitel und andere Kolumnentitel schreiben 'W$_*$-Algebras' groß |
| 423 | ◐ | Hence Fix($T$) is a self adjoint subspace of $M$ | selfadjoint (bzw. einheitlich self-adjoint) | Konsistenz | Uneinheitlich: 'self adjoint' (S. 423, 425, 457) neben 'selfadjoint' (S. 425, 426) |
| 425 | • | by the uniqueness theorem (*) on p. 422 for the absolut value | for the absolute value | Tippfehler | |
| 426 | • | we obtain that $\varphi$+ and $\varphi-$ in Fix(R). | we obtain that $\varphi$+ and $\varphi-$ are in Fix(R). | Grammatik | Verb fehlt |
| 428 | ◐ | This is no longer true in the non commutative case. | noncommutative (einheitlich) | Konsistenz | 'non commutative' (S. 428, 429, 454) neben 'noncommutative' (S. 438, 489, 490) |

| | | | | | |
|---|---|---|---|---|---|
| 429 | ◐ | (i) The fixed space of T is one dimensional ... (iii) The fixed space of the adjoint weak*-semigroup T' is one-dimensional. | einheitlich 'one-dimensional' | Konsistenz | Im selben Theorem uneinheitlich; 'one dimensional' auch S. 430, 452, 455 |
| 431 | ◐ | (see, Jacobs [11, Satz 6.1., p.77]) | (see Jacobs [11, Satz 6.1, p. 77]) | Satz | überflüssiges Komma nach 'see', Leerzeichen in 'p. 77' fehlt |
| 432 | • | 3.2 Spectral Properties of Uniformly Ergodic Semigoups | 3.2 Spectral Properties of Uniformly Ergodic Semigroups | Tippfehler | Abschnittstitel; der Fehler steht auch in den Kolumnentiteln S. 433, 435, 437, 439 (und vermutlich im Inhaltsverzeichnis) |
| 432 | • | the discret group $P\sigma(A) \cap$ iR | the discrete group | Tippfehler | |
| 432 | ◐ | fulfils the assertions of Theorem 1.10 | fulfils the assertions of Theorem 3.1.10 | Querverweis | Gemeint ist sicher Theorem 3.1.10 (Altnummerierung) |
| 433 | ◐ | we obtain from Lemma 1.1.b that for all $x \in N$ | we obtain from Lemma 3.1.1 (ii) that | Querverweis | Altnummerierung 'Lemma 1.1.b'; inhaltlich passt Lemma 3.1.1 (ii) |
| 434 | • | and the assertion follows from Lemma 3.1.6). | and the assertion follows from Lemma 3.1.6. | Satz | überzählige schließende Klammer |
| 437 | • | and the assocciated ergodic projection | and the associated ergodic projection | Tippfehler | |
| 437 | ◐ | one part of the corollary is proved | one part of the proposition is proved | Querverweis | Die Aussage ist als Proposition 3.2.3 nummeriert |
| 438 | • | Our main result Theorem 3.1.10 on extends Part-II, Theorem 3.3.6 | Our main result Theorem 3.1.10 extends Part-II, Theorem 3.3.6 | Tippfehler | überflüssiges 'on' |
| 438 | ○ | for all $\alpha \in$ R (Proposition 3.2.1 ). | (Proposition 3.2.1). | Satz | Leerzeichen vor der Klammer |
| 439 | • | G. Greiner, J. Voigt & M. Wolff: On the spectral bound ... J. Operator Theory (1981) 245–256. | J. Operator Theory 5 (1981) 245–256. | Literatur | Bandnummer fehlt (in den Updated Notes Part IV korrekt mit 5); gilt ebenso für Ref. [8] auf S. 463 |

## Teil IV, Kap. 4 – Asymptotics on C- *and W*-Algebras

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 442 | • | characterizes uniform exponentially stability of the semigroup | characterizes uniform exponential stability | Grammatik | |
| 442 | ◐ | From this the assumption follows by Part-I, Theorem 4.1.11. | From this the assertion follows | Tippfehler | |
| 443 | • | Then the following assertion are equivalent. | Then the following assertions are equivalent. | Grammatik | |
| 443 | ◐ | Proof. Since $s(A) = \omega 0$ by Theorem 4.1.3 | by Theorem 4.1.1 | Querverweis | Theorem 4.1.7 betrifft C*-Algebren mit Eins, dafür gilt Theorem 4.1.1; 4.1.3 behandelt Prädualräume |
| 443 | ○ | positive semigroups on C-*algebras with unit. Theorem 4.1.7 Let $\mathcal{M}$ be a C*-algebra | $C_*$-algebras (gleicher Stern wie sonst) | Konsistenz | Hier ein normales Sternchen statt des sonst verwendeten hochgestellten $*$; bitte im PDF prüfen |
| 443 | ○ | but $(T(t))t\geqslant 0$ is strongly stable | is uniformly stable (Terminologie aus Definition 4.1.6 (ii)) | Konsistenz | In Definition 4.1.6 heißt die starke Konvergenz gegen 0 'uniformly stable'; 'strongly stable' kommt dort nicht vor |
| 445 | • | whereas (c) and (d)), resp. (e) and (f) are equivalent by the Remark 4.2.1 (iii). | whereas (c) and (d), resp. (e) and (f), are equivalent by Remark 4.2.1 (iii). | Satz | überzählige Klammer, überflüssiger Artikel 'the' |
| 446 | • | (see Propositions 4.3.7, 4.3.8 &. 4.3.9) | (see Propositions 4.3.7, 4.3.8 & 4.3.9) | Satz | Punkt nach '&' streichen; 4.3.8 ist außerdem ein Theorem und 4.3.9 ein Corollary |
| 446 | • | Therefore the following compactness criterium is quite useful. | compactness criterion | Tippfehler | |
| 446 | • | the orbit $\{T(t)\varphi : t \in \mathrm{R}+\}$ is relatively weak compact | is relatively weakly compact | Grammatik | |
| 447 | ◐ | which is of Schwarz type. It follows as in (b) that | It follows as in (ii) that | Querverweis | Remark 4.3.2 hat die Teile (i)–(iii), ein Teil (b) existiert nicht |

| | | | | | |
|---|---|---|---|---|---|
| 448 | • | (see Krengel [14, Chap.2, Theorem1.4]) | (see Krengel [14, Chap. 2, Theorem 1.4]) | Satz | Leerzeichen fehlen; auf S. 461 fehlt ebenso das Leerzeichen in 'Chap.2' |
| 449 | • | By Proposition 3.1 the semigroup T is strongly ergodic. | By Proposition 4.3.1 the semigroup T is strongly ergodic. | Querverweis | |
| 449 | ◐ | (Leeuw & Glicksberg [15], Junghenn [13] & Krengel [14, § 2.4]) | (de Leeuw & Glicksberg [15], ...) | Konsistenz | Im Literaturverzeichnis steht 'K. de Leeuw' |
| 450 | ◐ | which is not empty by assumptions. | which is not empty by assumption. | Grammatik | |
| 450 | ○ | Halmos-von Neumann Theorem in commutative ergodic theory | Halmos–von Neumann theorem (einheitlich mit S. 455) | Konsistenz | S. 450 'Theorem', S. 455 'theorem' |
| 451 | • | If $u1$ and $u2$ in $U$, then | If $u1$ and $u2$ are in $U$, then | Grammatik | |
| 451 | • | the fact that $T0(t)$' is a *-automorphismus on $N$. | is a *-automorphism on $N$. | Tippfehler | deutsches Wort |
| 452 | • | as group of *-automorphism such that the fixed space | as a group of *-automorphisms | Grammatik | |
| 452 | • | We apply now these results to weak*-semigroup on B($H$). | We now apply these results to weak*-semigroups on B($H$). | Grammatik | |
| 452 | • | and $\varphi$ is a faithful T-invariant normal states. | a faithful T-invariant normal state. | Grammatik | |
| 452 | • | irreducible (equvalently, if the fixed space | (equivalently, if | Tippfehler | |
| 453 | ◐ | We show next, that for irreducible W$*$-dynamical systems | We show next that for irreducible | Satz | Komma nach deutscher Regel, im Englischen falsch |
| 453 | ◐ | pertaining to the eigenvalues on iR and let $Q$ be the faithful normal conditional expectation from $M$ onto $N$ (Proposition 4.3.7) | eigenvalues in iR ... (Theorem 4.3.5) | Querverweis | Die bedingte Erwartung stammt aus Theorem 4.3.5 (i), nicht aus Prop. 4.3.7; sonst steht 'eigenvalues in iR' |
| 454 | • | a predual $M_*$ of a von Neumann-algebra $M$ | of a von Neumann algebra $M$ | Satz | |
| 454 | ◐ | 0 is a pole of the resolvent of its generator $B$ with residuum $P$ = 1 ⊗ 1 | with residue $P$ = 1 ⊗ 1 | Konsistenz | Sonst immer 'residue' |
| | | forms a strongly continuous | one-parameter | | |

| | | | | | |
|---|---|---|---|---|---|
| 454 | ◐ | one parameter semigroup | semigroup | Konsistenz | |
| 456 | ○ | $H$0 is invariant under the extension to $H\varphi$ and for the multiplication maps we defined in Remark 3.1.3. | $H$0 is invariant under the extensions to $H\varphi$ and under the multiplication maps defined in Remark 3.1.3. | Grammatik | Satz grammatisch verunglückt |
| 457 | ○ | then by Lemma 3.1.2, there exists a sequence of normal states in Fix(R') with orthogonal supports | then by Lemma 4.4.1, there exists … | Querverweis | Die Existenz einer Folge von Zuständen mit orthogonalen Trägern ist Lemma 4.4.1; bitte prüfen |
| 459 | • | Without further comments, we will now use following facts. | we will now use the following facts. | Grammatik | |
| 459 | ◐ | if and only if it converges in $\sigma(M', M'')$-topology | converges in the $\sigma(M', M'')$-topology | Grammatik | |
| 461 | • | hence Theorem 4.4.2 (iv) is proved. | hence Theorem 4.4.5 (iv) is proved. | Querverweis | 4.4.2 ist eine Remark; gemeint ist Theorem 4.4.5 |
| 461 | • | exists in the strong operator topology with $A'$ is the generator of T'. | where $A'$ is the generator of T'. | Grammatik | |
| 463 | • | U. Groh & B. Kümmerer: Bibounded operators $W_*$-algebras. | Bibounded operators on W$_*$-algebras. | Literatur | 'on' fehlt im Titel |
| 463 | ○ | E. Størmer: On projection maps on von Neumann algebras. Math. Scand. 50 (1972) 42–50. | Math. Scand. 30 (1972) … | Literatur | Band 50 ist von 1982 (vgl. Groh & Kümmerer); 1972 entspricht Band 30. Bitte Band und Seiten prüfen |

## Updated Notes

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 468 | • | to the spectral theory of $C$0-semigroups, as in Chapter 3.6.6, was started | as in Part-I, Section 3.6.6 | Querverweis | 'Chapter' statt 'Section' |
| 468 | • | Using this, El-Mennaoui & Engel [30] gives a characterization | El-Mennaoui & Engel [30] give a characterization | Grammatik | |
| 469 | ◐ | (i.e., stability of the semigroup induced on an ultra power of the underlying Banach space) | ultrapower | Konsistenz | Sonst immer 'ultrapower' |

| | | | | | |
|---|---|---|---|---|---|
| 470 | • | Quantative estimates of the decay of the semigroup | Quantitative estimates | Tippfehler | |
| 471 | • | [16] J. Diestel: A survey of results related to the Dunford-Pettis property. … Contemp. Math. 2, 15-60 (1980). 1980. | … Contemp. Math. 2 (1980) 15–60. | Literatur | Jahr doppelt, Format weicht ab, Bindestrich statt Halbgeviertstrich |
| 471 | ○ | [15] R. Derndinger: Über das Spektrum positiver Generatoren. Math. Z. 172 (1980). | Math. Z. 172 (1980) 281–293. | Literatur | Seitenangaben fehlen (bitte prüfen) |
| 471 | ○ | Institute of Mathematics 2007 pp. 71–109 Available from: | … 2007 pp. 71–109. Available from: | Literatur | Punkt fehlt; ebenso bei Groh [13] S. 491 ('C∗-Algebras. 2026 Available from') |
| 474 | • | As examples for the rich litterature we mention | the rich literature | Tippfehler | |
| 474 | ◐ | by the work of I. Mezic [36] in the context of numerical problems using the dynamical mode decomposition | I. Mezić [36] … dynamic mode decomposition | Konsistenz | Im Literaturverzeichnis 'Mezić'; der übliche Fachbegriff lautet 'dynamic mode decomposition' |
| 475 | • | persistent irreducibilty implies that the generator | persistent irreducibility | Tippfehler | |
| 475 | • | In Gerlach [21] strong convergence of Feller semigoups is studied. | Feller semigroups | Tippfehler | |
| 475 | ○ | whose positive cone has non empty interior | non-empty interior | Konsistenz | sonst 'non-empty' |
| 477 | ◐ | [29] N. Jacob: Pseudo Differential Operators and Markov Processes. Vol. 1: … [30] N. Jacob: Pseudo Differential Operators and Markov Processes. Vol III: | Pseudo-Differential Operators and Markov Processes. Vol. I: … / Vol. III: | Literatur | Uneinheitlich gegenüber [28] 'Pseudo-Differential … Vol. II'; Punkt nach 'Vol' fehlt bei [30] |
| 480 | ◐ | a generalisation of Kato’s inequality … to characterise positivity … to characterise irreducibility | generalization / characterize (AE wie im übrigen Text) | Konsistenz | Sonst amerikanische Schreibung (behavior, characterize, analyzed) |
| 480 | ◐ | see Vogt & Voigt [75], Stein & Voigt [73], Stein et al. [72] | … Stein et al. [72]. | Satz | Schlusspunkt fehlt |
| 481 | • | Even the spectrum of the generator is independent on p in this case | independent of p | Grammatik | |

| | | | | | |
|---|---|---|---|---|---|
| 481 | • | The domain of its generator can be described explicitely | explicitly | Tippfehler | |
| 481 | ◐ | Kernel estimates for this semigroup are obtained in Elst & Ouhabaz [27] | … Elst & Ouhabaz [27]. | Satz | Schlusspunkt fehlt |
| 481 | ○ | (see also the extended notes of Part-II, Chapter 2 concerning Koopman semigroups) | the updated notes of Part-II, Chapter 2 | Konsistenz | Sonst 'Updated Notes' |
| 482 | ◐ | the Markov conjecture: Any generator of a strongly continuous positive semigroup on ℓ1 … has a bounded generator. | Any strongly continuous positive semigroup on ℓ1 which … extends to a group has a bounded generator. | Grammatik | 'generator … has a bounded generator' ist doppelt gemoppelt |
| 484 | ◐ | based on the Jacobs-DeLeeuw-Glicksberg Theorem | Jacobs–de Leeuw–Glicksberg Theorem | Konsistenz | S. 490: 'Jacobs-de Leeuw-Glicksberg' |
| 484 | ○ | Glück & Haase [38] introduce the notion semigroup at infinity | introduce the notion of a semigroup at infinity | Grammatik | |
| 489 | • | This also follows from the example given above above and the fact | from the example given above and the fact | Tippfehler | Wort doppelt |
| 490 | • | By Greiner et al. [12, Theorem 5.3.1]] one has | By Greiner et al. [12, Theorem 5.3.1] one has | Satz | doppelte schließende Klammer |
| 490 | ◐ | Chapter 4 As in Part-IV, Chapter 4, results from Glück & Mironchenko [10] | As in Part-II, Chapter 4, results from Glück & Mironchenko … | Querverweis | Der Abschnitt gehört selbst zu Part IV, Chapter 4; Glück & Mironchenko wird in den Updated Notes zu Part II, Chapter 4 zitiert |
| 491 | ◐ | C. W. Henson & L. C. j. Moore: Nonstandard hulls | C. W. Henson & L. C. Moore Jr.: | Literatur | BibTeX-Artefakt aus 'Jr.' |

## Index

| S. | | Ist | Soll | Art | Kommentar |
|---|---|---|---|---|---|
| 494 | ◐ | Kato’s equality, 157–161, 317, 318, 360 | Kato’s inequality, … | Tippfehler | Index; im Text 'Kato’s inequality' |
| 495 | ○ | F-Product semigroup, 25 | F-product semigroup, 25 | Konsistenz | Index, sonst 'F-product' |

*Hinweis:* Die Liste beruht auf dem aus dem PDF extrahierten Text. Verdächtige Stellen (zusammengeklebte Wörter, Trennungen) wurden am PDF-Bild gegengeprüft. Formeln wurden bewusst ausgelassen.