

CMPE 322/327 - Theory of Computation

Week 3: Nondeterministic Finite State Automata & Epsilon Transitions

Burak Ekici

March 7-11, 2022

Outline

- 1 A Quick Recap
- 2 Nondeterministic Finite Automata
- 3 Epsilon Transitions
- 4 Closure Properties

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with
 - 1 Q : finite set of states

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with
 - 1 Q : finite set of states
 - 2 Σ : input alphabet

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with
 - ① Q : finite set of states
 - ② Σ : input alphabet
 - ③ $\delta: Q \times \Sigma \rightarrow Q$: transition function

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with
 - 1 Q : finite set of states
 - 2 Σ : input alphabet
 - 3 $\delta : Q \times \Sigma \rightarrow Q$: transition function
 - 4 $s \in Q$: start state

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with
 - ① Q : finite set of states
 - ② Σ : input alphabet
 - ③ $\delta : Q \times \Sigma \rightarrow Q$: transition function
 - ④ $s \in Q$: start state
 - ⑤ $F \subseteq Q$: final (accept) states

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\delta : Q \times \Sigma \rightarrow Q$: transition function
- ④ $s \in Q$: start state
- ⑤ $F \subseteq Q$: final (accept) states

- $\hat{\delta} : Q \times \Sigma^* \rightarrow Q$ is inductively defined by

$$\hat{\delta}(q, \varepsilon) := q$$

$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\delta : Q \times \Sigma \rightarrow Q$: transition function
- ④ $s \in Q$: start state
- ⑤ $F \subseteq Q$: final (accept) states

- $\hat{\delta} : Q \times \Sigma^* \rightarrow Q$ is inductively defined by

$$\hat{\delta}(q, \varepsilon) := q$$

$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$

- string $x \in \Sigma^*$ is **accepted** by M if $\hat{\delta}(s, x) \in F$

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\delta: Q \times \Sigma \rightarrow Q$: transition function
- ④ $s \in Q$: start state
- ⑤ $F \subseteq Q$: final (accept) states

- $\hat{\delta}: Q \times \Sigma^* \rightarrow Q$ is inductively defined by

$$\hat{\delta}(q, \varepsilon) := q$$

$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$

- string $x \in \Sigma^*$ is **accepted** by M if $\hat{\delta}(s, x) \in F$
- string $x \in \Sigma^*$ is **rejected** by M if $\hat{\delta}(s, x) \notin F$

Definitions

- **deterministic finite automaton (DFA)** is quintuple $M = (Q, \Sigma, \delta, s, F)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\delta : Q \times \Sigma \rightarrow Q$: transition function
- ④ $s \in Q$: start state
- ⑤ $F \subseteq Q$: final (accept) states

- $\hat{\delta} : Q \times \Sigma^* \rightarrow Q$ is inductively defined by


$$\hat{\delta}(q, \varepsilon) := q$$


$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$


- string $x \in \Sigma^*$ is **accepted** by M if $\hat{\delta}(s, x) \in F$
- string $x \in \Sigma^*$ is **rejected** by M if $\hat{\delta}(s, x) \notin F$
- language accepted by M is given by $L(M) := \{x \mid \hat{\delta}(x, s) \in F\}$


Example (DFA \rightarrow Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$




Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ ① $Q = \{1, 2, 3\}$

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ ① $Q = \{1, 2, 3\}$ ② $\Sigma = \{a, b\}$

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ ① $Q = \{1, 2, 3\}$ ② $\Sigma = \{a, b\}$ ③ $\delta : Q \times \Sigma \rightarrow Q$


Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ ① $Q = \{1, 2, 3\}$ ② $\Sigma = \{a, b\}$ ③ $\delta : Q \times \Sigma \rightarrow Q$

δ	a	b
1	1	2
2	3	3
3	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 


- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta : Q \times \Sigma \rightarrow Q$
- ④ $s = 1$

δ	a	b
1	1	2
2	3	3
3	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta : Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3


Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- 1 $Q = \{1, 2, 3\}$
- 2 $\Sigma = \{a, b\}$
- 3 $\delta : Q \times \Sigma \rightarrow Q$
- 4 $s = 1$
- 5 $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b
1 1 1 1 1 2


Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- 1 $Q = \{1, 2, 3\}$
- 2 $\Sigma = \{a, b\}$
- 3 $\delta: Q \times \Sigma \rightarrow Q$
- 4 $s = 1$
- 5 $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b
1 1 1 1 1 2

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 


- 1 $Q = \{1, 2, 3\}$
- 2 $\Sigma = \{a, b\}$
- 3 $\delta : Q \times \Sigma \rightarrow Q$
- 4 $s = 1$
- 5 $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 


- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta : Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 


- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta : Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta : Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3


Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- 1 $Q = \{1, 2, 3\}$
- 2 $\Sigma = \{a, b\}$
- 3 $\delta : Q \times \Sigma \rightarrow Q$
- 4 $s = 1$
- 5 $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b
1 1 1 1 1 2


Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- 1 $Q = \{1, 2, 3\}$
- 2 $\Sigma = \{a, b\}$
- 3 $\delta : Q \times \Sigma \rightarrow Q$
- 4 $s = 1$
- 5 $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3


Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- 1 $Q = \{1, 2, 3\}$
- 2 $\Sigma = \{a, b\}$
- 3 $\delta : Q \times \Sigma \rightarrow Q$
- 4 $s = 1$
- 5 $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- 1 $Q = \{1, 2, 3\}$
- 2 $\Sigma = \{a, b\}$
- 3 $\delta : Q \times \Sigma \rightarrow Q$
- 4 $s = 1$
- 5 $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1 1 1 1 1 2

Example (DFA \rightarrow Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

Example (DFA \rightarrow Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
a	a	b	b	a	
1	1	1	2	3	3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

Example (DFA \rightarrow Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
a	a	b	b	a	
1	1	1	2	3	3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3


Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
a	a	b	b	a	
1	1	1	2	3	3

Example (DFA \rightarrow Regular Sets) $M = (Q, \Sigma, \delta, s, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

a a a a b $\in L(M)$


1 1 1 1 1 2

a a b b a

1 1 1 2 3 3

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
	a	a	b	b	a	$\notin L(M)$
1	1	1	2	3	3	

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$


δ	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	2	
	a	a	b	b	a	$\notin L(M)$
1	1	1	2	3	3	

$L(M) := \{x \mid \dots\}$

Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\delta: Q \times \Sigma \rightarrow Q$
- ④ $s = 1$
- ⑤ $F = \{2\}$

δ	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	2	
	a	a	b	b	a	$\notin L(M)$
1	1	1	2	3	3	

$$L(M) := \{x \mid x = a^n b, n \geq 0\}$$

Definition

set $A \subseteq \Sigma^*$ is **regular** if $A = L(M)$ for some DFA M

Definition

set $A \subseteq \Sigma^*$ is **regular** if $A = L(M)$ for some DFA M

Theorem

regular sets are effectively closed under **intersection**, **complement** and **union**

Outline

- 1 A Quick Recap
- 2 Nondeterministic Finite Automata
- 3 Epsilon Transitions
- 4 Closure Properties

Definitions

- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with

Definitions

- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with
 - 1 Q : finite set of states

Definitions

- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with
 - 1 Q : finite set of states
 - 2 Σ : input alphabet

Definitions

- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with
 - ① Q : finite set of states
 - ② Σ : input alphabet
 - ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$: transition function

Definitions


- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with
 - 1 Q : finite set of states
 - 2 Σ : input alphabet
 - 3 $\Delta : Q \times \Sigma \rightarrow 2^Q$: transition function
 - 4 $S \subseteq Q$: set of start states

Definitions

- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with
 - ① Q : finite set of states
 - ② Σ : input alphabet
 - ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$: transition function
 - ④ $S \subseteq Q$: set of start states
 - ⑤ $F \subseteq Q$: final (accept) states


Example

$N = (Q, \Sigma, \Delta, S, F)$



Example


$N = (Q, \Sigma, \Delta, S, F)$



① $Q = \{1, 2, 3\}$

Example


$N = (Q, \Sigma, \Delta, S, F)$



① $Q = \{1, 2, 3\}$


② $\Sigma = \{a, b\}$

Example

 $N = (Q, \Sigma, \Delta, S, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$


Example

 $N = (Q, \Sigma, \Delta, S, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$

Δ	a	b
1	{1, 2}	{1}
2	{3}	{1, 3}
3	{3}	\emptyset


Example

 $N = (Q, \Sigma, \Delta, S, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$
- ④ $S = \{1\}$

Δ	a	b
1	{1, 2}	{1}
2	{3}	{1, 3}
3	{3}	\emptyset

Example

 $N = (Q, \Sigma, \Delta, S, F)$ 

- ① $Q = \{1, 2, 3\}$
- ② $\Sigma = \{a, b\}$
- ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$
- ④ $S = \{1\}$
- ⑤ $F = \{3\}$

Δ	a	b
1	{1, 2}	{1}
2	{3}	{1, 3}
3	{3}	\emptyset

Definitions

- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$: transition function
- ④ $S \subseteq Q$: set of start states
- ⑤ $F \subseteq Q$: final (accept) states

- $\widehat{\Delta} : 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}(A, \epsilon) := A$$

$$\widehat{\Delta}(A, xa) := \bigcup_{q \in \widehat{\Delta}(A, x)} \Delta(q, a)$$

Definitions

- **nondeterministic finite automaton (NFA)** is quintuple $N = (Q, \Sigma, \Delta, S, F)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\Delta : Q \times \Sigma \rightarrow 2^Q$: transition function
- ④ $S \subseteq Q$: set of start states
- ⑤ $F \subseteq Q$: final (accept) states

- $\widehat{\Delta} : 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}(A, \epsilon) := A$$

$$\widehat{\Delta}(A, xa) := \bigcup_{q \in \widehat{\Delta}(A, x)} \Delta(q, a)$$

- string $x \in \Sigma^*$ is **accepted** by N if $\widehat{\Delta}(S, x) \cap F \neq \emptyset$

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup (q \in \widehat{\Delta}(A, ababb), a)$

1st rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup (q \in \widehat{\Delta}(A, ababb), a)$$

$$\bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$$

1st rec. call

2nd rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup (q \in \widehat{\Delta}(A, ababb), a)$$

$$\bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$$

$$\bigcup (q \in \bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$$

1st rec. call

2nd rec. call

3rd rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup (q \in \widehat{\Delta}(A, ababb), a)$$

1st rec. call

$$\bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$$

2nd rec. call

$$\bigcup (q \in \bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$$

3rd rec. call

$$\bigcup (q \in \bigcup (q \in \bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$$

4th rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup (q \in \widehat{\Delta}(A, ababb), a)$$

1st rec. call

$$\bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$$

2nd rec. call

$$\bigcup (q \in \bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$$

3rd rec. call

$$\bigcup (q \in \bigcup (q \in \bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$$

4th rec. call

$$\bigcup (q \in \bigcup (q \in \bigcup (q \in \bigcup (q \in \bigcup (q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$$

5th rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \epsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \varepsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in A \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \varepsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in A \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in B \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in A \Delta(q, a)) = B$

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \varepsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in A \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in B \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in A \Delta(q, a)) = B$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in C \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in B \Delta(q, b)) = C$

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \varepsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in A \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in B \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in A \Delta(q, a)) = B$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in C \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in B \Delta(q, b)) = C$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in D \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in C \Delta(q, a)) = D$

Example (Unfolding of the multistep function $\widehat{\Delta}$)Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \varepsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in A \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in B \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in A \Delta(q, a)) = B$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in C \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in B \Delta(q, b)) = C$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in D \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in C \Delta(q, a)) = D$
$\bigcup(q \in \bigcup(q \in E \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in D \Delta(q, b)) = E$

Example (Unfolding of the multistep function $\widehat{\Delta}$)Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \varepsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in A \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in B \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in A \Delta(q, a)) = B$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in C \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in B \Delta(q, b)) = C$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in D \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in C \Delta(q, a)) = D$
$\bigcup(q \in \bigcup(q \in E \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in D \Delta(q, b)) = E$
$\bigcup(q \in F \Delta(q, a))$	assuming $\bigcup(q \in E \Delta(q, b)) = F$

Example (Unfolding of the multistep function $\widehat{\Delta}$)

Let $x = ababba$ over the alphabet $\Sigma = \{a, b\}$

$\bigcup(q \in \widehat{\Delta}(A, ababb), a)$	1 st rec. call
$\bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, abab) \Delta(q, b)) \Delta(q, a))$	2 nd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, aba) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	3 rd rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, ab) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	4 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, a) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	5 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \widehat{\Delta}(A, \varepsilon) \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	6 th rec. call
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in A \Delta(q, a)) \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in B \Delta(q, b)) \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in A \Delta(q, a)) = B$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in \bigcup(q \in C \Delta(q, a)) \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in B \Delta(q, b)) = C$
$\bigcup(q \in \bigcup(q \in \bigcup(q \in D \Delta(q, b)) \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in C \Delta(q, a)) = D$
$\bigcup(q \in \bigcup(q \in E \Delta(q, b)) \Delta(q, a))$	assuming $\bigcup(q \in D \Delta(q, b)) = E$
$\bigcup(q \in F \Delta(q, a))$	assuming $\bigcup(q \in E \Delta(q, b)) = F$
G	assuming $\bigcup(q \in F \Delta(q, a)) = G$

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Proof.

We argue by induction on $|y|$:

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Proof.

We argue by induction on $|y|$:

- base case: $|y| = 0$ thus $y = \varepsilon$

$$\widehat{\Delta}(A, x\varepsilon) = \widehat{\Delta}(A, x) = \widehat{\Delta}(\widehat{\Delta}(A, x), \varepsilon)$$

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Proof.

We argue by induction on $|y|$:

- base case: $|y| = 0$ thus $y = \varepsilon$

$$\widehat{\Delta}(A, x\varepsilon) = \widehat{\Delta}(A, x) = \widehat{\Delta}(\widehat{\Delta}(A, x), \varepsilon)$$

- step case: $|y| > 0$ thus $y = zb$ s.t. $|z| = |y| - 1$ with IH: $\widehat{\Delta}(A, xz) = \widehat{\Delta}(\widehat{\Delta}(A, x), z)$

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Proof.

We argue by induction on $|y|$:

- base case: $|y| = 0$ thus $y = \varepsilon$

$$\widehat{\Delta}(A, x\varepsilon) = \widehat{\Delta}(A, x) = \widehat{\Delta}(\widehat{\Delta}(A, x), \varepsilon)$$

- step case: $|y| > 0$ thus $y = zb$ s.t. $|z| = |y| - 1$ with IH: $\widehat{\Delta}(A, xz) = \widehat{\Delta}(\widehat{\Delta}(A, x), z)$

$$\widehat{\Delta}(A, xzb) = \bigcup_{q \in \widehat{\Delta}(A, xz)} \Delta(q, b) \quad (\text{by definition of } \widehat{\Delta})$$

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Proof.

We argue by induction on $|y|$:

- base case: $|y| = 0$ thus $y = \varepsilon$

$$\widehat{\Delta}(A, x\varepsilon) = \widehat{\Delta}(A, x) = \widehat{\Delta}(\widehat{\Delta}(A, x), \varepsilon)$$

- step case: $|y| > 0$ thus $y = zb$ s.t. $|z| = |y| - 1$ with IH: $\widehat{\Delta}(A, xz) = \widehat{\Delta}(\widehat{\Delta}(A, x), z)$

$$\begin{aligned} \widehat{\Delta}(A, xzb) &= \bigcup_{q \in \widehat{\Delta}(A, xz)} \Delta(q, b) && \text{(by definition of } \widehat{\Delta}) \\ &= \bigcup_{q \in \widehat{\Delta}(\widehat{\Delta}(A, x), z)} \Delta(q, b) && \text{(by IH)} \end{aligned}$$

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Proof.

We argue by induction on $|y|$:

- base case: $|y| = 0$ thus $y = \varepsilon$

$$\widehat{\Delta}(A, x\varepsilon) = \widehat{\Delta}(A, x) = \widehat{\Delta}(\widehat{\Delta}(A, x), \varepsilon)$$

- step case: $|y| > 0$ thus $y = zb$ s.t. $|z| = |y| - 1$ with IH: $\widehat{\Delta}(A, xz) = \widehat{\Delta}(\widehat{\Delta}(A, x), z)$

$$\begin{aligned} \widehat{\Delta}(A, xzb) &= \bigcup_{q \in \widehat{\Delta}(A, xz)} \Delta(q, b) && \text{(by definition of } \widehat{\Delta}) \\ &= \bigcup_{q \in \widehat{\Delta}(\widehat{\Delta}(A, x), z)} \Delta(q, b) && \text{(by IH)} \\ &= \widehat{\Delta}(\widehat{\Delta}(A, x), zb) && \text{(by definition of } \widehat{\Delta}) \end{aligned}$$

Lemma ($\widehat{\Delta}$ distributes)

$$\widehat{\Delta}(A, xy) = \widehat{\Delta}(\widehat{\Delta}(A, x), y) \quad \forall A \subseteq Q_N \text{ and } x, y \in \Sigma^*$$

Proof.

We argue by induction on $|y|$:

- base case: $|y| = 0$ thus $y = \varepsilon$

$$\widehat{\Delta}(A, x\varepsilon) = \widehat{\Delta}(A, x) = \widehat{\Delta}(\widehat{\Delta}(A, x), \varepsilon)$$

- step case: $|y| > 0$ thus $y = zb$ s.t. $|z| = |y| - 1$ with IH: $\widehat{\Delta}(A, xz) = \widehat{\Delta}(\widehat{\Delta}(A, x), z)$

$$\begin{aligned} \widehat{\Delta}(A, xzb) &= \bigcup_{q \in \widehat{\Delta}(A, xz)} \Delta(q, b) && \text{(by definition of } \widehat{\Delta}) \\ &= \bigcup_{q \in \widehat{\Delta}(\widehat{\Delta}(A, x), z)} \Delta(q, b) && \text{(by IH)} \\ &= \widehat{\Delta}(\widehat{\Delta}(A, x), zb) && \text{(by definition of } \widehat{\Delta}) \\ &= \widehat{\Delta}(\widehat{\Delta}(A, x), y) \end{aligned}$$

Theorem

every set accepted by NFA is regular

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - 1 $Q_M := 2^{Q_N}$

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - 1 $Q_M := 2^{Q_N}$
 - 2 $\delta_M(A, a) := \widehat{\Delta}(A, a) \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - ① $Q_M := 2^{Q_N}$
 - ② $\delta_M(A, a) := \widehat{\Delta}(A, a) \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$
 - ③ $s_M := S_N$

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - ① $Q_M := 2^{Q_N}$
 - ② $\delta_M(A, a) := \widehat{\Delta}(A, a) \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$
 - ③ $s_M := S_N$
 - ④ $F_M := \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\}$

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - ① $Q_M := 2^{Q_N}$
 - ② $\delta_M(A, a) := \widehat{\Delta}(A, a) \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$
 - ③ $s_M := S_N$
 - ④ $F_M := \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\}$
- claim: $\widehat{\delta_M}(A, x) = \widehat{\Delta}(A, x) \quad \forall A \subseteq Q \text{ and } x \in \Sigma^*$

Theorem

every set accepted by NFA is regular

Proof.

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - ① $Q_M := 2^{Q_N}$
 - ② $\delta_M(A, a) := \widehat{\Delta}(A, a) \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$
 - ③ $s_M := S_N$
 - ④ $F_M := \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\}$
- claim: $\widehat{\delta_M}(A, x) = \widehat{\Delta}(A, x) \quad \forall A \subseteq Q \text{ and } x \in \Sigma^*$
 proof: by induction on $|x|$ see next slide

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

- base case: $|x| = 0$ thus $x = \varepsilon$

$$\widehat{\delta}_M(A, \varepsilon) = A = \widehat{\Delta}_N(A, \varepsilon)$$

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

- base case: $|x| = 0$ thus $x = \varepsilon$

$$\widehat{\delta}_M(A, \varepsilon) = A = \widehat{\Delta}_N(A, \varepsilon)$$

- step case: $|x| > 0$ thus $x = ya$ s.t. $|y| = |x| - 1$ with IH: $\widehat{\delta}_M(A, y) = \widehat{\Delta}_N(A, y)$

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

- base case: $|x| = 0$ thus $x = \varepsilon$

$$\widehat{\delta}_M(A, \varepsilon) = A = \widehat{\Delta}_N(A, \varepsilon)$$

- step case: $|x| > 0$ thus $x = ya$ s.t. $|y| = |x| - 1$ with IH: $\widehat{\delta}_M(A, y) = \widehat{\Delta}_N(A, y)$

$$\widehat{\delta}_M(A, ya) = \delta_M(\widehat{\delta}_M(A, y), a) \quad (\text{by definition of } \widehat{\delta}_M)$$

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

- base case: $|x| = 0$ thus $x = \varepsilon$

$$\widehat{\delta}_M(A, \varepsilon) = A = \widehat{\Delta}_N(A, \varepsilon)$$

- step case: $|x| > 0$ thus $x = ya$ s.t. $|y| = |x| - 1$ with IH: $\widehat{\delta}_M(A, y) = \widehat{\Delta}_N(A, y)$

$$\begin{aligned} \widehat{\delta}_M(A, ya) &= \delta_M(\widehat{\delta}_M(A, y), a) && \text{(by definition of } \widehat{\delta}_M) \\ &= \delta_M(\widehat{\Delta}_N(A, y), a) && \text{(by induction hypothesis IH)} \end{aligned}$$

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

- base case: $|x| = 0$ thus $x = \varepsilon$

$$\widehat{\delta}_M(A, \varepsilon) = A = \widehat{\Delta}_N(A, \varepsilon)$$

- step case: $|x| > 0$ thus $x = ya$ s.t. $|y| = |x| - 1$ with IH: $\widehat{\delta}_M(A, y) = \widehat{\Delta}_N(A, y)$

$$\begin{aligned}\widehat{\delta}_M(A, ya) &= \delta_M(\widehat{\delta}_M(A, y), a) && \text{(by definition of } \widehat{\delta}_M) \\ &= \delta_M(\widehat{\Delta}_N(A, y), a) && \text{(by induction hypothesis IH)} \\ &= \widehat{\Delta}_N(\widehat{\Delta}_N(A, y), a) && \text{(by definition of } \delta_M)\end{aligned}$$

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

- base case: $|x| = 0$ thus $x = \varepsilon$

$$\widehat{\delta}_M(A, \varepsilon) = A = \widehat{\Delta}_N(A, \varepsilon)$$

- step case: $|x| > 0$ thus $x = ya$ s.t. $|y| = |x| - 1$ with IH: $\widehat{\delta}_M(A, y) = \widehat{\Delta}_N(A, y)$

$$\begin{aligned}\widehat{\delta}_M(A, ya) &= \delta_M(\widehat{\delta}_M(A, y), a) && \text{(by definition of } \widehat{\delta}_M) \\ &= \delta_M(\widehat{\Delta}_N(A, y), a) && \text{(by induction hypothesis IH)} \\ &= \widehat{\Delta}_N(\widehat{\Delta}_N(A, y), a) && \text{(by definition of } \delta_M) \\ &= \widehat{\Delta}_N(A, ya) && \text{(by distributivity of } \widehat{\Delta})\end{aligned}$$

proof of the claim

claim: $\widehat{\delta}_M(A, x) = \widehat{\Delta}_N(A, x) \quad \forall A \subseteq Q_N \text{ and } x \in \Sigma^*$

- base case: $|x| = 0$ thus $x = \varepsilon$

$$\widehat{\delta}_M(A, \varepsilon) = A = \widehat{\Delta}_N(A, \varepsilon)$$

- step case: $|x| > 0$ thus $x = ya$ s.t. $|y| = |x| - 1$ with IH: $\widehat{\delta}_M(A, y) = \widehat{\Delta}_N(A, y)$

$$\begin{aligned}\widehat{\delta}_M(A, ya) &= \delta_M(\widehat{\delta}_M(A, y), a) && \text{(by definition of } \widehat{\delta}_M) \\ &= \delta_M(\widehat{\Delta}_N(A, y), a) && \text{(by induction hypothesis IH)} \\ &= \widehat{\Delta}_N(\widehat{\Delta}_N(A, y), a) && \text{(by definition of } \delta_M) \\ &= \widehat{\Delta}_N(A, ya) && \text{(by distributivity of } \widehat{\Delta}) \\ &= \widehat{\Delta}_N(A, x)\end{aligned}$$



Proof. (NFA regularity)

statement: $L(M) = L(N)$

Proof. (NFA regularity)

statement: $L(M) = L(N)$

$$\forall x \in \Sigma^*, x \in L(M) \iff \widehat{\delta}_M(s_M, x) \in F_M \quad (\text{by definition of acceptance})$$

Proof. (NFA regularity)

statement: $L(M) = L(N)$

$$\begin{aligned} \forall x \in \Sigma^*, x \in L(M) &\iff \widehat{\delta}_M(s_M, x) \in F_M && \text{(by definition of acceptance)} \\ &\iff \widehat{\delta}_M(s_N, x) \in \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\} && \text{(by definition of } s_M \text{ and } F_M) \end{aligned}$$

Proof. (NFA regularity)

statement: $L(M) = L(N)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M) &\iff \widehat{\delta}_M(s_M, x) \in F_M && \text{(by definition of acceptance)} \\
 &\iff \widehat{\delta}_M(s_N, x) \in \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\} && \text{(by definition of } s_M \text{ and } F_M) \\
 &\iff \widehat{\Delta}_N(s_N, x) \in \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\} && \text{(by claim proven in slide 12)}
 \end{aligned}$$

Proof. (NFA regularity)

statement: $L(M) = L(N)$

$$\begin{aligned}\forall x \in \Sigma^*, x \in L(M) &\iff \widehat{\delta}_M(s_M, x) \in F_M && \text{(by definition of acceptance)} \\ &\iff \widehat{\delta}_M(s_N, x) \in \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\} && \text{(by definition of } s_M \text{ and } F_M) \\ &\iff \widehat{\Delta}_N(s_N, x) \in \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\} && \text{(by claim proven in slide 12)} \\ &\iff \widehat{\Delta}_N(s_N, x) \cap F_N \neq \emptyset && \text{(by set comprehension)}\end{aligned}$$


Proof. (NFA regularity)

statement: $L(M) = L(N)$

$$\begin{aligned} \forall x \in \Sigma^*, x \in L(M) &\iff \widehat{\delta}_M(s_M, x) \in F_M && \text{(by definition of acceptance)} \\ &\iff \widehat{\delta}_M(s_N, x) \in \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\} && \text{(by definition of } s_M \text{ and } F_M) \\ &\iff \widehat{\Delta}_N(s_N, x) \in \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\} && \text{(by claim proven in slide 12)} \\ &\iff \widehat{\Delta}_N(s_N, x) \cap F_N \neq \emptyset && \text{(by set comprehension)} \\ &\iff x \in L(N) && \text{(by definition of acceptance)} \end{aligned}$$



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$

Δ	a	b

Δ	a	b

Example



$$A = \emptyset$$

$$B = \{1\}$$

$$C = \{2\}$$

$$D = \{3\}$$

$$E = \{1, 2\}$$

$$F = \{1, 3\}$$


$$G = \{2, 3\}$$

$$H = \{1, 2, 3\}$$

Δ	<i>a</i>	<i>b</i>
A	A	A

Δ	<i>a</i>	<i>b</i>

Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$

Δ	<i>a</i>	<i>b</i>
A	A	A
B	E	B

Δ	<i>a</i>	<i>b</i>

Example



$$A = \emptyset$$

$$B = \{1\}$$

$$C = \{2\}$$

$$D = \{3\}$$

$$E = \{1, 2\}$$

$$F = \{1, 3\}$$


$$G = \{2, 3\}$$

$$H = \{1, 2, 3\}$$

Δ	<i>a</i>	<i>b</i>
A	A	A
B	E	B
C	D	F

Δ	<i>a</i>	<i>b</i>

Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$

Δ	<i>a</i>	<i>b</i>
A	A	A
B	E	B
C	D	F
D	D	A

Δ	<i>a</i>	<i>b</i>


Example


 $A = \emptyset$
 $B = \{1\}$
 $C = \{2\}$
 $D = \{3\}$
 $E = \{1, 2\}$
 $F = \{1, 3\}$
 $G = \{2, 3\}$
 $H = \{1, 2, 3\}$

Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$


$F = \{1, 3\}$

$G = \{2, 3\}$


$H = \{1, 2, 3\}$

Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

*a*bbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

ab**b**baababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

ab**bb**aababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbb**a**ababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbbaababbabbbaababba



Example




$A = \emptyset$
 $B = \{1\}$
 $C = \{2\}$
 $D = \{3\}$

$E = \{1, 2\}$
 $F = \{1, 3\}$
 $G = \{2, 3\}$
 $H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
A	A	A
B	E	B
C	D	F
D	D	A

Δ	<i>a</i>	<i>b</i>
E	H	F
F	H	B
G	D	F
H	H	F

abbbaabababbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababba**bb**baababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$A = \emptyset$

$B = \{1\}$

$C = \{2\}$

$D = \{3\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$G = \{2, 3\}$

$H = \{1, 2, 3\}$


Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>
<i>B</i>	<i>E</i>	<i>B</i>
<i>C</i>	<i>D</i>	<i>F</i>
<i>D</i>	<i>D</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>
<i>E</i>	<i>H</i>	<i>F</i>
<i>F</i>	<i>H</i>	<i>B</i>
<i>G</i>	<i>D</i>	<i>F</i>
<i>H</i>	<i>H</i>	<i>F</i>

abbbaababbabbbaababba



Example



$B = \{1\}$

$E = \{1, 2\}$

$F = \{1, 3\}$


$H = \{1, 2, 3\}$

Δ	a	b
B	E	B

Δ	a	b
E	H	F
F	H	B
H	H	F

abbbaababbabbbaababba

remove inaccessible states



Question

Every regular set is accepted by ...

- A ... an NFA having exactly one final state,
- B ... a DFA having exactly one final state,
- C ... an NFA having exactly one start state.

Outline

- 1 A Quick Recap
- 2 Nondeterministic Finite Automata
- 3 Epsilon Transitions**
- 4 Closure Properties

Definitions

- NFA with ε -transitions (NFA_ε) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

Definitions

- NFA with ε -transitions (NFA_ε) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


Definitions

- NFA with ε -transitions (NFA_ε) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that
 - ① $\varepsilon \notin \Sigma$
 - ② $N_\varepsilon = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

Definitions

- NFA with ε -transitions (NFA_ε) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that
 - 1 $\varepsilon \notin \Sigma$
 - 2 $N_\varepsilon = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$


Example



Definitions

- NFA with ε -transitions (NFA_ε) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that
 - 1 $\varepsilon \notin \Sigma$
 - 2 $N_\varepsilon = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$
- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$


Example



Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that
 - 1 $\varepsilon \notin \Sigma$
 - 2 $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$
- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


② $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$
- $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \varepsilon) = C_{\varepsilon}(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_{\varepsilon}(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


② $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$
- $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \varepsilon) = C_{\varepsilon}(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_{\varepsilon}(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

$$\widehat{\Delta}_N(\{1\}, b) =$$

Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


② $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$
- $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \varepsilon) = C_{\varepsilon}(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_{\varepsilon}(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

$$\widehat{\Delta}_N(\{1\}, b) = C_{\varepsilon}(\{1\})$$

Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


② $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$
- $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \varepsilon) = C_{\varepsilon}(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_{\varepsilon}(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

$$\widehat{\Delta}_N(\{1\}, b) = C_{\varepsilon}(\Delta(1, b)) \cup C_{\varepsilon}(\Delta(2, b)) \cup C_{\varepsilon}(\Delta(3, b))$$

Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


② $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$
- $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \varepsilon) = C_{\varepsilon}(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_{\varepsilon}(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

$$\widehat{\Delta}_N(\{1\}, b) = C_{\varepsilon}(\emptyset) \cup C_{\varepsilon}(\{2\}) \cup C_{\varepsilon}(\emptyset)$$

Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


② $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$
- $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \varepsilon) = C_{\varepsilon}(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_{\varepsilon}(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

$$\widehat{\Delta}_N(\{1\}, b) = \emptyset \cup \{2, 3\} \cup \emptyset$$

Definitions

- NFA with ε -transitions (NFA $_{\varepsilon}$) is sextuple $N = (Q, \Sigma, \varepsilon, \Delta, S, F)$ such that

① $\varepsilon \notin \Sigma$


② $N_{\varepsilon} = (Q, \Sigma \cup \{\varepsilon\}, \Delta, S, F)$ is NFA over alphabet $\Sigma \cup \{\varepsilon\}$

- $\Delta: Q \times \Sigma \cup \{\varepsilon\} \rightarrow 2^Q$
- ε -closure of set $A \subseteq Q$ is defined as $C_{\varepsilon}(A) = \bigcup \{\widehat{\Delta}_{N_{\varepsilon}}(A, x) \mid x \in \{\varepsilon\}^*\}$
- $\widehat{\Delta}_N: 2^Q \times \Sigma^* \rightarrow 2^Q$ is inductively defined by

$$\widehat{\Delta}_N(A, \varepsilon) = C_{\varepsilon}(A)$$

$$\widehat{\Delta}_N(A, xa) = \bigcup \{C_{\varepsilon}(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, x)\}$$

Example



$$C_{\varepsilon}(\{1\}) = \{1, 2, 3\}$$

$$\widehat{\Delta}_N(\{1\}, b) = \{2, 3\}$$

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, ba)\}$$

1st rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, ba)\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, b)\}\}$$

1st rec. call

2nd rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, ba)\}$$

1st rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, b)\}\}$$

2nd rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in \widehat{\Delta}_N(A, \varepsilon)\}\}\}$$

3rd rec. call

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, ba)\}$$

1st rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, b)\}\}$$

2nd rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in \widehat{\Delta}_N(A, \varepsilon)\}\}\}$$

3rd rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in C_\varepsilon(A)\}\}\}$$

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, ba)\}$$

1st rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, b)\}\}$$

2nd rec. call

$$\bigcup \left\{ C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \left\{ C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in \widehat{\Delta}_N(A, \varepsilon)\} \right\} \right\}$$

3rd rec. call

$$\bigcup \left\{ C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \left\{ C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in C_\varepsilon(A)\} \right\} \right\}$$

$$\bigcup \left\{ C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \mathbf{B}\} \right\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in C_\varepsilon(A)\} = B$$

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, ba)\}$$

1st rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, b)\}\}$$

2nd rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in \widehat{\Delta}_N(A, \varepsilon)\}\}\}$$

3rd rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in C_\varepsilon(A)\}\}\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in B\}\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in C_\varepsilon(A)\} = B$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \textcolor{red}{C}\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in B\} = C$$

Example (Unfolding of the multistep function $\widehat{\Delta}_N$)

Let $x = baa$ over the alphabet $\Sigma = \{a, b\}$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, ba)\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \widehat{\Delta}_N(A, b)\}\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in \widehat{\Delta}_N(A, \varepsilon)\}\}\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in C_\varepsilon(A)\}\}\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in \bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in B\}\}$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in C\}$$

D

1st rec. call

2nd rec. call

3rd rec. call

$$\bigcup \{C_\varepsilon(\Delta(q, b)) \mid q \in C_\varepsilon(A)\} = B$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in B\} = C$$

$$\bigcup \{C_\varepsilon(\Delta(q, a)) \mid q \in C\} = D$$

Lemma

$C_\varepsilon(A)$ is least extension of A that is closed under ε -transitions:

$$q \in C_\varepsilon(A) \implies \Delta_{N_\varepsilon}(q, \varepsilon) \subseteq C_\varepsilon(A)$$

Lemma

$C_\varepsilon(A)$ is least extension of A that is closed under ε -transitions:

$$q \in C_\varepsilon(A) \implies \Delta_{N_\varepsilon}(q, \varepsilon) \subseteq C_\varepsilon(A)$$

Theorem

every set accepted by NFA_ε is regular

Lemma

$C_\varepsilon(A)$ is least extension of A that is closed under ε -transitions:

$$q \in C_\varepsilon(A) \implies \Delta_{N_\varepsilon}(q, \varepsilon) \subseteq C_\varepsilon(A)$$

Theorem

every set accepted by NFA_ε is regular

Proof. (by construction)

- $NFA_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$

Lemma

$C_\varepsilon(A)$ is least extension of A that is closed under ε -transitions:

$$q \in C_\varepsilon(A) \implies \Delta_{N_\varepsilon}(q, \varepsilon) \subseteq C_\varepsilon(A)$$

Theorem

every set accepted by NFA_ε is regular

Proof. (by construction)

- $NFA_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$
- $L(N_1) = L(N_2)$ for $NFA N_2 = (Q, \Sigma, \Delta_2, S, F_2)$ with

Lemma

$C_\varepsilon(A)$ is least extension of A that is closed under ε -transitions:

$$q \in C_\varepsilon(A) \implies \Delta_{N_\varepsilon}(q, \varepsilon) \subseteq C_\varepsilon(A)$$

Theorem

every set accepted by NFA_ε is regular

Proof. (by construction)

- $NFA_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$
- $L(N_1) = L(N_2)$ for $NFA N_2 = (Q, \Sigma, \Delta_2, S, F_2)$ with
 - ① $\Delta_2(q, a) := \widehat{\Delta_1}(\{q\}, a) \quad \forall q \in Q \ \forall a \in \Sigma$

Lemma

$C_\varepsilon(A)$ is least extension of A that is closed under ε -transitions:

$$q \in C_\varepsilon(A) \implies \Delta_{N_\varepsilon}(q, \varepsilon) \subseteq C_\varepsilon(A)$$


Theorem

every set accepted by NFA_ε is regular

Proof. (by construction)

- $NFA_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$
- $L(N_1) = L(N_2)$ for $NFA N_2 = (Q, \Sigma, \Delta_2, S, F_2)$ with
 - ① $\Delta_2(q, a) := \widehat{\Delta_1}(\{q\}, a) \quad \forall q \in Q \ \forall a \in \Sigma$
 - ② $F_2 := \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$


Example



Example

$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	$\{1\}$	\emptyset	\emptyset	$\{2\}$
2	\emptyset	$\{2\}$	\emptyset	$\{3\}$
3	\emptyset	\emptyset	$\{3\}$	\emptyset




Example

$\text{NFA}_\epsilon N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	$\{1\}$	\emptyset	\emptyset	$\{2\}$
2	\emptyset	$\{2\}$	\emptyset	$\{3\}$
3	\emptyset	\emptyset	$\{3\}$	\emptyset

$\text{NFA } N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\epsilon(\{q\}) \cap F_1 \neq \emptyset\}$



Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c



Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c



$$\Delta_2(1, a) = \widehat{\Delta}_1(\{1\}, a) = \bigcup \{C_\varepsilon(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \varepsilon)\}$$

Example


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
------------	-----	-----	-----



$$\begin{aligned}
 \Delta_2(1, a) &= \widehat{\Delta}_1(\{1\}, a) = \bigcup \{C_{\epsilon}(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\} \\
 &= C_{\epsilon}(\Delta_1(1, a)) \cup C_{\epsilon}(\Delta_1(2, a)) \cup C_{\epsilon}(\Delta_1(3, a))
 \end{aligned}$$

Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
------------	-----	-----	-----



$$\begin{aligned}
 \Delta_2(1, a) &= \widehat{\Delta}_1(\{1\}, a) = \bigcup \{C_\varepsilon(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \varepsilon)\} \\
 &= C_\varepsilon(\Delta_1(1, a)) \cup C_\varepsilon(\Delta_1(2, a)) \cup C_\varepsilon(\Delta_1(3, a)) \\
 &= C_\varepsilon(\{1\}) \cup C_\varepsilon(\emptyset) \cup C_\varepsilon(\emptyset)
 \end{aligned}$$

Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
1	{1, 2, 3}		



$$\begin{aligned}
 \Delta_2(1, a) &= \widehat{\Delta}_1(\{1\}, a) = \bigcup \{C_\varepsilon(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \varepsilon)\} \\
 &= C_\varepsilon(\Delta_1(1, a)) \cup C_\varepsilon(\Delta_1(2, a)) \cup C_\varepsilon(\Delta_1(3, a)) \\
 &= C_\varepsilon(\{1\}) \cup C_\varepsilon(\emptyset) \cup C_\varepsilon(\emptyset) \\
 &= \{1, 2, 3\}
 \end{aligned}$$

Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
1	{1, 2, 3}		



Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
1	{1, 2, 3}		



$$\Delta_2(1, b) = \widehat{\Delta}_1(\{1\}, b) = \bigcup \{C_\varepsilon(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \varepsilon)\}$$

Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
1	{1, 2, 3}		



$$\begin{aligned}
 \Delta_2(1, b) &= \widehat{\Delta}_1(\{1\}, b) = \bigcup \{C_\varepsilon(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \varepsilon)\} \\
 &= C_\varepsilon(\Delta_1(1, b)) \cup C_\varepsilon(\Delta_1(2, b)) \cup C_\varepsilon(\Delta_1(3, b))
 \end{aligned}$$

Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
1	{1, 2, 3}		



$$\begin{aligned}
 \Delta_2(1, b) &= \widehat{\Delta}_1(\{1\}, b) = \bigcup \{C_\varepsilon(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \varepsilon)\} \\
 &= C_\varepsilon(\Delta_1(1, b)) \cup C_\varepsilon(\Delta_1(2, b)) \cup C_\varepsilon(\Delta_1(3, b)) \\
 &= C_\varepsilon(\emptyset) \cup C_\varepsilon(\{2\}) \cup C_\varepsilon(\emptyset)
 \end{aligned}$$

Example


NFA_ε $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	∅	∅	{2}
2	∅	{2}	∅	{3}
3	∅	∅	{3}	∅

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
1	{1, 2, 3}	{2, 3}	




$$\begin{aligned}
 \Delta_2(1, b) &= \widehat{\Delta}_1(\{1\}, b) = \bigcup \{C_\varepsilon(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \varepsilon)\} \\
 &= C_\varepsilon(\Delta_1(1, b)) \cup C_\varepsilon(\Delta_1(2, b)) \cup C_\varepsilon(\Delta_1(3, b)) \\
 &= C_\varepsilon(\emptyset) \cup C_\varepsilon(\{2\}) \cup C_\varepsilon(\emptyset) \\
 &= \{2, 3\}
 \end{aligned}$$

Example (cont'd)

$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with


Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset



$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ_2	a	b	c
1	{1, 2, 3}	{2, 3}	{3}
2	\emptyset	{2, 3}	{3}
3	\emptyset	\emptyset	{3}



Outline

- 1 A Quick Recap
- 2 Nondeterministic Finite Automata
- 3 Epsilon Transitions
- 4 Closure Properties

Theorem

regular sets are effectively closed under **concatenation**

Theorem

regular sets are effectively closed under **concatenation**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
 $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$

Theorem

regular sets are effectively closed under **concatenation**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- without loss of generality $Q_1 \cap Q_2 = \emptyset$

Theorem

regular sets are effectively closed under **concatenation**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
 $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- without loss of generality $Q_1 \cap Q_2 = \emptyset$
- $AB = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S_1, F_2)$ with

Theorem

regular sets are effectively closed under **concatenation**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
 $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- without loss of generality $Q_1 \cap Q_2 = \emptyset$
- $AB = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S_1, F_2)$ with
 $\textcircled{1} Q \quad \quad \quad := \quad Q_1 \cup Q_2$

Theorem

regular sets are effectively closed under **concatenation**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- without loss of generality $Q_1 \cap Q_2 = \emptyset$
- $AB = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S_1, F_2)$ with

$$\begin{aligned} \textcircled{1} Q &:= Q_1 \cup Q_2 \\ \textcircled{2} \Delta(q, a) &:= \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \end{cases} \end{aligned}$$

Theorem

regular sets are effectively closed under **concatenation**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
 $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- without loss of generality $Q_1 \cap Q_2 = \emptyset$
- $AB = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S_1, F_2)$ with
 - 1 $Q := Q_1 \cup Q_2$
 - 2 $\Delta(q, a) := \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ \Delta_2(q, a) & \text{if } q \in Q_2 \text{ and } a \in \Sigma \end{cases}$

Theorem

regular sets are effectively closed under **concatenation**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
 $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- without loss of generality $Q_1 \cap Q_2 = \emptyset$
- $AB = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S_1, F_2)$ with

$$\begin{aligned} \textcircled{1} Q &:= Q_1 \cup Q_2 \\ \textcircled{2} \Delta(q, a) &:= \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ \Delta_2(q, a) & \text{if } q \in Q_2 \text{ and } a \in \Sigma \\ S_2 & \text{if } q \in F_1 \text{ and } a = \epsilon \end{cases} \end{aligned}$$

Theorem


regular sets are effectively closed under **concatenation**

Proof. (by construction)


- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
 $B = L(N_2)$ for NFA $N_2 = (Q_2, \Sigma, \Delta_2, S_2, F_2)$
- without loss of generality $Q_1 \cap Q_2 = \emptyset$
- $AB = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S_1, F_2)$ with
 - 1 $Q := Q_1 \cup Q_2$
 - 2 $\Delta(q, a) := \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ \Delta_2(q, a) & \text{if } q \in Q_2 \text{ and } a \in \Sigma \\ S_2 & \text{if } q \in F_1 \text{ and } a = \epsilon \\ \emptyset & \text{otherwise} \end{cases}$

Example


$\{x \in \{a\}^* \mid |x| \text{ is divisible by } 3\}$




$\{x \in \{a\}^* \mid |x| \text{ is divisible by } 4\}$



Example



Example



Theorem

regular sets are effectively closed under **asterate**

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with
 - 1 $Q := Q_1 \cup \{s\}$

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with
 - ① $Q := Q_1 \cup \{s\}$
 - ② $S := \{s\}$

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with

$$\textcircled{1} Q \quad := \quad Q_1 \cup \{s\}$$

$$\textcircled{2} S \quad := \quad \{s\}$$

$$\textcircled{3} F \quad := \quad \{s\}$$

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with
 - ① $Q := Q_1 \cup \{s\}$
 - ② $S := \{s\}$
 - ③ $F := \{s\}$
 - ④ $\Delta(q, a) := \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \end{cases}$

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with
 - ① $Q := Q_1 \cup \{s\}$
 - ② $S := \{s\}$
 - ③ $F := \{s\}$
 - ④ $\Delta(q, a) := \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ S_1 & \text{if } q = s \text{ and } a = \epsilon \end{cases}$

Theorem

regular sets are effectively closed under **asterate**

Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with
 - ① $Q := Q_1 \cup \{s\}$
 - ② $S := \{s\}$
 - ③ $F := \{s\}$
 - ④ $\Delta(q, a) := \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ S_1 & \text{if } q = s \text{ and } a = \epsilon \\ S & \text{if } q \in F_1 \text{ and } a = \epsilon \end{cases}$


Theorem

regular sets are effectively closed under **asterate**


Proof. (by construction)

- $A = L(N_1)$ for NFA $N_1 = (Q_1, \Sigma, \Delta_1, S_1, F_1)$
- $A^* = L(N)$ for NFA _{ϵ} $N = (Q, \Sigma, \epsilon, \Delta, S, F)$ with
 - ① $Q := Q_1 \cup \{s\}$
 - ② $S := \{s\}$
 - ③ $F := \{s\}$
 - ④ $\Delta(q, a) := \begin{cases} \Delta_1(q, a) & \text{if } q \in Q_1 \text{ and } a \in \Sigma \\ S_1 & \text{if } q = s \text{ and } a = \epsilon \\ S & \text{if } q \in F_1 \text{ and } a = \epsilon \\ \emptyset & \text{otherwise} \end{cases}$

Example


 $\{a\}^* \{b\}^*$ 

Example

 $\{a\}^* \{b\}^*$ 

Example

$$(\{a\}^* \{b\}^*)^* = \{a, b\}^*$$



Thanks! & Questions?